

Chapter 8

Measurements of Ice Nucleating Particles and Ice Residuals

DANIEL J. CZICZO,^{a,b} LUIS LADINO,^c YVONNE BOOSE,^d ZAMIN A. KANJI,^d
PIOTR KUPISZEWSKI,^e SARA LANCE,^f STEPHAN MERTES,^g AND HEIKE WEX^g

^a *Department of Earth, Atmospheric and Planetary Sciences, Massachusetts
Institute of Technology, Cambridge, Massachusetts*

^b *Department of Civil and Environmental Engineering, Massachusetts
Institute of Technology, Cambridge, Massachusetts*

^c *Centro de Ciencias de la Atmósfera, Universidad Nacional Autónoma de México,
Ciudad Universitaria, Mexico City, Mexico*

^d *Institute for Atmospheric and Climate Science, ETH Zurich, Zurich, Switzerland*
^e *Paul Scherrer Institute, Villigen, Switzerland*

^f *University at Albany, State University of New York, Albany, New York*

^g *Leibniz Institute for Tropospheric Research, Leipzig, Germany*

ABSTRACT

It has been known that aerosol particles act as nuclei for ice formation for over a century and a half (see Dufour). Initial attempts to understand the nature of these ice nucleating particles were optical and electron microscope inspection of inclusions at the center of a crystal (see Isono; Kumai). Only within the last few decades has instrumentation to extract ice crystals from clouds and analyze the residual material after sublimation of condensed-phase water been available (see Cziczo and Froyd). Techniques to ascertain the ice nucleating potential of atmospheric aerosols have only been in place for a similar amount of time (see DeMott et al.). In this chapter the history of measurements of ice nucleating particles, both in the field and complementary studies in the laboratory, are reviewed. Remaining uncertainties and artifacts associated with measurements are described and suggestions for future areas of improvement are made.

1. A history of ice nucleating particle measurements

a. The difference between ice nucleating particles and ice residuals

The idea that small particles that originate from Earth's surface initiate ice formation in the atmosphere was suggested by Dufour (1862), who found that hailstones contained small sand, ash, or chaff. It is now understood that these inclusions often initiate nucleation of the ice phase either via deposition of water vapor to their surface or from within a supercooled liquid water droplet; they are termed ice nucleating particles (INPs; all acronyms additionally defined in Table 8-1) (Vali et al. 2015). After ice formation, riming, scavenging, and aggregation may lead to the collection of additional particles by a growing ice crystal and

chemical reactions may take place on the ice crystals' surface. Dissolution of the original INP and ice multiplication processes can form ice particles that do not contain an INP. Particles remaining after evaporation of ice in hydrometeors are called ice residuals (IRs). To date, attempts to compare ice number density to residuals have proven difficult because of shattering of ice crystals on cloud probes (Korolev and Isaac 2005; Cziczo and Froyd 2014). For these reasons, IRs and INPs cannot currently be assumed to be identical.

Nevertheless, atmospheric IRs have been studied to understand which particles initiated the formation of ice crystals. One technique is to examine particles located at the crystallographic center of ice crystals by means of microscopy (Rogers and Vali 1978). Newly formed ice crystals likely contain only the INP as they have not been

Corresponding author e-mail: Daniel J. Cziczo, djciczo@mit.edu

Publisher's Note: This chapter was revised on 12 April 2017 to include acknowledgments originally missing in the Acknowledgments section.

DOI: 10.1175/AMSMONOGRAPHS-D-16-0008.1

© 2017 American Meteorological Society. For information regarding reuse of this content and general copyright information, consult the [AMS Copyright Policy](https://www.ametsoc.org/PUBSReuseLicenses) (www.ametsoc.org/PUBSReuseLicenses).

TABLE 8-1. List of acronyms.

Expansion	Acronym
Aerosol interaction and dynamics in the atmosphere	AIDA
Attenuated total reflectance FTIR	ATR-FTIR
Bielefeld ice nucleation array	BINARY
Black carbon	BC
Cloud condensation nuclei	CCN
Continuous flow diffusion chamber	CFDC
Colorado State University-immersion mode ice spectrometer	CSU-IS
Contact ice nucleation chamber	CINC
Cloud and Aerosol Characterization Experiment	CLACE
Collision ice nucleation chamber	CLINCH
Cold stage	CS
Counterflow virtual impactor	CVI
Droplet evaporation unit	DEU
Drop freezing arrays	DFA
Differential scanning calorimeter	DSC
Electrodynamic balance	EDB
Energy-dispersive X-ray spectroscopy	EDX
Electron microscopy	EM
Environmental SEM	ESEM
Field emission gun	FEG-SEM
Fast Fourier transformation	FFT
Fast ice nucleation chamber	FINCH
Frankfurt Ice Deposition Freezing Experiment	FRIDGE
Fourier transform infrared spectroscopy	FTIR
Immersion mode cooling chamber	IMCA
Ice nucleation active site densities	INAS
Ice nucleating particles	INP
Ice residuals	IR
Ice selective inlet	ISI
Leipzig aerosol cloud interaction simulator	LACIS
Limit of detection	LOD
Manchester ice cloud chamber	MICC
Massachusetts Institute of Technology-collision efficiency chamber	MIT-CEC
Mixed-phase cloud	MPC
Microlitre nucleation by immersed particle instrument	μ l-NIPI
Near-edge absorption fine structure spectroscopy	NEXAFS
North Carolina	NC
Optical trap	OT
Pumped counterflow virtual impactor	PCVI
Portable immersion mode cooling chamber	PIMCA
Portable ice nucleation chamber	PINC
Scanning EM	SEM
Spectrometer ice nuclei	SPIN
Single-particle mass spectrometer	SPMS
Scanning transmission X-ray microscopy	STXM
Transmission (or tunneling) EM	TEM
Temperature gradient diffusion chamber	TGDC
Time of flight	ToF
Ultraviolet	UV
Vacuum chamber	VC
Wind tunnel	WT
X-ray diffraction	XRD
Zurich ice nucleation chamber	ZINC

exposed to extensive processing. They have been sampled using counterflow virtual impactors (CVIs) from aircraft to study cirrus IRs (e.g., [Ogren et al. 1985](#); [Noone et al. 1988](#); [Twohy et al. 1997](#); [Cziczo et al. 2004](#)) or in special ground-based inlets to study IRs from mixed-phase clouds (MPCs) ([Mertes et al. 2007](#); [Kupiszewski et al. 2015](#)).

Sampling INPs follows a different approach whereby particles, either airborne or on a substrate, are exposed to known thermodynamic conditions and the occurrence of ice formation is monitored. Various techniques have been developed since the first half of the twentieth century to study the different ice formation mechanisms ([Kanji et al. 2017](#), chapter 1). An overview of the most important techniques and results is also given here.

b. Ground-based studies

For ground-based examinations there are three main approaches: investigating collected precipitation (IRs), studying the formation of ice on ambient particles (INPs), and examining MPCs directly (IRs).

1) MEASUREMENTS OF INPs AND IRs IN COLLECTED PRECIPITATION

[Isono \(1955\)](#) in Japan and [Kumai \(1961\)](#) in the United States analyzed the particles found in collected ice crystals by means of electron microscopy (EM) and both found clay minerals to be the main IRs. Although mineral material is still considered the dominant INP type by number, in the following years biological IRs were also detected ([Szyrmer and Zawadzki 1997](#) and references therein). The use of drop freezing arrays (DFA) by [Bigg \(1953\)](#) was an early technique that allowed determination of freezing onset temperature and frozen fractions. [Vali \(1966\)](#) estimated atmospheric INP concentrations, by the definition here assuming INP number concentration was equivalent to that measured for IRs, based on volume of melted hail and rainwater from initial ice-phase precipitation.

2) AMBIENT AEROSOL PARTICLES AS INPs

To determine the INP concentration, two methods have been used: online (such as cloud or continuous flow chambers) and offline (where particles are collected on a substrate and subsequently exposed to controlled conditions).

INPs were initially studied in the laboratory starting as early as the mid-1940s, when [Vonnegut \(1947\)](#), motivated by cloud seeding, not atmospheric INPs, sought to identify and characterize the activity of potent INPs. Vonnegut used an early cloud chamber to show that a man-made INP, silver iodide, nucleated ice between -4° and -8° C. The main mechanisms for chambers being used at that time were either expansion (i.e., lowering the pressure) or on injection of hot steam in a cooled chamber (i.e.,




FIG. 8-1. The range of INP concentration observed for ambient aerosols vs temperature. The common temperature ranges of CFDC (Table 8-2) and cold-stage (Table 8-3) instruments are given for context (DeMott et al. 2011 and references therein).

supersaturation). Online methods were also developed as early as the 1940s for in situ aircraft measurements of INPs (e.g., Palmer 1949; Langer et al. 1967). With secondary ice formation and crystal shatter artifacts not yet known, INP and ice crystal concentrations were often assumed to be identical. This changed when Mossop (1970) and Hallett and Mossop (1974) first suggested secondary ice production mechanisms in clouds and when Korolev and Isaac (2005) quantified ice shattering on cloud probes.

Today, the most common in situ and laboratory chambers are variants of the continuous flow diffusion chamber (CFDC; Rogers 1988), which uses supersaturation derived from two ice-coated plates held at different temperatures (section 2a). Several larger, nonportable systems are currently in use for laboratory studies of specific materials. They are based on the expansion principle (e.g., Möhler et al. 2003; Connolly et al. 2012), on heat-vapor diffusion (Hartmann et al. 2011), or mixing of cold and dry air with warm and humidified air (Bundke et al. 2008) (see Kanji et al. 2017, chapter 1, and section 2).

Offline methods for INP determination were initially derived from aerosol filter collections. Subsequent analysis is performed by exposing the filter to controlled supersaturation and temperature and observing ice formation. This is either done in deposition/condensation mode by controlling the temperature and humidity above the filter or for aerosol immersed in water droplets prior to cooling (e.g., Schnell 1977; Rosinski et al. 1986).

DeMott et al. (2015) found good agreement when comparing INP concentrations derived by online and offline methods. As a whole, measurements have demonstrated an exponential increase in INP concentration with decreasing temperatures (Fig. 8-1; see additional description in Kanji et al. 2017, chapter 1). With

variability up to seven orders of magnitude (DeMott et al. 2010, 2016; Petters and Wright 2015), it remains to be determined what fraction of this is due to natural versus instrumental differences (DeMott et al. 2011).

Historically, laboratory measurements were sometimes motivated by determination of ice activity of various samples without considerable connection to their atmospheric abundance. Foremost among these were man-made substances such as silver iodide that were studied for their use in weather modification (Vonnegut 1947). Natural samples such as soil dusts, leaf litter, seawater, and bacterial samples were also studied, often motivated by collections of fog water and precipitation (e.g., Schnell and Vali 1976; Vali et al. 1976; Schnell 1977).

3) IRS IN MIXED-PHASE CLOUDS

There exist two ground-based CVI inlet systems for sampling IRs of ice crystals contained in MPCs. These are the Ice-CVI (Mertes et al. 2007) and Ice Selective Inlet (ISI; Kupiszewski et al. 2015), both of which have been deployed at the high-altitude research station Jungfraujoch in the Swiss Alps (3580 m MSL; section 2b).

The Ice-CVI was operated during the Cloud and Aerosol Characterization Experiment (CLACE) for which Mertes et al. (2007) presented an ice activated fraction as a function of particle size. Assuming the IRs to be the original INPs, they concluded that the particles' efficiency to nucleate ice increases with size. A higher concentration of IRs with diameters below $1\ \mu\text{m}$ was found than those in the supermicrometer range. A higher abundance of submicrometer-diameter IRs was also found by Kupiszewski et al. (2016) although these findings are not in agreement with the finding of larger, often supermicrometer, particles by Cziczo et al. (2013) and Mason et al. (2016).

Several studies have been carried out to determine IR chemical composition. It should be noted that these are location-specific findings and, because of the limited number of measurements to date, these should not yet be taken as general findings. An enrichment of black carbon (BC) mass in IRs was observed using a filter-based light transmission method (CLACE 3; Cozic et al. 2008; Mertes et al. 2007) and using EM (CLACE 5; Ebert et al. 2011). Although the signals from ice nucleation and scavenging were not directly differentiated, the size limit of the ice crystals sampled ($<20\ \mu\text{m}$ diameter) was noted to allow minimal scavenging time after nucleation. In contrast, no enrichment of BC in IRs was found for CLACE 6 (Kamphus et al. 2010) and CLACE 2013 (Kupiszewski et al. 2016; Worringer et al. 2015) using single-particle mass spectrometry, laser-induced incandescence measurements, and EM. Another chemical fingerprint in the Ice-CVI measurements was the detection of lead in the IRs, which was substantially enhanced, compared to the

background aerosol population (Cziczo et al. 2009; Ebert et al. 2011; Worringen et al. 2015) and is known to be an efficient INP type (Cziczo et al. 2009). In general, the aforementioned studies identified mineral dust (silicates) as the dominant IR, and presumably INP, component. The existence of biological particles in the IRs was investigated by Kupiszewski et al. (2015) who found an enrichment of fluorescent particles compared to the background aerosol during a Saharan dust event in the sub-2- μm IRs.

c. Aircraft-based studies

The first airborne efforts to understand the nature of ice nucleation in clouds was performed in the early 1990s using a CVI to separate ice crystals from other particles (Noone et al. 1992; Ström and Heintzenberg 1994; Heintzenberg et al. 1996; Ström et al. 1997; Seifert et al. 2003, 2004). These initial studies sublimated condensed-phase water and performed counts, and later sizing, of the residual particles. As described in section 3b, it is highly likely that these initial efforts were subject to undetected artifacts associated with ice impaction on aircraft and inlet surfaces.

Later studies collected particles for offline analysis, for example, by EM (Petzold et al. 1998; Twohy and Gandrud 1998; Twohy and Poellot 2005; Targino et al. 2006; Cziczo et al. 2013), and these are more fully described in section 2c. Since the early 2000s, the most common technique is the online, real-time mass spectrometer described in section 2d. (Cziczo et al. 2004; Pratt et al. 2009; Froyd et al. 2010; Cziczo et al. 2013).

The other means by which INPs can be studied from aircraft platforms (or at high-altitude sites) is to induce nucleation in a portable CFDC (section 2); the concept is to use controlled saturation and temperature to induce ice formation on a subset of the ambient aerosol (DeMott et al. 2010). When the chambers are utilized in free-tropospheric air masses, the assumption is the particles studied are relevant to cirrus formation (DeMott et al. 2003). One particular example is the work of Chen et al. (1998) who used an ice chamber to induce ice formation on upper-tropospheric aerosol as well as contrail-affected air masses. Ice crystals were separated and sublimated with the residuals collected for offline analysis using EM.

2. Measurements methods

a. Ice nucleation chambers

No single chamber has been developed to investigate all ice nucleation mechanisms or the full terrestrial temperature range; chambers are therefore designed to study specific ice nucleation mechanisms and temperature ranges (DeMott et al. 2011). As shown in Tables 8-2 and 8-3, these chambers tend to fall into three broad categories: 1) large chambers in fixed laboratories that accept collected or prepared samples,

2) offline chambers that expose collected or prepared samples to low temperature and ice saturated conditions, and 3) portable chambers for both field and laboratory use.

The main purpose of a cloud chamber is to provide a controlled water vapor content and temperature at which particles can nucleate ice. Aerosol size, concentration, residence time, and composition are then varied so that the ice nucleation onset or frozen fractions can be determined. Note that in some instruments care is taken to be at supersaturated conditions continuously while the temperature is changed [e.g., Leipzig aerosol cloud interaction simulator (LACIS), immersion mode cooling chamber (IMCA); note acronyms defined in Tables 8-1–8-3] while in others the temperature is constant and RH is varied (e.g., most CFDCs and some cold-stage instruments). A subset of chambers, for example, those that use expansion of the air parcel, allow both parameters to be varied at the same time [e.g., aerosol interaction and dynamics in the atmosphere (AIDA)].

For the laboratory chambers, particles are typically directly injected into the cell, with the exception of vacuum chamber (VC) and cold stage (CS), where particles are deposited on a substrate. Droplets with particles immersed are kept within the instruments during immersion freezing experiments conducted in a differential scanning calorimeter (DSC), wind tunnel (WT), optical trap (OT), or electrodynamic balance (EDB). In contrast, in continuous flow instruments [such the LACIS, Zurich ice nucleation chamber (ZINC), portable ice nucleation chamber (PINC), IMCA, portable immersion mode cooling chamber (PIMCA), contact ice nucleation chamber (CINC), Manchester ice cloud chamber (MICC), spectrometer ice nuclei (SPIN), collision ice nucleation chamber (CLINCH), and Massachusetts Institute of Technology–collision efficiency chamber (MIT-CEC) or the WT, EDB, DSC, and OT] the particles pass through the instruments during the experiments. Since particle size (surface area) has been shown to play a role in ice activity there has been a shift to monodisperse studies (i.e., use of mobility selection before the chamber).

The primary goal of the detection system of most cloud chambers is quantification of ice crystals. A secondary goal, discussed in the next sections, is often compositional analysis. To quantify ice crystal number, most instruments use an optical system to either directly capture images of ice crystals or determine number by means of incidents of light scattering from a continuous laser beam at the terminus of the chamber. More sophisticated optical systems that use laser depolarization to distinguish between ice crystals, liquid droplets, and unactivated aerosol particles are now also in use.

b. Separation techniques

The measurement of IRs involves the use of a CVI, first described by Ogren et al. (1985) for aircraft

TABLE 8-2. Examples of laboratory-based contemporary instrumentation to study ice nucleation. Homogeneous freezing (Ho), deposition nucleation (DE), immersion freezing (IM), condensation freezing (CD), contact freezing (CT).

Type of instrument	Abbreviation	Full name	Freezing mode	Type of analysis	References
Chamber	AIDA	Aerosol interaction and dynamics in the atmosphere	Ho, DE, IM, CD	Online	Möhler et al. (2003)
	ZINC	Zurich ice nucleation chamber	Ho, DE, CD, IM	Online	Stetzer et al. (2008)
	LACIS	Leipzig aerosol cloud interaction simulator	Ho, IM	Online	Hartmann et al. (2011)
	MICC	Manchester ice cloud chamber	Ho, DE, IM	Online	Connolly et al. (2012)
	IMCA	Immersion mode cooling chamber	Ho, IM	Online	Lüönd et al. (2010)
	CLINCH	Collision ice nucleation chamber	Ho, CT	Online	Ladino et al. (2011)
	CINC	Contact ice nucleation chamber	Ho, CT	Online	Niehaus et al. (2014)
	MIT-CEC	Massachusetts Institute of Technology–collision efficiency chamber	Ho, CT	Online	Ardon-Dryer et al. (2015)
Cell	WT	Wind tunnel	Ho, DE, IM, CD, CT	Online	Diehl et al. (2011)
	EDB	Electrodynamic balance	Ho, IM, CT	Online	Davis (1997); Hoffmann et al. (2013a,b)
	VC	Vacuum chamber	Ho, DE	Online	Shilling et al. (2006)
	DSC	Differential scanning calorimeter	Ho, IM	Online	Zuberi et al. (2001)
	μ l-NIPI	Microlitre nucleation by immersed particle instrument	Ho, IM	Online	Whale et al. (2015)
	CSU-IS	Colorado State University–immersion mode ice spectrometer	Ho, IM	Online	Hill et al. (2014)
	NC State-CS	Cold-stage-supported droplet assay	Ho, IM	Online	Wright and Petters (2013)
	CS	Cold stage	Ho, DE, CD, IM	Online and Offline	Koop et al. (1998); Durant and Shaw (2005)
	BINARY	Bielefeld ice nucleation array	Ho, IM	Online	Budke and Koop (2015)
	OT	Optical trap	Ho, CT	Online	Davis et al. (2015)

applications, where a controlled airflow is blown out of the CVI inlet tip. Smaller interstitial particles are decelerated, stopped, and rejected, whereas larger cloud particles can overcome the counterflow and are impacted into a dry, particle-free flow. The condensed phase of the collected cloud particles is evaporated, releasing the residuals. The CVI separation is only based on inertia and does not differentiate between droplets and ice.

1) AIRCRAFT-BASED INLETS

The deployment of aircraft CVI inlets in cirrus has been used to separate IRs from ambient aerosol because of the absence of droplets in these completely glaciated clouds (e.g., Cziczo et al. 2004, 2013; Froyd et al. 2010; Heintzenberg et al. 1996; Seifert et al. 2004; Stith et al. 2011; Twohy and Gandrud 1998; Twohy and Poellot 2005). The lower cut size of the transmitted particles for a given CVI design depends on the aircraft velocity and the magnitude of the counterflow, with typical cut sizes on the order of a few micrometers aerodynamic diameter (Anderson et al. 1993; Noone et al. 1988; Twohy et al. 2003). Much less discussed in the literature is the upper cutoff diameter of a

CVI. In general, because of geometrical and flow rate limitations, only crystals up to a size of $50\ \mu\text{m}$ can be stopped and sublimated without impaction to a wall surface inside traditional CVI inlets (Cziczo and Froyd 2014). As Cziczo and Froyd (2014) point out, cirrus and ice clouds often contain crystals larger than $50\ \mu\text{m}$, which means that only the small size fraction has, to date, been sampled. Moreover, the larger crystals that cannot be stopped impact onto internal surfaces; creating artifacts (see section 3b). A further artifact, which is also relevant for ground-based CVI inlets, is the unintentional transmission of interstitial particles below the CVI cut size due to collision, coalescence, or riding the wake of ice particles (Pekour and Cziczo 2011).

2) GROUND-BASED INLETS

Because of the lower altitudes of mountain sites, ground-based inlets are typically operated in MPC conditions and therefore face the additional challenge of cloud phase separation. Since cloud droplets are of a similar aerodynamic size as small crystals, a traditional CVI is insufficient. As described in section 1b, two CVI-based separation techniques (Ice-CVI and ISI) have

TABLE 8-3. Examples of portable instruments (field and laboratory) to study ice nucleation. N_{CCN} is the number concentration of ambient CCN at the sampling location.

Type of instrument	Abbreviation	Full name	Freezing mode	Type of analysis	INP sensitivity (L^{-1})	References
Chamber	PINC	Portable ice nucleation chamber	Ho, DE, CD, IM	Online	0.1	Chou et al. (2011)
	PIMCA	Portable immersion mode cooling chamber	Ho, IM	Online	$0.1 * N_{\text{CCN}}/\text{L}$	Kohn et al. (2016)
	SPIN	Spectrometer for ice nuclei	Ho, DE, IM, CD	Online	1.0	Garimella et al. (2016)
	CSU-CFDC	Colorado State University–continuous flow diffusion chamber	Ho, DE, CD, IM	Online	0.01	Rogers (1988) ; Rogers et al. (2001)
	UT-CFDC	University of Toronto–continuous flow diffusion chamber	Ho, DE, CD	Online	0.01	Kanji and Abbatt (2009)
	FINCH	Fast ice nucleation chamber	IM, DE, CD	Online	0.01	Bundke et al. (2008)
	TGDC	Temperature gradient diffusion chamber	Ho, DE, CD	Online	0.01	Bailey and Hallett (2002)
	FRIDGE	Frankfurt Ice Deposition Freezing Experiment	IM, CD, DE	Offline	0.01	Bundke et al. (2008)
Cell	CS (DFT)	Mixing cloud chamber	DE	Offline	0.01	López and Ávila (2013)
		Cold stage	Ho, DE, CT, IM	Offline	0.01	Knopf et al. (2010) ; Mason et al. (2015a)
Cell	DFA	Drop freezing array	Ho, IM	Offline	1.0e^{-6}	Bigg (1953) ; Conen et al. (2012)

been developed for ground-based use. Both inlets use a series of additional components to extract small ($5\text{--}20\ \mu\text{m}$) crystals from MPCs.

The Ice-CVI is mounted vertically. Air is sampled through an omnidirectional inlet, which reduces transmission of cloud particles larger than $50\ \mu\text{m}$ ([Noone et al. 1992](#)). A virtual impactor downstream removes cloud particles larger than $20\ \mu\text{m}$. Phase is separated by a two-stage preimpactor (cutoff diameters 10 and $4\ \mu\text{m}$), where supercooled drops freeze on the impaction plates while crystals bounce off ([Tenberken-Pöttsch et al. 2000](#)). The flow is then directed into a conventional CVI ([Mertes et al. 2001](#)), where interstitial particles below $5\ \mu\text{m}$ are removed.

In the ISI droplets are removed from the sample flow. This is carried out via the Wegener–Bergeron–Findeisen process, in a so-called droplet evaporation unit (DEU), a chamber with ice-coated inner walls that induce droplet evaporation via the difference in saturation vapor pressures over water and ice. Directly upstream and downstream of the DEU, optical particle size spectrometers count and size cloud particles contained in the sample flow. The phase and habit of cloud particles transmitted through the DEU are ascertained using a particle phase discriminator ([Kaye et al. 2008](#); [Vochezer et al. 2016](#)) and interstitial particles are removed from the sample flow using a pumped counter-flow virtual impactor (PCVI; [Boulter et al. 2006](#)), with a cutoff size of $\sim 5\ \mu\text{m}$ ([Kupiszewski et al. 2015](#)).

Both the Ice-CVI and ISI inlet systems have limitations and may suffer from sampling artifacts. The two systems sample through omnidirectional inlets on which crystals could shatter; the released fragments could then make it through the system. The use of impaction plates in the Ice-CVI could produce additional shatter and potential abrasion of material from the plate surfaces, although the kinetic energy of ice under these conditions has been shown to be low ([Vidaurre and Hallett 2009](#)). Comparisons of Ice-CVI IR and small ice particle concentrations are similar and suggest minimal artifacts ([Kupiszewski et al. 2016](#); [Mertes et al. 2007](#)). The ISI is known to partially evaporate ice crystals within the inlet, thereby leading to rejection of some IRs ([Kupiszewski et al. 2015](#)). Another limitation of both inlet systems is the inability to distinguish between primary and secondary ice.

c. Analysis of INPs with microscopy and other optical techniques

INP surface topography, surface area, and other physical and chemical characteristics have been probed using optical and other microscopy techniques. In some cases the technique [e.g., scanning EM (SEM)] can be destructive and/or alter the particle. Less destructive methods can be used to probe the physical and chemical properties of potential INPs before or after subjecting those particles to freezing conditions. A homogeneous aerosol population can also be split between independent freezing and aerosol characterization experiments to




FIG. 8-2. Electron micrographs of ice crystals formed by deposition ice nucleation on a small fraction of mineral dust particles [(left) illite; (right) gypsum] within an ESEM (Zimmermann et al. 2007).

prevent experimental artifacts from influencing subsequent experiments. The techniques discussed in this section generally require that the particle is stationary, which is most frequently accomplished by depositing the particle onto a “cold stage” or an EM “grid” (Fig. 8-2).

Optical microscopy, sensitive to supermicrometer sizes, is typically used to determine when freezing has occurred. EM, with higher resolution, is typically used to determine surface area and topography (Hoffmann et al. 2013a). Limitations of SEM are exposure of the sample to vacuum (thereby potentially losing volatile material) and metallic coating required for samples with low electrical conductivity. Nevertheless, INPs of many different types have been analyzed in the laboratory using SEM (Alpert et al. 2011; Hiranuma et al. 2015a; Shilling et al. 2006; China et al. 2015). Cirrus IRs containing mineral dust, metallic particles, soot, and biological material have also been analyzed by SEM (Cziczo et al. 2013).

The field emission gun (FEG-SEM) technique can overcome problems resulting from low electrical conductivity of sample materials due to the smaller electron beam spot size used. Environmental SEM (ESEM) utilizes differential pumping of the region near the sample and the electron gun, and the higher pressure near the sample allows for electrical discharge while also preventing evaporation of volatile material. Ambient multicomponent aerosol, volcanic ash, and mineral dust INPs have been characterized using ESEM (Hoffmann et al. 2013a; Knopf et al. 2010; Wang et al. 2012a,b), as have IRs collected from orographic MPCs and cirrus (Ebert et al. 2011). By adding a cold-stage and carefully controlling humidity and temperature, ice nucleation experiments can take place within an ESEM chamber (Zimmermann et al. 2007).

Transmission (or tunneling) EM (TEM) has resolving power that can surpass SEM, and has been used to investigate very fine structures (~ 10 nm). This capability has been used to identify surface defects and morphology of

mineral dust and soot (Freedman 2015; Niedermeier et al. 2011; Sihvonen et al. 2014; China et al. 2015; Zolles et al. 2015) as well as internally mixed organic/inorganic INPs (Veghte et al. 2014). Phase and crystal lattice information at sub-10-nm can be obtained using fast Fourier transformation (FFT) analysis of high-resolution TEM images (Ebert et al. 2011). TEM has also been used to analyze cirrus IRs (Heintzenberg et al. 1996; Stith et al. 2011; Twohy and Poellot 2005; Twohy and Gandrud 1998).

Elemental composition and mixing state of INPs can be determined through energy-dispersive X-ray spectroscopy (EDX), which can be added to an EM instrument (Broadley et al. 2012; Ebert et al. 2011; Heintzenberg et al. 1996; Hiranuma et al. 2015a; Hoffmann et al. 2013b; Knopf et al. 2010; Stith et al. 2011; Twohy and Poellot 2005; Twohy and Gandrud 1998; Wang et al. 2012b; Zimmermann et al. 2007; Zolles et al. 2015).

Scanning transmission X-ray microscopy (STXM) near-edge absorption fine structure spectroscopy (NEXAFS) is used in the analysis of thin-film polymers and to provide chemical functionality relevant for biogenic and organic INPs (Knopf et al. 2010; Moffet et al. 2010; Wang et al. 2012a; Wilson et al. 2015).

Other tools to characterize INPs include Raman spectroscopy, fluorescence microscopy, Fourier transform infrared (FTIR) spectroscopy, attenuated total reflectance FTIR (ATR-FTIR), elastic X-ray diffraction (XRD), and powder XRD (Alpert et al. 2011; Baustian et al. 2013; Broadley et al. 2012; Brooks et al. 2014; Freedman 2015; Hiranuma et al. 2015b; Marcolli 2014; Mason et al. 2015b; Schill et al. 2014; Shilling et al. 2006; Sihvonen et al. 2014; Wise et al. 2012; Zolles et al. 2015).

d. Mass spectrometry of ice nucleating particles

Single-particle mass spectrometers (SPMSs) have been used since the early 1990s for the study of particles. Reviewed by Murphy (2007), the SPMS allows for the determination of single-particle chemical composition, typically of all aerosol components, both in situ and in real time. Modern SPMS instruments focus particles using an aerodynamic inlet and particles are detected and sized using scattered light as they pass through one or two laser beams set a fixed distance apart. Because of the optical detection step, SPMSs are limited to a lower size threshold, typically ~ 150 -nm geometric diameter particles that scatter sufficient light (Murphy 2007). It should be noted that this remains below the expected INP size mode (DeMott et al. 2010). A second laser, typically in the near UV, is fired to ablate the particle and ionize that material. Extracted ions are separated in single- or dual-polarity time-of-flight (ToF) mass spectrometers, thereby producing a complete mass spectrum on a single-particle basis. SPMS is generally considered

qualitative since signal is affected by the particle matrix and instrumental variability. Since its first airborne deployment for analysis of cirrus residuals in the 2001 CRYSTAL-FACE study by [Cziczo et al. \(2004\)](#), SPMS has become the most common technique for IRs analysis since ([Cziczo et al. 2004](#); [Pratt et al. 2009](#); [Froyd et al. 2010](#); [Cziczo et al. 2013](#)).

3. Uncertainties and limitations of contemporary measurements

a. Ice nucleation chambers

Most of the available portable online INP counters are limited in operation to temperatures relevant for MPC conditions ($> -40^{\circ}\text{C}$). Lower sampling temperatures (-40° to -60°C) would require chamber cold walls to be cooled to $\sim -60^{\circ}\text{C}$ and lower, which is challenging to achieve. The focus on warmer temperatures, albeit circumstantial, means there is a gap in knowledge about INP concentrations at cirrus conditions. On the warmer end, CFDCs are limited to sampling temperatures $< -8^{\circ}\text{C}$ since the warm wall temperatures would approach close to 0°C , resulting in melting of the ice layer.

CFDCs require human intervention for icing the chamber walls, a semiautomated procedure at best, limiting their use in continuous monitoring applications. Additionally, they have to be reiced after a few hours because of frost formation on the cold wall ([Rogers et al. 2001](#); [Chou et al. 2011](#)). This can be extended somewhat, to ~ 12 h, in horizontally oriented chambers ([Kanji and Abbatt 2009](#)).

CFDCs require laminar flow conditions, resulting in low sample flow rates (~ 1 lpm). The result of atmospheric measurements in remote areas at the higher-than-cirrus temperature range where CFDCs are most effective is that INP concentrations are often $< 1 \text{ L}^{-1}$. This then means longer sampling times are necessary for statistical significance. An exception is the fast ice nucleation chamber (FINCH), a turbulent mixing flow chamber that samples 5–10 lpm of ambient air ([Worringen et al. 2015](#); [Bundke et al. 2008](#)). Offline techniques that can use higher flow rates (> 20 lpm) to impact particles on substrates are also important for an increase of signal ([Mason et al. 2015a,b](#)).

CS and DFA techniques have the advantage of a low limit of detection (LOD) but require aerosol sampling and storage prior to ice nucleation analysis. It is not clear if effects of sample handling, storage, and impaction of particles onto substrates exist on the particle structure and morphology. Longer sampling times are typically necessary (> 5 h) for size-resolved collections onto filters ([Mason et al. 2015a](#); [Wang et al. 2012b](#)).

These limitations illustrate the need for higher instrument sensitivity in remote locations where the INP

concentration is low ($< 1 \text{ L}^{-1}$). [Boose et al. \(2016\)](#) showed that this requires accurate determination of signal to noise and LOD; reporting only INP concentrations $>$ LOD without reporting the frequency of measurements that fall below LOD can result in a significant overestimate of INP for a given location.

Another uncertainty is the determination of ice nucleation rate often achieved by using the residence time of particles in chambers. Since this time includes equilibration, nucleation, and growth, the nucleation rate is necessarily a lower limit ([Archuleta et al. 2005](#); [Kanji and Abbatt 2010](#)). This can be overcome in static chambers with high cooling rates if the time between ice nucleation and growth to detectable sizes is small relative to the time particles are exposed to given temperature and RH condition.

Although the EDB is a powerful technique to study different ice nucleation processes such as contact freezing or ice multiplication, and their experiments allow for long experimental time scales, it requires a large number of electrical charges on droplet surfaces for levitation ([Hoffmann et al. 2013a](#)). The effect of charge on the freezing process remains to be fully considered. For example, the excess or scarcity of electrical charges can substantially change the collection efficiencies between aerosol particles and cloud droplets, affecting the efficiency of contact freezing ([Ladino et al. 2013](#); [Ardon-Dryer et al. 2015](#)).

The use of polydisperse aerosol samples has inhibited the quantification of freezing efficiency in some instruments, such as the wind tunnel, EDB, or a dynamic CS ([Ladino et al. 2013](#)). Methods employing monodisperse particles also need to be considered carefully since they can lead to an overestimation of ice nucleation active site densities (INAS) if corrections for multiply charged particles are not applied ([Hartmann et al. 2016](#)). In general, measurements can be improved when accurate size distributions are acquired.

For CS methods in deposition and immersion nucleation, coagulation of smaller particles or the use of larger particles up to $10\text{-}\mu\text{m}$ may limit the application of the ice nucleation data generated ([Eastwood et al. 2008](#); [Kanji and Abbatt 2006](#)). While large aerosols have been shown to undergo long-range transport ([Young et al. 2016](#)), they may not bear atmospheric relevance for ice cloud formation at the upper limit of the investigated particle sizes since they are rare at cirrus altitudes.

Contact and immersion freezing studies conducted with EDB and WT experiments use droplet sizes from 60 to $350 \mu\text{m}$ ([Hoffmann et al. 2013a,b](#); [Diehl et al. 2014](#)). These may not be representative of ice nucleation processes occurring in typical cloud droplets of $\sim 20 \mu\text{m}$. In contact freezing studies, low collision efficiencies

pose a significant challenge to determining atmospheric relevance (Ladino et al. 2013). Although laboratory studies have shown that contact freezing is an efficient ice nucleation mechanism, it remains necessary to demonstrate relevance in clouds to understand if it competes with other freezing mechanisms.

b. Inertial separation and inlet artifacts

It has now been demonstrated that artifacts are produced by the collision of ice crystals with aircraft and inlet surfaces (Murphy et al. 2004). The reason for this is that ice crystals and large droplets have significant inertia and do not follow gas flow lines (Korolev and Isaac 2005). A surface impaction is not necessary; ice crystals may also break up upon the deceleration required for sampling (Pekour and Cziczo 2011). The generation of artifacts can best be visualized as crystals breaking into multiple fragments and/or the pitting of surfaces by ice impact (Korolev and Isaac 2005; Murphy et al. 2004).

Cziczo and Froyd (2014) gave a comprehensive account of the various inlet artifacts that include, but are not limited to, 1) inlet material, 2) aircraft material (e.g., paint), 3) combinations of these with real IRs, and 4) liberation of particles that were previously deposited on the inlet surface. Cziczo and Froyd (2014) noted that artifact content most likely went unnoticed in pre-SPMS composition analyses. Evidence supporting the generation of artifacts by ice impact have also been provided by EM: Kojima et al. (2004) detected abundant aluminum-, silicon-, and zinc-containing particles, which were suggested to be aircraft paint chipped by ice crystals. Perring et al. (2013) showed a black carbon artifact that appeared in a forward-facing (but not rear-facing) inlet when cirrus ice was present.

As a result of these observations, modern CVI inlets are constructed with distinct materials that lend an ability to detect artifacts. These include titanium (Stith et al. 2011) and gold (Froyd et al. 2010) as well as design elements such as orthogonal pickoff (Cziczo et al. 2013) that minimize artifact concentration.

4. Challenges to improving observations

There are less than five aircraft-capable ice chambers in existence at the publication of this monograph. This limitation has a direct impact on the ability of our community to understand the spatial and temporal variability of INPs. The SPMSs currently in use for both mountaintop and airborne studies of INPs are of a similarly limited number. The result is that many flight or deployment opportunities go unrealized not because of a lack of interest but because of a lack of instrumentation.

As mentioned in the previous sections, the CVI technique has allowed for investigation of IR properties

for over two decades. Airborne inlets are, to date, inertial separation systems only because of the complexity, mass, and size limitations of phase discrimination in the Ice-CVI and ISI. Furthermore, airborne CVIs are limited in upper cutpoint by the stopping distance of large ice at aircraft speeds. This has resulted in (often unrealized) artifact generation as well as no current information on the composition of IRs from crystals larger than $\sim 50\text{-}\mu\text{m}$ diameter (Cziczo and Froyd 2014).

A concerted effort toward reporting global INP concentrations has been suggested since the 1950s but never realized. Using a suite of methods/instruments at specific locations, this challenging task is now realizable with sufficient instrumentation and funding. Models are currently limited by a lack of data and this, in turn, limits our ability to correctly predict ice cloud formation and radiative effects (Hoose and Möhler 2012).

Acknowledgments. The authors would like to thank the many sponsors who have provided funding for the monograph: Leibniz Institute for Tropospheric Research (TROPOS), Forschungszentrum Jülich (FZJ), and Deutsches Zentrum für Luft- und Raumfahrt (DLR), Germany; ETH Zurich, Switzerland; National Center for Atmospheric Research (NCAR), United States; the Met Office, United Kingdom; the University of Illinois, United States; Environment and Climate Change Canada (ECCC), Canada; National Science Foundation (NSF), AGS 1723548, National Aeronautics and Space Administration (NASA), United States; the International Commission on Clouds and Precipitation (ICCP), the European Facility for Airborne Research (EUFAR), and Droplet Measurement Technologies (DMT), United States. D.J.C. acknowledges funding from the Victor Starr Career Development Chair. Y.B. acknowledges funding from the Swiss National Science Foundation (200020 150169/1). S.M. acknowledges funding from the German Research Foundation Research Unit INUIT (grant STR453/7-1) and DFG Transregional Collaborative Research Center SFB/TR 172 (project B03). H.W. acknowledges funding from the German Research Foundation (DFG project WE 4722/1-2) within the DFG Research Unit FOR 1525 INUIT. NCAR is sponsored by the NSF. Any opinions, findings, and conclusions or recommendations expressed in this publication are those of the author(s) and do not necessarily reflect the views of the National Science Foundation.

REFERENCES

- Alpert, P., J. Aller, and D. Knopf, 2011: Ice nucleation from aqueous NaCl droplets with and without marine diatoms. *Atmos. Chem. Phys.*, **11**, 5539–5555, doi:10.5194/acp-11-5539-2011.
- Anderson, T. L., R. J. Charlson, and D. S. Covert, 1993: Calibration of a counterflow virtual impactor at aerodynamic diameters

- from 1 to 15 μm . *Aerosol Sci. Technol.*, **19**, 317–329, doi:10.1080/02786829308959639.
- Archuleta, C. M., P. J. DeMott, and S. M. Kreidenweis, 2005: Ice nucleation by surrogates for atmospheric mineral dust and mineral dust/sulfate particles at cirrus temperatures. *Atmos. Chem. Phys.*, **5**, 2617–2634, doi:10.5194/acp-5-2617-2005.
- Ardon-Dryer, K., Y.-W. Huang, and D. J. Cziczo, 2015: Laboratory studies of collection efficiency of sub-micrometer particles by cloud droplets on a single droplet basis. *Atmos. Chem. Phys.*, **15**, 9159–9171, doi:10.5194/acp-15-9159-2015.
- Bailey, M., and J. Hallett, 2002: Nucleation effects on the habit of vapour grown ice crystals from -18 to -42°C . *Quart. J. Roy. Meteor. Soc.*, **128**, 1461–1483, doi:10.1002/qj.200212858304.
- Baustian, K., M. Wise, E. Jensen, G. Schill, M. Freedman, and M. Tolbert, 2013: State transformations and ice nucleation in amorphous (semi-) solid organic aerosol. *Atmos. Chem. Phys.*, **13**, 5615–5628, doi:10.5194/acp-13-5615-2013.
- Bigg, E. K., 1953: The formation of atmospheric ice crystals by the freezing of droplets. *Quart. J. Roy. Meteor. Soc.*, **79**, 510–519, doi:10.1002/qj.49707934207.
- Boose, Y., and Coauthors, 2016: Ice nucleating particle measurements at 241 K during winter months at 3580 m MSL in the Swiss Alps. *J. Atmos. Sci.*, **73**, 2203–2228, doi:10.1175/JAS-D-15-0236.1.
- Boulter, J. E., D. J. Cziczo, A. M. Middlebrook, D. S. Thomson, and D. M. Murphy, 2006: Design and performance of a pumped counterflow virtual impactor. *Aerosol Sci. Technol.*, **40**, 969–979, doi:10.1080/02786820600840984.
- Broadley, S., B. J. Murray, R. J. Herbert, J. D. Atkinson, S. Dobbie, T. L. Malkin, E. Condliffe, and L. Neve, 2012: Immersion mode heterogeneous ice nucleation by an illite rich powder representative of atmospheric mineral dust. *Atmos. Chem. Phys.*, **12**, 287–307, doi:10.5194/acp-12-287-2012.
- Brooks, S. D., K. Suter, and L. Olivarez, 2014: Effects of chemical aging on the ice nucleation activity of soot and polycyclic aromatic hydrocarbon aerosols. *J. Phys. Chem.*, **118A**, 10036–10047, doi:10.1021/jp508809y.
- Budke, C., and T. Koop, 2015: BINARY: An optical freezing array for assessing temperature and time dependence of heterogeneous ice nucleation. *Atmos. Meas. Tech.*, **8**, 689–703, doi:10.5194/amt-8-689-2015.
- Bundke, U., B. Nillius, R. Jaenicke, T. Wetter, H. Klein, and H. Bingemer, 2008: The Fast Ice Nucleus Chamber FINCH. *Atmos. Res.*, **90**, 180–186, doi:10.1016/j.atmosres.2008.02.008.
- Chen, Y., S. M. Kreidenweis, L. M. McInnes, D. C. Rogers, and P. J. DeMott, 1998: Single particle analyses of INPs in the upper troposphere and lower stratosphere. *Geophys. Res. Lett.*, **25**, 1391–1394, doi:10.1029/97GL03261.
- China, S., and Coauthors, 2015: Morphology of diesel soot residuals from supercooled water droplets and ice crystals: Implications for optical properties. *Environ. Res. Lett.*, **10**, 114010, doi:10.1088/1748-9326/10/11/114010.
- Chou, C., O. Stetzer, E. Weingartner, Z. Jurányi, Z. A. Kanji, and U. Lohmann, 2011: Ice nuclei properties within a Saharan dust event at the Jungfrauoch in the Swiss Alps. *Atmos. Chem. Phys.*, **11**, 4725–4738, doi:10.5194/acp-11-4725-2011.
- Conen, F., S. Henne, C. E. Morris, and C. Alewell, 2012: Atmospheric ice nucleators active $\geq -12^\circ\text{C}$ can be quantified on PM₁₀ filters. *Atmos. Meas. Tech.*, **5**, 321–327, doi:10.5194/amt-5-321-2012.
- Connolly, P. J., C. Emersic, and P. R. Field, 2012: A laboratory investigation into the aggregation efficiency of small ice crystals. *Atmos. Chem. Phys.*, **12**, 2055–2076, doi:10.5194/acp-12-2055-2012.
- Cozic, J., S. Mertes, B. Verheggen, D. J. Cziczo, S. J. Gallavardin, S. Walter, U. Baltensperger, and E. Weingartner, 2008: Black carbon enrichment in atmospheric ice particle residuals observed in lower tropospheric mixed phase clouds. *J. Geophys. Res.*, **113**, D15209, doi:10.1029/2007JD009266.
- Cziczo, D. J., and K. D. Froyd, 2014: Sampling the composition of cirrus ice residuals. *Atmos. Res.*, **142**, 15–31, doi:10.1016/j.atmosres.2013.06.012.
- , D. M. Murphy, P. K. Hudson, and D. S. Thompson, 2004: Single particle measurements of the chemical composition of cirrus ice residue during CRYSTAL-FACE. *J. Geophys. Res.*, **109**, doi:10.1029/2003JD004032.
- , and Coauthors, 2009: Inadvertent climate modification due to anthropogenic lead. *Nat. Geosci.*, **2**, 333–336, doi:10.1038/geo499.
- , and Coauthors, 2013: Clarifying the dominant sources and mechanisms of cirrus cloud formation. *Science*, **340**, doi:10.1126/science.1234145.
- Davis, E. J., 1997: A history of single particle levitation. *Aerosol Sci. Technol.*, **26**, 212–254, doi:10.1080/02786829708965426.
- Davis, R. D., S. Lance, J. A. Gordon, S. B. Ushijima, and M. A. Tolbert, 2015: Contact efflorescence as a pathway for crystallization of atmospherically relevant particles. *Proc. Natl. Acad. Sci. USA*, **112**, 15 815–15 820, doi:10.1073/pnas.1522860113.
- DeMott, P. J., D. J. Cziczo, A. J. Prenni, D. M. Murphy, S. M. Kreidenweis, D. S. Thomson, R. Borys, and D. C. Rogers, 2003: Measurements of the concentration and composition of nuclei for cirrus formation. *Proc. Natl. Acad. Sci. USA*, **100**, 14 655–14 660, doi:10.1073/pnas.2532677100.
- , and Coauthors, 2010: Predicting global atmospheric ice nuclei distributions and their impacts on climate. *Proc. Natl. Acad. Sci. USA*, **107**, 11 217–11 222, doi:10.1073/pnas.0910818107.
- , and Coauthors, 2011: Resurgence in ice nuclei measurement research. *Bull. Amer. Meteor. Soc.*, **92**, 1623–1635, doi:10.1175/2011BAMS3119.1.
- , and Coauthors, 2015: Integrating laboratory and field data to quantify the immersion freezing ice nucleation activity of mineral dust particles. *Atmos. Chem. Phys.*, **15**, 393–409, doi:10.5194/acp-15-393-2015.
- , and Coauthors, 2016: Sea spray aerosol as a unique source of INPs. *Proc. Natl. Acad. Sci. USA*, **113**, 5797–5803, doi:10.1073/pnas.1514034112.
- Diehl, K., S. K. Mitra, M. Szakáll, N. von Blohn, S. Borrmann, and H. R. Pruppacher, 2011: The Mainz vertical wind tunnel facility: A review of 25 years of laboratory experiments on cloud physics and chemistry. *Wind Tunnels: Aerodynamics, Models and Experiments*, J. D. Pereira, Ed., Nova Press, 69–92.
- , M. Debertshäuser, O. Eppers, H. Smithhüsen, S. K. Mitra, and S. Borrmann, 2014: Particle surface area dependence of mineral dust in immersion freezing mode: Investigations with freely suspended drops in an acoustic levitator and a vertical wind tunnel. *Atmos. Chem. Phys.*, **14**, 12 343–12 355, doi:10.5194/acp-14-12343-2014.
- Dufour, L., 1862: Über das Gefrieren des Wassers und über die Bildung des Hagels. *Ann. Phys. Chem.*, **190**, 530–554, doi:10.1002/andp.18621901203.
- Durant, A., and R. Shaw, 2005: Evaporation freezing by contact nucleation inside-out. *Geophys. Res. Lett.*, **32**, L20814, doi:10.1029/2005GL024175.
- Eastwood, M. L., S. Cremel, C. Gehrke, E. Girard, and A. K. Bertram, 2008: Ice nucleation on mineral dust particles: Onset conditions, nucleation rates and contact angles. *J. Geophys. Res.*, **113**, D22203, doi:10.1029/2008JD010639.

- Ebert, M., A. Worringer, N. Benker, S. Mertes, E. Weingartner, and S. Weinbruch, 2011: Chemical composition and mixing-state of IRs sampled within mixed-phase clouds. *Atmos. Chem. Phys.*, **11**, 2805–2816, doi:10.5194/acp-11-2805-2011.
- Freedman, M. A., 2015: Potential sites for ice nucleation on aluminosilicate clay minerals and related materials. *J. Phys. Chem. Lett.*, **6**, 3850–3858, doi:10.1021/acs.jpcclett.5b01326.
- Froyd, K. D., D. M. Murphy, P. Lawson, D. Baumgardner, and R. L. Herman, 2010: Aerosols that form subvisible cirrus at the tropical tropopause. *Atmos. Chem. Phys.*, **10**, 209–218, doi:10.5194/acp-10-209-2010.
- Garimella, S., and Coauthors, 2016: The SPectrometer for Ice Nuclei (SPIN): An instrument to investigate ice nucleation. *Atmos. Meas. Tech.*, **9**, 2781–2795, doi:10.5194/amt-9-2781-2016.
- Hallett, J., and S. C. Mossop, 1974: Production of secondary ice particles during the riming process. *Nature*, **249**, 26–28, doi:10.1038/249026a0.
- Hartmann, S., D. Niedermeier, J. Voigtländer, T. Clauss, R. A. Shaw, H. Wex, A. Kiselev, and F. Stratmann, 2011: Homogeneous and heterogeneous ice nucleation at LACIS: Operating principle and theoretical studies. *Atmos. Chem. Phys.*, **11**, 1753–1767, doi:10.5194/acp-11-1753-2011.
- , H. Wex, T. Clauss, S. Augustin-Bauditz, D. Niedermeier, M. Rösch, and F. Stratmann, 2016: Immersion freezing of kaolinite: Scaling with particle surface area. *J. Atmos. Sci.*, **73**, 263–278, doi:10.1175/JAS-D-15-0057.1.
- Heintzenberg, J., K. Okada, and J. Ström, 1996: On the composition of non-volatile material in upper tropospheric aerosols and cirrus crystals. *Atmos. Res.*, **41**, 81–88, doi:10.1016/0169-8095(95)00042-9.
- Hill, T. C., B. F. Moffett, P. J. DeMott, D. G. Georgakopoulos, W. L. Stump, and G. D. Franc, 2014: Measurement of ice nucleation-active bacteria on plants and in precipitation by quantitative PCR. *Appl. Environ. Microbiol.*, **80**, 1256–1267, doi:10.1128/AEM.02967-13.
- Hiranuma, N., and Coauthors, 2015a: A comprehensive laboratory study on the immersion freezing behavior of illite NX particles: A comparison of 17 ice nucleation measurement techniques. *Atmos. Chem. Phys.*, **15**, 2489–2518, doi:10.5194/acp-15-2489-2015.
- , and Coauthors, 2015b: Ice nucleation by cellulose and its potential contribution to ice formation in clouds. *Nat. Geosci.*, **8**, 273–277, doi:10.1038/ngeo2374.
- Hoffmann, N., D. Duft, A. Kiselev, and T. Leisner, 2013a: Contact freezing efficiency of mineral dust aerosols studied in an electrodynamic balance: Quantitative size and temperature dependence for illite particles. *Faraday Discuss.*, **165**, 383–390, doi:10.1039/c3fd00033h.
- , A. Kiselev, D. Rzesanke, D. Duft, and T. Leisner, 2013b: Experimental quantification of contact freezing in an electrodynamic balance. *Atmos. Meas. Tech.*, **6**, 2373–2382, doi:10.5194/amt-6-2373-2013.
- Hoose, C., and O. Möhler, 2012: Heterogeneous ice nucleation on atmospheric aerosols: A review of results from laboratory experiments. *Atmos. Chem. Phys.*, **12**, 9817–9854, doi:10.5194/acp-12-9817-2012.
- Isono, K., 1955: On ice-crystal nuclei and other substances found in snow crystals. *J. Meteor.*, **12**, 456–462, doi:10.1175/1520-0469(1955)012<0456:OICNAO>2.0.CO;2.
- Kamphus, M., and Coauthors, 2010: Chemical composition of ambient aerosol, ice residues and cloud droplet residues in mixed-phase clouds: Single particle analysis during the Cloud and Aerosol Characterization Experiment (CLACE 6). *Atmos. Chem. Phys.*, **10**, 8077–8095, doi:10.5194/acp-10-8077-2010.
- Kanji, Z. A., and J. P. D. Abbatt, 2006: Laboratory studies of ice formation via deposition mode nucleation onto mineral dust and n-hexane soot samples. *J. Geophys. Res.*, **111**, D16204, doi:10.1029/2005JD006766.
- , and —, 2009: The University of Toronto Continuous Flow Diffusion Chamber (UT-CFDC): A simple design for ice nucleation studies. *Aerosol Sci. Technol.*, **43**, 730–738, doi:10.1080/02786820902889861.
- , and —, 2010: Ice nucleation onto Arizona Test Dust at cirrus temperatures: Effect of temperature and aerosol size on onset relative humidity. *J. Phys. Chem.*, **114A**, 935–941, doi:10.1021/jp908661m.
- , L. A. Ladino, H. Wex, Y. Boose, M. Burkert-Kohn, D. Cziczo, and M. Krämer, 2017: Heterogeneous ice nucleation. *Ice Formation and Evolution in Clouds and Precipitation: Measurement and Modeling Challenges, Meteor. Monogr.*, No. 58, Amer. Meteor. Soc., doi:10.1175/AMSMONOGRAPHSD-16-0006.1.
- Kaye, P. H., E. Hirst, R. S. Greenaway, Z. Ulanowski, E. Hesse, P. J. DeMott, C. Saunders, and P. Connolly, 2008: Classifying atmospheric ice crystals by spatial light scattering. *Opt. Lett.*, **33**, 1545–1547, doi:10.1364/OL.33.001545.
- Knopf, D. A., B. Wang, A. Laskin, R. C. Moffet, and M. K. Gilles, 2010: Heterogeneous nucleation of ice on anthropogenic organic particles collected in Mexico City. *Geophys. Res. Lett.*, **37**, L11803, doi:10.1029/2010GL043362.
- Kohn, M., U. Lohmann, A. Welti, and Z. A. Kanji, 2016: Immersion mode ice nucleation measurements with the new Portable Immersion Mode Cooling Chamber (PIMCA). *J. Geophys. Res.*, **121**, 4713–4733, doi:10.1002/2016JD024761.
- Kojima, T., P. R. Buseck, J. C. Wilson, J. M. Reeves, and M. J. Mahoney, 2004: Particles from tropical convective systems: Cloud tops and cirrus anvils. *J. Geophys. Res.*, **109**, D12201, doi:10.1029/2003JD004504.
- Koop, T., H. P. Ng, L. T. Molina, and M. J. Molina, 1998: A new optical technique to study aerosol phase transitions: The nucleation of ice from H₂SO₄ aerosols. *J. Phys. Chem.*, **102A**, 8924–8931, doi:10.1021/jp9828078.
- Korolev, A., and G. A. Isaac, 2005: Shattering during sampling by OAPs and HVPS. Part I: Snow particles. *J. Atmos. Oceanic Technol.*, **22**, 528–542, doi:10.1175/JTECH1720.1.
- Kumai, M., 1961: Snow crystals and the identification of the nuclei in the northern United States of America. *J. Meteor.*, **18**, 139–150, doi:10.1175/1520-0469(1961)018<0139:SCATIO>2.0.CO;2.
- Kupiszewski, P., and Coauthors, 2015: The Ice Selective Inlet: A novel technique for exclusive extraction of pristine ice crystals in mixed-phase clouds. *Atmos. Meas. Tech.*, **8**, 3087–3106, doi:10.5194/amt-8-3087-2015.
- , and Coauthors, 2016: Ice residual properties in mixed-phase clouds at the high-alpine Jungfrauoch site. *J. Geophys. Res. Atmos.*, **121**, 12 343–12 362, doi:10.1002/2016JD024894.
- Ladino, L. A., O. Stetzer, F. Lüönd, A. Welti, and U. Lohmann, 2011: Contact freezing experiments of kaolinite particles with cloud droplets. *J. Geophys. Res.*, D22202, doi:10.1029/2011JD015727.
- , —, and U. Lohmann, 2013: Contact freezing: A review of experimental studies. *Atmos. Chem. Phys.*, **13**, 9745–9769, doi:10.5194/acp-13-9745-2013.
- Langer, G., J. Rosinski, and C. P. Edwards, 1967: A continuous ice nucleus counter and its application to tracking in the troposphere. *J. Appl. Meteor.*, **6**, 114–125, doi:10.1175/1520-0450(1967)006<0114:ACINCA>2.0.CO;2.
- López, M. L., and E. E. Ávila, 2013: Measurements of natural deposition ice nuclei in Córdoba, Argentina. *Atmos. Chem. Phys.*, **13**, 3111–3119, doi:10.5194/acp-13-3111-2013.

- Lüönd, F., O. Stetzer, A. Welti, and U. Lohmann, 2010: Experimental study on the ice nucleation ability of size-selected kaolinite particles in the immersion mode. *J. Geophys. Res.*, **115**, D14201, doi:10.1029/2009JD012959.
- Marcilli, C., 2014: Deposition nucleation viewed as homogeneous or immersion freezing in pores and cavities. *Atmos. Chem. Phys.*, **14**, 2071–2104, doi:10.5194/acp-14-2071-2014.
- Mason, R. H., and Coauthors, 2015a: The micro-orifice uniform deposit impactor–droplet freezing technique (MOUDI-DFT) for measuring concentrations of ice nucleating particles as a function of size: Improvements and initial validation. *Atmos. Meas. Tech.*, **8**, 2449–2462, doi:10.5194/amt-8-2449-2015.
- , and Coauthors, 2015b: Ice nucleating particles at a coastal marine boundary layer site: Correlations with aerosol type and meteorological conditions. *Atmos. Chem. Phys.*, **15**, 12 547–12 566, doi:10.5194/acp-15-12547-2015.
- , and Coauthors, 2016: Size-resolved measurements of ice-nucleating particles at six locations in North America and one in Europe. *Atmos. Chem. Phys.*, **16**, 1637–1651, doi:10.5194/acp-16-1637-2016.
- Mertes, S., A. Schwarzenböck, P. Laj, W. Wobrock, J.-M. Pichon, G. Orsi, and J. Heintzenberg, 2001: Changes of cloud microphysical properties during the transition from supercooled to mixed-phase conditions during CIME. *Atmos. Res.*, **58**, 267–294, doi:10.1016/S0169-8095(01)00095-3.
- , and Coauthors, 2007: Counterflow virtual impactor based collection of small ice particles in mixed-phase clouds for the physico-chemical characterization of tropospheric ice nuclei: Sampler description and first case study. *Aerosol Sci. Technol.*, **41**, 848–864, doi:10.1080/02786820701501881.
- Moffet, R. C., and Coauthors, 2010: Microscopic characterization of carbonaceous particle aging in the outflow from Mexico City. *Atmos. Chem. Phys.*, **10**, 961–976, doi:10.5194/acp-10-961-2010.
- Möhler, O., and Coauthors, 2003: Experimental investigation of homogeneous freezing of sulphuric acid particles in the aerosol chamber AIDA. *Atmos. Chem. Phys.*, **3**, 211–223, doi:10.5194/acp-3-211-2003.
- Mossop, S. C., 1970: Concentrations of ice crystals in clouds. *Bull. Amer. Meteor. Soc.*, **51**, 474–479, doi:10.1175/1520-0477(1970)051<0474:COICIC>2.0.CO;2.
- Murphy, D. M., 2007: The design of single particle laser mass spectrometers. *Mass Spectrom. Rev.*, **26**, 150–165, doi:10.1002/mas.20113.
- , D. J. Cziczo, P. K. Hudson, D. S. Thomson, J. C. Wilson, T. Kojima, and P. R. Buseck, 2004: Particle generation and resuspension in aircraft inlets when flying in clouds. *Aerosol Sci. Technol.*, **38**, 401–409, doi:10.1080/02786820490443094.
- Niedermeier, D., and Coauthors, 2011: Experimental study of the role of physicochemical surface processing on the IN ability of mineral dust particles. *Atmos. Chem. Phys.*, **11**, 11 131–11 144, doi:10.5194/acp-11-11131-2011.
- Niehaus, J., K. W. Bunker, S. China, A. Kostinski, C. Mazzoleni, and W. Cantrell, 2014: A technique to measure ice nuclei in the contact mode. *J. Atmos. Oceanic Technol.*, **31**, 913–922, doi:10.1175/JTECH-D-13-00156.1.
- Noone, K. J., J. A. Ogren, J. Heintzenberg, R. J. Charlson, and D. S. Covert, 1988: Design and calibration of a counterflow virtual impactor for sampling of atmospheric fog and cloud droplets. *Aerosol Sci. Technol.*, **8**, 235–244, doi:10.1080/02786828808959186.
- , H.-C. Hansson, and R. K. A. M. Mallant, 1992: Droplet sampling from crosswinds: An inlet efficiency calibration. *J. Aerosol Sci.*, **23**, 153–164, doi:10.1016/0021-8502(92)90051-V.
- Ogren, J. A., J. Heintzenberg, and R. J. Charlson, 1985: In-situ sampling of clouds with a droplet to aerosol converter. *Geophys. Res. Lett.*, **12**, 121–124, doi:10.1029/GL012i003p00121.
- Palmer, H. P., 1949: Natural ice-particle nuclei. *Quart. J. Roy. Meteor. Soc.*, **75**, 17–22, doi:10.1002/qj.49707532303.
- Pekour, M., and D. J. Cziczo, 2011: Wake capture, particle breakup and other artifacts associated with counterflow virtual impaction. *Aerosol Sci. Technol.*, **45**, 758–764, doi:10.1080/02786826.2011.558942.
- Perring, A. E., J. P. Schwarz, R. S. Gao, A. J. Heymsfeld, C. G. Schmitt, M. Schnaiter, and D. W. Fahey, 2013: Evaluation of a perpendicular inlet for airborne sampling of interstitial sub-micron aerosol. *Aerosol Sci. Technol.*, **47**, 1066–1072, doi:10.1080/02786826.2013.821196.
- Petters, M. D., and T. P. Wright, 2015: Revisiting ice nucleation from precipitation samples. *Geophys. Res. Lett.*, **42**, 8758–8766, doi:10.1002/2015GL065733.
- Petzold, A., J. Ström, S. Ohlsson, and F. P. Schroder, 1998: Elemental composition and morphology of ice-crystal residual particles in cirrus clouds and contrails. *Atmos. Res.*, **49**, 21–34, doi:10.1016/S0169-8095(97)00083-5.
- Pratt, K. A., and Coauthors, 2009: In situ detection of biological particles in cloud ice-crystals. *Nat. Geosci.*, **2**, 398–401, doi:10.1038/ngeo521.
- Rogers, D. C., 1988: Development of a continuous flow thermal gradient diffusion chamber for ice nucleation studies. *Atmos. Res.*, **22**, 149–181, doi:10.1016/0169-8095(88)90005-1.
- , P. J. DeMott, S. M. Kreidenweis, and Y. Chen, 2001: A continuous-flow diffusion chamber for airborne measurements of ice nuclei. *J. Atmos. Oceanic Technol.*, **18**, 725–741, doi:10.1175/1520-0426(2001)018<0725:ACFDCF>2.0.CO;2.
- Rogers, R. R., and G. Vali, 1978: Recent developments in meteorological physics. *Phys. Rep.*, **48**, 65–177, doi:10.1016/0370-1573(78)90078-9.
- Rosinski, J., P. L. Haagenson, C. T. Nagamoto, and F. Parungo, 1986: Ice-forming nuclei of maritime origin. *J. Aerosol Sci.*, **17**, 23–46, doi:10.1016/0021-8502(86)90004-2.
- Schill, G. P., D. O. De Haan, and M. A. Tolbert, 2014: Heterogeneous ice nucleation on simulated secondary organic aerosol. *Environ. Sci. Technol.*, **48**, 1675–1682, doi:10.1021/es4046428.
- Schnell, R. C., 1977: Ice nuclei in seawater, fog water and marine air off the coast of Nova Scotia: Summer 1975. *J. Atmos. Sci.*, **34**, 1299–1305, doi:10.1175/1520-0469(1977)034<1299:INISFW>2.0.CO;2.
- , and G. Vali, 1976: Biogenic ice nuclei: Part I. Terrestrial and marine sources. *J. Atmos. Sci.*, **33**, 1554–1564, doi:10.1175/1520-0469(1976)033<1554:BINPIT>2.0.CO;2.
- Seifert, M., J. Ström, R. Krejci, A. Minikin, A. Petzold, J.-F. Gayet, U. Schumann, and J. Olvarlez, 2003: In situ observations of particles remaining from evaporated cirrus crystals: Comparing clean and polluted air masses. *Atmos. Chem. Phys.*, **3**, 1037–1049, doi:10.5194/acp-3-1037-2003.
- , and Coauthors, 2004: Thermal stability analysis of particles incorporated in cirrus crystals and of non-activated particles in between the cirrus crystals: Comparing clean and polluted air masses. *Atmos. Chem. Phys.*, **4**, 1343–1353, doi:10.5194/acp-4-1343-2004.
- Shilling, J. E., T. J. Fortin, and M. A. Tolbert, 2006: Depositional ice nucleation on crystalline organic and inorganic solids. *J. Geophys. Res.*, **111**, D05207, doi:10.1029/2005JD006664.
- Sihvonen, S. K., G. P. Schill, N. A. Lykтей, D. P. Veghte, M. A. Tolbert, and M. A. Freedman, 2014: Chemical and physical transformations of aluminosilicate clay minerals due to

- acid treatment and consequences for heterogeneous ice nucleation. *J. Phys. Chem.*, **118A**, 8787–8796, doi:10.1021/jp504846g.
- Stetzer, O., B. Baschek, F. Lüönd, and U. Lohmann, 2008: The Zurich Ice Nucleation Chamber (ZINC)—A new instrument to investigate atmospheric ice formation. *Aerosol Sci. Technol.*, **42**, 64–74, doi:10.1080/02786820701787944.
- Stith, J. L., C. H. Twohy, P. J. DeMott, D. Baumgardner, T. Campos, R. Gao, and J. Anderson, 2011: Observations of ice nuclei and heterogeneous freezing in a western Pacific extratropical storm. *Atmos. Chem. Phys.*, **11**, 6229–6249, doi:10.5194/acp-11-6229-2011.
- Ström, J., and J. Heintzenberg, 1994: Water vapor, condensed water and crystal concentration in orographically influenced cirrus clouds. *J. Atmos. Sci.*, **51**, 2368–2383, doi:10.1175/1520-0469(1994)051<2368:WVCWAC>2.0.CO;2.
- , B. Strauss, T. Anderson, J. Schröder, J. Heintzenberg, and P. Wendling, 1997: In situ observations of the microphysical properties of young cirrus clouds. *J. Atmos. Sci.*, **54**, 2542–2553, doi:10.1175/1520-0469(1997)054<2542:ISOOTM>2.0.CO;2.
- Szyrmer, W., and I. Zawadzki, 1997: Biogenic and anthropogenic sources of ice-forming nuclei: A review. *Bull. Amer. Meteor. Soc.*, **78**, 209–228, doi:10.1175/1520-0477(1997)078<0209:BAASOI>2.0.CO;2.
- Targino, A. C., R. Krejci, K. J. Noone, and P. Glantz, 2006: Single particle analysis of ice crystal residuals observed in orographic wave clouds over Scandinavia during INTACC experiment. *Atmos. Chem. Phys.*, **6**, 1977–1990, doi:10.5194/acp-6-1977-2006.
- Tenberken-Pötzsch, B., M. Schwikowski, and H. W. Gäggeler, 2000: A method to sample and separate ice crystals and supercooled cloud droplets in mixed phase clouds for subsequent chemical analysis. *Atmos. Environ.*, **34**, 3629–3633, doi:10.1016/S1352-2310(00)00140-0.
- Twohy, C. H., and B. W. Gandrud, 1998: Electron microscope analysis of residual particles from aircraft contrails. *Geophys. Res. Lett.*, **25**, 1359–1362, doi:10.1029/97GL03162.
- , and M. R. Poellot, 2005: Chemical characteristics of IR nuclei in anvil cirrus clouds: Evidence for homogeneous and heterogeneous ice formation. *Atmos. Chem. Phys.*, **5**, 2289–2297, doi:10.5194/acp-5-2289-2005.
- , A. J. Schanot, and W. A. Cooper, 1997: Measurement of condensed water content in liquid and ice clouds using an airborne counterflow virtual impactor. *J. Atmos. Oceanic Technol.*, **14**, 197–202, doi:10.1175/1520-0426(1997)014<0197:MOCWCI>2.0.CO;2.
- , J. W. Strapp, and M. Wendisch, 2003: Performance of a counterflow virtual impactor in the NASA Icing Research Tunnel. *J. Atmos. Oceanic Technol.*, **20**, 781–790, doi:10.1175/1520-0426(2003)020<0781:POACVI>2.0.CO;2.
- Vali, G., 1966: Sizes of atmospheric ice nuclei. *Nature*, **212**, 384–385, doi:10.1038/212384a0.
- , M. Christensen, R. W. Fresh, E. L. Galyan, L. R. Maki, and R. C. Schnell, 1976: Biogenic ice nuclei. Part II: Bacterial sources. *J. Atmos. Sci.*, **33**, 1565–1570, doi:10.1175/1520-0469(1976)033<1565:BINPIB>2.0.CO;2.
- , P. J. DeMott, O. Möhler, and T. F. Whale, 2015: Technical note: A proposal for ice nucleation terminology. *Atmos. Chem. Phys.*, **15**, 10263–10270, doi:10.5194/acp-15-10263-2015.
- Veghte, D. P., D. R. Bittner, and M. A. Freedman, 2014: Cryo-transmission electron microscopy imaging of the morphology of submicrometer aerosol containing organic acids and ammonium sulfate. *Anal. Chem.*, **86**, 2436–2442, doi:10.1021/ac403279f.
- Vidaurre, G., and J. Hallett, 2009: Particle impact and breakup in aircraft measurement. *J. Atmos. Oceanic Technol.*, **26**, 972–983, doi:10.1175/2008JTECHA1147.1.
- Vochezer, P., E. Järvinen, R. Wagner, P. Kupiszewski, T. Leisner, and M. Schnaiter, 2016: In situ characterization of mixed phase clouds using the small ice detector and the particle phase discriminator. *Atmos. Meas. Tech.*, **9**, 159–177, doi:10.5194/amt-9-159-2016.
- Vonnegut, B., 1947: The nucleation of ice formation by silver iodide. *J. Appl. Phys.*, **18**, 593–595, doi:10.1063/1.1697813.
- Wang, B. B., A. T. Lambe, P. Massoli, T. B. Onasch, P. Davidovits, D. R. Worsnop, and D. A. Knopf, 2012a: The deposition ice nucleation and immersion freezing potential of amorphous secondary organic aerosol: Pathways for ice and mixed-phase cloud formation. *J. Geophys. Res.*, **117**, D16209, doi:10.1029/2012JD018063.
- , A. Laskin, T. Roedel, M. K. Gilles, R. C. Moffet, A. V. Tivanski, and D. A. Knopf, 2012b: Heterogeneous ice nucleation and water uptake by field-collected atmospheric particles below 273 K. *J. Geophys. Res.*, **117**, D00V19, doi:10.1029/2012JD017446.
- Whale, T. F., and Coauthors, 2015: A technique for quantifying heterogeneous ice nucleation in microliter supercooled water droplets. *Atmos. Meas. Tech.*, **8**, 2437–2447, doi:10.5194/amt-8-2437-2015.
- Wilson, T. W., and Coauthors, 2015: A marine biogenic source of atmospheric ice-nucleating particles. *Nature*, **525**, 234–238, doi:10.1038/nature14986.
- Wise, M. E., K. J. Baustian, T. Koop, M. A. Freedman, E. J. Jensen, and M. A. Tolbert, 2012: Depositional ice nucleation onto crystalline hydrated NaCl particles: A new mechanism for ice formation in the troposphere. *Atmos. Chem. Phys.*, **12**, 1121–1134, doi:10.5194/acp-12-1121-2012.
- Worringen, A., and Coauthors, 2015: Single-particle characterization of ice-nucleating particles and ice particle residuals sampled by three different techniques. *Atmos. Chem. Phys.*, **15**, 4161–4178, doi:10.5194/acp-15-4161-2015.
- Wright, T. P., and M. D. Petters, 2013: The role of time in heterogeneous freezing nucleation. *J. Geophys. Res. Atmos.*, **118**, 3731–3743, doi:10.1002/jgrd.50365.
- Young, G., and Coauthors, 2016: Size-segregated compositional analysis of aerosol particles collected in the European Arctic during the ACCACIA campaign. *Atmos. Chem. Phys.*, **16**, 4063–4079, doi:10.5194/acp-16-4063-2016.
- Zimmermann, F., M. Ebert, A. Worringen, L. Schütz, and S. Weinbruch, 2007: Environmental scanning electron microscopy (ESEM) as a new technique to determine the ice nucleation capability of individual atmospheric particles. *Atmos. Environ.*, **41**, 8219–8227, doi:10.1016/j.atmosenv.2007.06.023.
- Zolles, T., J. Burkart, T. Häusler, B. Pummer, R. Hitzemberger, and H. Grothe, 2015: Identification of ice nucleation active sites on feldspar dust particles. *J. Phys. Chem.*, **119A**, 2692–2700, doi:10.1021/jp509839x.
- Zuberi, B., A. K. Bertram, T. Koop, L. T. Molina, and M. J. Molina, 2001: Heterogeneous freezing of aqueous particles induced by crystallized (NH₄)₂SO₄, ice, and letovicite. *J. Phys. Chem.*, **105A**, 6458–6464, doi:10.1021/jp010094e.