

Near Infrared Radiation in Northern Greenland

S. J. BOLSENGA¹

21 November 1966 and 17 December 1966

1. Introduction

Measurements of near infrared ($0.7\text{--}3.0\ \mu$) direct solar and global (direct+diffuse) radiation were taken on clear days near Thule, on the edge of the Icecap, in

¹ Presently at the Great Lakes Research Center, U. S. Lake Survey, Detroit, Mich. This project was sponsored by and conducted at the U. S. Army Cold Regions Research & Engineering Laboratory, Hanover, N. H.

Northern Greenland ($76^{\circ}24'N$, $68^{\circ}19'W$) during May and June 1964. Since published information on near infrared radiation in polar regions is limited, this note presents a brief analysis of the measurements. Instrumentation consisted of Epply pyranometers, 2 m above the surface, and normal incidence pyrhemometers installed on a manually operated altitude-azimuth mount. All instruments were equipped with Schott RG-8 filters.




FIG. 1. Near infrared global radiation G_{IR} as a function of the solar altitude γ .

Pyranometers were leveled and filters cleaned daily or as required. Measurements were recorded by a potentiometric strip-chart recorder which was calibrated daily and periodic time checks were from a time standard radio signal.

2. Results

Fig. 1 shows the diurnal pattern of near infrared global radiation on a day of nearly average atmospheric conditions as observed during the period. Maximum solar altitude γ was about 33° or only 3° less than at the summer solstice. Mean atmospheric turbidity T_p , computed according to IGY recommendations (International Council of Scientific Unions, 1958), and atmospheric water vapor content w_T , computed from radio-

sonde data, were 2.4 and 0.16 cm, respectively. For the entire measurement period T_p averaged 2.6 with a range of 2.3–3.5 and w_T averaged 0.16 cm with a range of 0.13–0.18 cm.

Near infrared global radiation constituted about 51% of the total global radiation for all days of measurement and remained in the range 50–53%.

Fig. 2 shows near infrared and total solar beam transmittance as a function of γ . Computations were made using the solar constant (International Council of Scientific Unions, 1958) for extraterrestrial total solar radiation and by graphical integration of extraterrestrial spectral solar intensity curves (Fritz, 1951) for extraterrestrial near infrared solar radiation. Near infrared solar beam transmittance ranged between 68% when $\gamma = 36^\circ$ to 47% when $\gamma = 11^\circ$. Total solar beam transmittance varied from about 59% at $\gamma = 36^\circ$ to 29% at $\gamma = 10^\circ$.

The highest value of diffuse sky radiation, calculated from global and normal incidence radiation, was $0.12\ ly\ min^{-1}$ in the near infrared and $0.26\ ly\ min^{-1}$ in the total solar spectrum. Both values occurred on 28 May at 0818 TST when $\gamma = 28^\circ$. T_p and w_T were the highest observed during the entire period. The lowest value of diffuse radiation, at $\gamma \geq 15^\circ$, was $0.03\ ly\ min^{-1}$ for the near infrared and $0.11\ ly\ min^{-1}$ for the total solar spectrum. Both values occurred on 1 June at 1954 TST when $\gamma = 15^\circ$. T_p was fairly low (2.5) and w_T was the lowest recorded during the entire period.

Diffuse radiation constituted about 16% of the global radiation in the near infrared and about 23% in the total solar spectrum. Gerdel and Diamond (1956) found that diffuse radiation was about 19% of global radiation at a site on the Greenland icecap with a continuous snow surface. Higher atmospheric turbidity at the edge of the icecap probably contributes to the higher percentage of diffuse radiation to global radiation. Turbidity measurements made near the location of Gerdel and Diamond's observations averaged 1.9 during a one month period in August and September 1963.




FIG. 2. Near infrared T_{IR} and total T_T solar beam transmittance as a function of the solar altitude γ .

REFERENCES

Fritz, 1951: Solar radiant energy and its modification by the earth and its atmosphere. *Compendium of Meteorology*, Boston, Amer. Meteor. Soc., 13-33.
 Gerdel, R. W., and M. Diamond, 1956: Radiation measurements

on the Greenland ice cap. U. S. Army Snow, Ice and Permafrost Research Establishment, Research Report 19, 9 pp.
 International Council of Scientific Unions, 1958: *Annals of the International Geophysical Year*. Vol. 5, London, Pergamon Press, 371-405.

A Rational Approximation for Saturation Vapor Pressure over the Temperature Range of Sea Water

W. E. LANGLOIS

IBM Research Laboratory, San Jose, Calif.

4 November 1966 and 7 December 1966

It is an immediate consequence of the Clausius-Clapeyron equation that the saturation vapor pressure e_s of an ideal gas in equilibrium with its own condensate is a decaying exponential function of $L/(R_s T)$, where L is the latent heat, R_s the specific gas constant for the substance, and T the Kelvin temperature. For water vapor in equilibrium with liquid water and e_s in millibars, the handbook (Berry *et al.*, 1945, p. 343) form of the complete relation is

$$\ln\left(\frac{e_s}{6.105}\right) = 25.22\left(\frac{T-273}{T}\right) - 5.31 \ln\left(\frac{T}{273}\right), \quad (1)$$

which accounts for the dependence of latent heat upon temperature.

When e_s is to be evaluated by computer, for example in a general circulation model with hydrological cycle, computational efficiency is substantially improved if we replace (1) by a rational approximation. We begin by selecting a value T_0 near the center of the range to be treated. Expanding the logarithm in (1), we obtain

$$e_s = A \exp\{B(1-T_0/T) - C(1-T_0/T)^2 + O[(1-T_0/T)^3]\},$$

where

$$A = 6.105 \exp\{25.22(1-273/T_0) + 5.31 \ln(273/T_0)\},$$

$$B = 25.22(273/T_0) - 5.31, \quad C = 2.655.$$

Next we employ the continued fraction expansion for the exponential function (Wall, 1948, p. 348). Truncating at the fifth term, rewriting as a simple fraction, and neglecting terms of order $(1-T_0/T)^8$ in numerator and denominator yields the approximate formula

$$e_s \doteq \frac{T^2 - c_1 T + c_2}{c_3 T^2 - c_4 T + c_5}, \quad (2)$$

with

$$c_1 = 2(B^2 + 3B - 6C)T_0/D,$$

$$c_2 = (B^2 - 6C)T_0^2/D,$$

$$c_3 = (12 + B^2 - 6B + 6C)/AD,$$

$$c_4 = 2(B^2 - 3B + 6C)T_0/AD,$$

$$c_5 = (B^2 + 6C)T_0^2/AD,$$




FIG. 1. Error curves for the formulas.

where

$$D = 12 + B^2 + 6B - 6C.$$

To cover the temperature range of sea water, we choose $T_0 = 287$. The approximation (2) then becomes

$$e_s \doteq \frac{T^2 - 488.56T + 60009.3}{0.0361622T^2 - 24.209T + 4104.45}. \quad (3)$$

Comparing results obtained from (3) with the observed values (Byers, 1959, p. 158) yields the error curve shown in Fig. 1. The corresponding curve for the handbook formula is included for comparison. The derivation of (3) indicates that the curves should touch at 14°C, but the rational approximation was improved somewhat by judicious rounding of the c_i . The originally computed values, to seven significant figures, were

$$c_1 = 488.5568, \quad c_2 = 60009.27$$

$$c_3 = 0.03616217, \quad c_4 = 24.20942, \quad c_5 = 4104.454.$$

REFERENCES

Berry, F. A., Jr., E. Bolla and N. R. Beers, 1945: *Handbook of Meteorology*. New York, McGraw-Hill Book Co., 343 pp.
 Byers, H. R., 1959: *General Meteorology*. New York, McGraw-Hill Book Co., 158 pp.
 Wall, H. S., 1948: *Analytic Theory of Continued Fractions*. Princeton, N. J., D. Van Nostrand Company, Inc., 348 pp.