Reply

M. C. SHEN

The University of Wisconsin, Madison

3 December 1966

Although linearized equations are insufficient to explain the phenomenon of waves of permanent type, which is essentially nonlinear in nature, they do yield important information about the so-called critical speeds associated with such waves. In general, an eigenvalue problem based upon the linearized equations can be set up, and a relation $F(c^2, k^2) = 0$ between the wave speed c and the wave number k is then obtained,

where k^2 may assume continuous values from 0 to ∞ . It has been shown (Peters and Stoker, 1960) that at the limiting speeds b of c as k tends to zero, the linear waves are unstable and the linear theory fails. However, it turns out that the values of b so obtained are exactly the same as the critical speeds found from the nonlinear theory, near which stable, solitary and cnoidal waves will appear.

In a previous paper (Shen, 1965) on solitary waves in polytropic, compressible media with the relation $p/p_b = (\rho/\rho_b)^n$, $n \geq 1$, between pressure p and density ρ , the linear theory was first formulated, and a linear equation was obtained for the perturbation vertical velocity V in terms of x and ψ instead of x and y . The critical speed b was found to be equal to $(p_b/\rho_b)^{1/n}$ for $n > 1$. For $n = 1$, the linear theory yields no information and a nonlinear analysis is especially necessary for this case. If we only formally require the boundedness of $\rho_0 U$ and $\rho_0 V$ where U is the perturbation horizontal velocity and $\rho_0 = \exp(-gy/RT)$, it is found that for $c^2 > RT$, $0 < k^2 < \infty$, non-trivial solutions for the linearized equations exist in the sense defined above, and for $c^2 \leq RT$, $0 \leq k^2 < \infty$, the solution is a trivial one. The only indication of these results is that $(RT)^{1/2}$ may play the role of critical speed in the nonlinear analysis. This is, in fact, the case. However, no unboundedness is involved in the nonlinear formulation, and we find the solitary wave equation as follows:

$$m_0 a'''(\sigma) + m_1 a'(\sigma) a(\sigma) + m_2 a'(\sigma) = 0,$$

$$m_0 = -2(n-1)^2/(2n-1)(3n-2),$$

$$m_1 = -(3n-1)/n,$$

$$m_2 = 1.$$

We see that for $n = 1$ only the trivial solution is possible if the condition at $x = -\infty$ is imposed. It has been shown (Shen, 1967), however, that unsteady waves do exist near the critical speed $(RT)^{1/2}$, and solitary waves will appear if the variation of the horizontal velocity in the state of equilibrium is taken into consideration. Finally, we remark that it is feasible to construct a rigorous proof of the non-existence of solitary waves in an isothermal medium along the lines given in Krasovskii (1960) or Morawetz (1964).

REFERENCES

- Krasovskii, Y. P., 1960: Theory of steady-state waves of large amplitude. *Soviet Physics—Doklady*, **5**, 62–65.
- Morawetz, C. S., 1964: Non-existence of transonic flow past a profile. *Commun. Pure Appl. Math.*, **17**, 357–367.
- Peters, A. S., and J. J. Stoker, 1960: Solitary waves in liquids having non-constant density. *Commun. Pure Appl. Math.*, **13**, 115–164.
- Shen, M. C., 1965: Solitary waves in polytropic compressible media. *Proc. 9th Midwestern Mechanics Conference*, The University of Wisconsin, Madison, (in press).
- , 1967: On long waves with cross wind in an atmosphere. *J. Fluid Mech.* (to be published).