Reply

ABDUL JABBAR ABDULLAH

State University of New York at Albany

12 December 1966

Dr. Wexler is to be commended for calling my attention to three points which seem to remain ambiguous about my theoretical model, and which need further elaboration.

The first point is that the motion of the individual cells within the band must be distinguished from the motion of the band itself. I concur with Dr. Wexler that the two motions must be distinguished from each other. According to my model the band is to be associated with the gravity wave after breaking. The individual cells may be associated with the convection cells that arise from instability triggered by the breaking. It appears, therefore, that although the two phenomena are distinct, yet the appearance of the cells must be organized by the breaking mechanism. The motion of the individual cells themselves may not be the same as that of the band.

The second point is related to the speed with which the band moves relative to the basic current. The velocity of the basic current, as implied in my model, is best approximated by the weighted mean velocity over the depth of the fluid involved in the lower layer. This mean velocity, as may easily be inferred, is usually smaller than the wind velocity above the frictional layers. Hence, even though the band may be moving with a speed which is greater than the mean speed of the basic current, its speed may be comparable

with that of the wind at some upper level. At the same time my theory does not necessarily predict that the bands move faster than the basic current. There is provision in the theory for the generation of two possible families of gravitational waves, one moving with the velocity $u+c$ and the other with the velocity $u-c$. This latter family may be responsible for the slowly moving, or the stationary, bands which are occasionally observed. The main point remains that the bands are not simply drifted with the winds, but they have a different velocity.

The main point in mentioning the radial movement of the individual cells is that the bands grow radially with great rapidity. The apparent movement of the cells was taken to indicate the rate at which the band spreads along its length. The fact that this rate is quite rapid is well borne out by observations. As to the numerical magnitude of this growth, it may be mentioned that my theory leaves three parameters as arbitrary. They may be determined from detailed observations. These parameters are A , c and ω which appear in Eqs. (32) of my paper. By a suitable combination of these parameters it is possible to obtain the right order of magnitude.

It seems to me that some more careful and detailed observations are needed in order to judge the validity of my model.