

The Chemical Composition of the Venus Atmosphere Based on the Data of the Interplanetary Station Venera 4

A. P. VINOGRADOV, U. A. SURKOV AND C. P. FLORENSKY

U.S.S.R. Academy of Sciences, Moscow

(Manuscript received 18 March 1968)

The general goal of the Venera 4 landing probe was the investigation of the physical and chemical characteristics of the Venus atmosphere. The chemical composition of the Venus atmosphere was investigated by gas analyzers which were constructed especially for this purpose. There were eleven gas analyzers which were divided into two groups. The first group consisted of five analyzers and operated at a pressure of approximately 550 mm; the second contained six gas analyzers and operated at a pressure of about 1500 mm.

There were many methods of investigation of the chemical constitution. The most simple and reliable physical-chemical methods were selected. The kinetic and thermodynamic reactions were investigated in the laboratory under conditions similar to those expected on Venus. Two types of analyzers, amplitude and threshold, were used. Some gas analyzers were cylindrical tubes divided into two parts by a thin membrane, one part containing the chemical absorber selected for the investigated component. Both parts were evacuated and sealed. In the operation process both parts were opened and the investigated gas penetrated simultaneously into them. Then they were closed, and the difference of the pressure in the two parts produced by absorption was measured.

For analyzing minor constituents more sensitive methods were used. These included measurements of the electrical resistance of the specially selected chemical absorber, the heat conductivity, etc. Investigations were made of the content of CO₂, N₂, O₂ and H₂O. Due to the previous uncertainty of the chemical composition of the Venus atmosphere, the sensors were designed with a large range.

The results of the investigation of the chemical composition of the Venus atmosphere are shown in Table 1. The presence of CO₂ is confirmed by four sensors. We conclude that the content of CO₂ in the Venus atmosphere is not less than 90%.

Both nitrogen sensors show the absence of a significant amount of this element in the Venus atmosphere. The first sensor gives an upper limit of 7% and the second sensor an upper limit of 2.5%. However, the second sensor is less accurate. Adopting a conservative point of view, we take 7% as the upper limit of nitrogen.

The oxygen content lies between the two limits obtained by two threshold sensors utilizing various methods. In the first sensor the tungsten filament was adjusted at the limit of about 3 mm partial pressure O₂, corresponding to 0.5% of oxygen at 550 mm pressure. This filament was burned out immediately. The second sensor was based on the selected absorption of O₂ by sublimated phosphorous vapor, in which, as a consequence of this reaction, P₂O₅ absorbs water vapor. This sensor was adjusted at 1.6% of (O₂+H₂O) and no overloading occurred.

The water content was estimated by three sensors. The first sensor, operating on the principle of the variation of the resistivity of P₂O₅, detected the presence of H₂O in an amount larger than 0.65 mg liter⁻¹. This corresponded to 0.1% H₂O for a condensation temperature higher than 22°C. The second sensor, based on a similar reaction, also showed the presence of H₂O to be larger than 0.65 mg liter⁻¹. The third sensor, operating on the pressure difference principle (absorption by CaCl₂), showed that the partial pressure of water vapor is smaller than 11 mm (<0.7%).

Summarizing, we have adopted the following composition (in volume per cent) of the Venus atmosphere:

CO ₂	90±10% (probably >90%)
O ₂	>0.4%, <1.6% (probably ~1%)
N ₂	<7% (probably <2.5%)
H ₂ O	from 1–8 mg liter ⁻¹




In general, the Venus atmosphere is oxidizing and it corresponds to the greenhouse model.

It is very interesting to compare the present atmospheres of Venus and earth. The overall amounts of degassing products are approximately the same. But on earth, carbon is bound by carbonate sedimentary rocks, and its amount is equal to about 2·10²³ gm CO₂. Therefore, if one sublimates all the CO₂ from the carbonates and hydrosphere of earth to its atmosphere, the surface pressures would be increased by a factor of about 40. In other words, in order to obtain the Venus pressure of 20 atm, it is enough to set free about half of the CO₂

TABLE 1. Results of chemical analyses of the Venus atmosphere.

Analysis conditions	Component under determination	Type of cell	Method of measurement	Limits on measurements (% by volume)	Results of measurements (% by volume)
$H \sim 26 \pm 1$ km	CO ₂	Threshold	Heat conductivity	Threshold 1	>1
$P_0 \sim 550$ mm	CO ₂	Amplitude	KOH absorption	7-100	90±10
$t \sim 25 \pm 10$ C (first group of analyzers)	N ₂	Amplitude	Preliminary absorption of O ₂ , CO ₂ at 1000C	7-100	<7
	O ₂	Threshold	W 800C outburning	Threshold 0.4	>0.4
	H ₂ O	Threshold	Electroconductivity of moistened P ₂ O ₅	Threshold 0.1	>0.1
$H \sim 19 \pm 1$ km	CO ₂	Amplitude	KOH absorption	2-30	>30
$P_0 \sim 1500$ mm	CO ₂	Threshold	KOH absorption	Threshold 1	>1
$t \sim 90 \pm 10$ C (second group of analyzers)	N ₂	Amplitude	Preliminary absorption of O ₂ , CO ₂ at 1000C	2.5-50	≤2.5
	O ₂ (+H ₂ O)	Threshold	P vaporizing	Threshold 1.6	<1.6
	H ₂ O	Threshold	Electroconductivity of moistened P ₂ O ₅	Threshold 0.05	>0.05
	H ₂ O	Amplitude	CaCl ₂ absorption	Threshold 0.7	<0.7

bound in the terrestrial rocks. This process may be imagined by chemical reactions of the following types:



These reactions develop from the left to the right side if the temperature is about 300C.

The only source of nitrogen in the Venus atmosphere may be NH₄. On earth NH₄ is degassing by volcanic action in the form of NH₄Cl, a stable salt sublimating only at 350C. On Venus there is much more CO₂ and (NH₄)₂CO₃ is therefore produced. This salt is less stable, decomposing at a temperature of 58C to form NH₃ which is then oxidized by oxygen to N₂, the oxygen being produced by photodissociation of water.

Let us discuss briefly the question of the origin of the Venus atmosphere. On planets of equal sizes endogenic processes, i.e., melting of the crust and degassing, are similar and develop by a zone melting mechanism. The subsequent evolution depends on the distance from the sun, the mass of the planet, and other exogenic conditions.

Let us draw some conclusions. Since Venus is closer to the sun than earth, its effective temperature as a naked planet would be higher by 50C, leading to the

emission into the atmosphere of a large amount of CO₂ and H₂O, and to self-heating of the atmosphere. The oxygen is produced by photodissociation of water and is absorbed by surface rocks. The other product of photodissociation is CO, but this recombines to CO₂ with oxygen formed by H₂O photodissociation. The hydrogen was dissipated due to the high temperature and the absence of a magnetic field. This process led to the self-heating of the atmosphere and to an increase of the greenhouse effect. Some contribution to the thermal balance of the planet is due now to internal heating. When the temperature of the surface reaches approximately 250-300C, many carbonates react with silicates, releasing large amounts of CO₂. The conditions on Venus obtained by Venera 4 exclude the existence of a hydrosphere because water boils at a temperature higher than 100C.

Therefore, endogenic processes on both Venus and earth were probably similar. But the exogenic processes depending primarily on the distance from the sun were different. This led to the formation of different atmospheres on these planets, the Venus atmosphere becoming very heavy. Such conditions and an aggressive state of the atmosphere led to a deep denudation of the surface rocks. The intensive circulation probably produced the flattening of the planet's surface.