

Wavelength Dependence of the Absorption of Black Carbon Particles: Predictions and Results from the TARFOX Experiment and Implications for the Aerosol Single Scattering Albedo

ROBERT W. BERGSTROM

Bay Area Environmental Research Institute, Sonoma, California

PHILIP B. RUSSELL

NASA Ames Research Center, Moffett Field, California

PHILLIP HIGNETT

Meteorological Support Group, Ministry of Defense, London, United Kingdom

(Manuscript received 19 December 2000, in final form 23 May 2001)

ABSTRACT

Measurements are presented of the wavelength dependence of the aerosol absorption coefficient taken during the Tropical Aerosol Radiative Forcing Observational Experiment (TARFOX) over the northern Atlantic. The data show an approximate λ^{-1} variation between 0.40 and 1.0 μm . The theoretical basis of the wavelength variation of the absorption of solar radiation by elemental carbon [or black carbon (BC)] is explored. For a wavelength independent refractive index the small particle absorption limit simplifies to a λ^{-1} variation in relatively good agreement with the data. This result implies that the refractive indices of BC were relatively constant in this wavelength region, in agreement with much of the data on refractive indices of BC. However, the result does not indicate the magnitude of the refractive indices.

The implications of the wavelength dependence of BC absorption for the spectral behavior of the aerosol single scattering albedo are discussed. It is shown that the single scattering albedo for a mixture of BC and nonabsorbing material decreases with wavelength in the solar spectrum (i.e., the percentage amount of absorption increases). This decrease in the single scattering albedo with wavelength for black carbon mixtures is different from the increase in single scattering albedo for most mineral aerosols (dusts). This indicates that, if generally true, the spectral variation of the single scattering albedo can be used to distinguish aerosol types. It also highlights the importance of measurements of the spectral variation of the aerosol absorption coefficient and single scattering albedo.

1. Introduction

Absorption of solar radiation by black carbon (BC) particles is important in understanding the effects of atmospheric particles on climate. Particles with no absorption have a negative (cooling) forcing while particles with substantial absorption can have a positive (warming) forcing. As Heintzenberg et al. (1997) noted, "Even though BC constitutes only a few percent of the aerosol mass, it can have a significant positive forcing." Simple calculations show that a small amount of BC can change a negative aerosol forcing to a positive aerosol forcing. Jacobson recently found that the magnitude of the direct radiative forcing from black carbon itself

exceeds that due to CH_4 and may be the second most important component of global warming after CO_2 in terms of direct forcing (Jacobson 2001b). In addition to the direct climate effect, the Indian Ocean Experiment recently showed that BC particles were apparently responsible for reduced cloudiness over the Indian Ocean (Ackerman et al. 2000).

Black carbon particles have received increased interest recently since Hansen et al. (2000) indicate that it is perhaps more cost effective to control black carbon particles than CO_2 at the present time. However, before embarking on a large-scale control strategy for BC particles, it is important to fully understand their radiative properties and effects.

Black carbon particles enter the atmosphere as a result of incomplete combustion processes, whether from motor vehicles, industrial flames, or biomass burning. If the flame temperature is too low or there is a limited

Corresponding author address: Robert W. Bergstrom, Bay Area Environmental Research Institute, 645 Anthony Ct., Sonoma, CA 95476.
E-mail: bergstro@sky.arc.nasa.gov




FIG. 1. Aerosol absorption coefficients as a function of wavelength for three TARFOX days: (a) 21, (b) 25, and (c) 27 July.

supply of oxygen, not all of the fuel is converted to carbon oxides and carbon soot results. As combustion is ubiquitous, carbon soot is found virtually everywhere [e.g., off the eastern coast of United States (Novakov et al. 1997); off the western coast of Europe (Putaud et al. 2000) and Novakov et al. 2000); off the Indian Peninsula (Ramanathan et al. 1996); in China (Chameides et al. 1999); in Brazil (Kaufman et al. 1998); over the Pacific Ocean (Kaneyasu and Murayama 2000), and in

remote areas (Bodhaine 1995)]. Global simulations of carbonaceous aerosols show a worldwide distribution of BC with maxima over the heavily industrialized areas and biomass burning regions (Liousse et al. 1996).

In this paper we present measurements of the wavelength dependence of the aerosol absorption coefficient due to BC obtained during the Tropical Aerosol Radiative Forcing Observational Experiment (TARFOX) experiment. We then discuss the effects of particle size




FIG. 1. (Continued)

and refractive indices on the absorption coefficient of BC and, in particular, the wavelength dependence of the absorption coefficient. The data show that the absorption seen in the TARFOX experiment has an approximate λ^{-1} dependence that is in good agreement with the small particle limit for BC particles for a constant refractive index.

We also briefly discuss the implications of the results in terms of the single scattering albedo of atmospheric aerosols. We show that for situations where the absorption is due to BC, the single scattering albedo of the aerosol mixture should decrease with wavelength. This is in contrast to most mineral aerosols (dusts) where the single scattering albedo usually increases with wavelength (Sokolik and Toon 1997, 1999).

2. Measurements of the aerosol absorption coefficient during the TARFOX experiment

The TARFOX field program was carried out off the eastern coast of the United States in the summer of 1996 (Russell et al. 1999a). Samples of the aerosol were obtained aboard the Met Office Hercules Aircraft (C130) on three particular days: 21, 25, and 27 July 1996. The filters were exposed during the level passes when the solar radiation values were measured (Hignett et al. 1999). The samples were taken back to the Met Office laboratory and the absorption coefficients at wavelengths between 0.45 and 1.0 μm were measured using the integrating sandwich method (Clarke 1982; Foot and Kilsby 1989).

a. Integrating sandwich method

The integrating sandwich technique was developed to minimize the influence of the scattering properties of the measured material on the absorption determination (Clarke 1982). The material is placed between two high-reflectance diffusing wafers and the resultant attenuation of light is measured. The method is discussed in detail elsewhere (Clarke 1982; Foot and Kilsby 1989; Horvath 1993; Heintzenberg et al. 1997). Clarke (1982) estimated the error to be $\pm 25\%$.

The major uncertainties in the method arise from the additional scattering in the layer from other particles (Horvath 1997b) and knowing how many passes the light makes through the media. This causes additional apparent absorption and leads to an overestimation of the absorption coefficient. However, in this study we are concerned with the wavelength dependence of the absorption coefficient and not the absolute magnitude of the absorption coefficient. We have included the standard deviation of the individual measurements as a measure of the intrinsic variability similar to Hignett et al. (1999) who showed some of this data in their Fig. 4.

b. Absorption coefficient results

The measured absorption coefficients are shown in Fig. 1. Only the data collected when the airplane was in the aerosol layers are plotted. The relative standard deviations are larger at longer wavelengths and for smaller absorption coefficients. To isolate the wavelength dependence, the data are normalized by the ab-




FIG. 2. Aerosol absorption coefficients as a function of wavelength normalized by the value at (a) $0.5 \mu m$, (b) 25 July, and (c) 27 July.

sorption coefficient at $0.55 \mu m$ and shown in Fig. 2. The results show clearly the decrease in absorption coefficient with wavelength. As seen, the wavelength variation over the 3 days is roughly the same. The absorption decreases quite quickly with wavelength between 0.4 and $0.6 \mu m$ and then levels off at the longer

wavelengths. There is a relatively large spread in the value at $1.0 \mu m$.

The aerosol conditions on two of the TARFOX days were quite similar (25 and 27 July), but the conditions on 21 July were very different (low optical depth on 21 July but very hazy on 25 and 27 July. The fact that the




FIG. 2. (Continued)

wavelength dependence is very similar on all 3 days is notable and possibly indicates that the absorbing material has a common origin.

The aerosol absorption observed in TARFOX was attributed to BC (Novakov et al. 1997) since the correlation of the absorption coefficient with measured carbonaceous aerosol amount was quite high ($r^2 = 0.69$) and the observed absorption coefficient indicated roughly 10% of the carbonaceous aerosol was BC. As we do not have detailed chemical information on the aerosol encountered in the TARFOX experiment, it is possible that there were other absorbing hydrocarbons present. Jacobson has pointed out that other materials such as benzene absorb in the UV in a similar manner as BC particles (Jacobson 2001a). However, for simplicity, we will refer to the absorption measured in the samples as absorption due to BC particles.

Since the samples were collected on filters, the absorption measurements are for the dry aerosol only. We do not consider here the effects of water or other coatings on the particles (Ackerman and Toon 1981; Fuller et al. 1999; Redemann et al. 2001). While important for the absolute magnitude of the absorption, these effects probably have less influence on the wavelength dependence of the absorption coefficient. Jacobson (2000) shows some sample calculations for externally mixed, internally mixed, and shell-core configurations of BC. While the magnitudes of the calculated absorption coefficients differ, the wavelength dependence is very sim-

ilar for all three cases for the spectral region with which we are concerned (0.45–1.0 μm).

3. Calculations of the absorption coefficient of BC

The optical properties of BC have been studied in the context of industrial flames (Dalzell and Sarofim 1969; Lee and Tien 1981; Chang and Charalampopoulos 1990; Stagg and Charalampopoulos 1993; paint pigments (Medalia and Richards 1972), urban atmospheres (Horvath 1993; Horvath et al. 1997), the nuclear winter debate (Turco et al. 1991; Golitsyn et al. 1988; Penner 1986), and more recently, biomass burning (Kaufman et al. 1999; Reid et al. 1998).

Fuller et al. (1999) have recently explored in some detail the effects of various sizes, shapes, and configurations on the extinction of light by BC particles. They considered the effects of material composition and morphology of soot and the electrostatics of spherules agglomerated into loose (ramiform) aggregates. They found a wide variation in the mass absorption coefficient at 0.55 μm depending upon the variations in optical constants. Their results suggest the importance of determining the physical state of the soot particles and the immediate environment when ascribing characteristic values for the absorption and scattering efficiencies.

However, Fuller et al. (1999) did not directly address the wavelength dependence of the absorption by BC particles and our results suggest that a less detailed anal-

ysis may be sufficient for explanation of the wavelength dependence of the absorption. Therefore, we discuss here only the small particle limit and the effect of the refractive indices on the absorption coefficient.

a. Size effects

In general, the radiative properties of atmospheric aerosols are a strong function of size. Electron microscope analyses show that most of the total BC mass is composed of fractal agglomerates made up of primary spherules with a diameter of 0.03–0.05 μm (Seinfeld and Pandis 1998, Fig. 13.1; Martins et al. 1998). For very small spherical absorbing particles (roughly $r < 0.01 \mu\text{m}$ for $\lambda = 0.55 \mu\text{m}$) the absorption per mass (specific absorption or mass absorption coefficient) becomes independent of size (van de Hulst 1957; Bergstrom 1973; Bohren and Huffman 1983; Horvath 1993; Colbeck et al. 1989; Martins et al. 1998). This size-independent specific absorption coefficient relationship corresponds to the so-called Rayleigh absorption limit of

$$\begin{aligned} \kappa &= (\text{constant}/\rho\lambda)\text{Im}[(m - 1/m - 2)] \\ &= (\text{constant}/\rho\lambda)nk/[(n^2 - k^2 + 2)^2 + (2nk)^2], \quad (1) \end{aligned}$$

where κ is the specific or mass absorption coefficient, m is the complex index of refraction ($m = n - ik$), and ρ is the density of the particle. An important result of Eq. (1) is that for small particles and a constant refractive index the wavelength dependence of the mass absorption coefficient is simply λ^{-1} .

Equation (1) is only strictly applicable to absorption by a spherical particle with a uniform index of refraction. Berry and Percival (1986) developed a fractal theory to describe the absorption by the BC fractal agglomerates. This theory also predicts that the specific absorption of these agglomerates should not be a strong function of size. [However, it is a strong function of the optical constants and the physical state of the particle (Fuller et al. 1999).] Experimental results are in good agreement with the fractal model predictions (Colbeck 1989; Petzold et al. 1997, Fig. 5).

Data from Horvath (1993) show a decrease in the specific absorption with size, but Petzold et al. (1997) argue that these larger black carbon particles that are not fractal clusters (and hence have less specific absorption) are a small amount of the total BC mass.

For very large spherical particles the absorption (again, for constant refractive index) becomes wavelength-independent (Bergstrom 1973). Thus we can infer that the wavelength dependence of the absorption coefficient of a BC particle with a constant refractive index should be about λ^{-1} to $\lambda^{-1.3}$ or less. (The Ångström exponent for absorption by a BC particle with effective radius of $\sim 0.02 \mu\text{m}$ is 1.3 at $0.55 \mu\text{m}$.) This result is independent of what the absolute value of the refractive index is (or the magnitude of the specific absorption).

b. Refractive index of BC

It is well established that the refractive indices of BC particles vary greatly depending upon the details of the combustion process (fuel type, flame temperature, etc.). There is a wide variation in the reported value of refractive indices of BC (Twitty and Weinman 1971, Fig. 1; Colbeck et al. 1989, Table 1; Fuller et al. 1999). Bohren (1986) noted that there is about a factor of 5 variation in the imaginary part of the index of refraction for what is called “soot.”

The wide variation in the measured refractive indices most likely comes from the amount of other hydrocarbons contained in the particles. Batten (1985) notes that the imaginary part of the refractive index is inversely correlated with the H/C ratio. In other words, the closer the particle is to pure carbon (i.e., graphite) the larger the refractive indices. Also, as mentioned above, other hydrocarbon compounds may absorb in a similar manner to BC further complicating the measurement of the refractive indices.

Undoubtedly, there is a large variation in the absorption coefficient of the carbon particles that exist in the atmosphere depending upon their sources and the changes that occur while in the atmosphere (coatings, chemical reactions). For example, Fuller et al. (1999) classified BC into five different categories with different refractive indices and densities for each group.

To explore the effect of the refractive index on the wavelength dependence of the absorption, we show two different reported values for the spectral variation of the refractive indices of flame soots (Lee and Tien 1981; Chang and Charalampopoulos 1990) in Fig. 3. Both studies represented the molecular structure of the soot as graphite (i.e., three bound electrons and one free electron). Then they computed values for the damped harmonic oscillator, etc., in the Kramers–Kronig formula (Bohren and Huffman 1983). They fit their transmission data and came up with the index of refraction as a function of wavelength for the particular type of flame soot being studied.

The wavelength dependence of the refractive indices of Lee and Tien is larger than that for the Chang and Charalampopoulos values. The imaginary part of the Lee and Tien refractive index varies from 1.0 at $0.3 \mu\text{m}$ to a minimum of 0.38 at $0.5 \mu\text{m}$ and then increases to 0.9 at $1.0 \mu\text{m}$. By contrast, the Chang and Charalampopoulos refractive indices have relatively little wavelength variation similar to other estimates of the refractive indices of flame soot (Dalzell and Sarofim 1969; Bergstrom 1972). It should be emphasized that different flame soots (acetylene, propane, diesel fuel, etc.) have different values for the refractive indices. These two choices of the refractive indices illustrate the range in the wavelength dependence in the region of 0.3 to $1.0 \mu\text{m}$ from relatively constant to varying by more than a factor of 2.

The wavelength variation in the Lee and Tien re-




FIG. 3. Refractive indices for flame soots from Lee and Tien (1981) and Chang and Charalampopoulos (1990).

refractive indices results from their expression for the temperature dependence of the electron damping coefficients. Lee and Tien show this temperature dependence in their Fig. 5 where the variation of the imaginary index with wavelength increases significantly as the temperature decreases. Stagg and Charalampopoulos (1993) indicated that there was no significant temperature variation in their determined refractive indices of flame soot. They question the use of the temperature dependence of the electron damping coefficients used by Lee and Tien (1981).

In the UV, the increase in refractive indices is termed a “glassy carbon” resonance by Chang and Charalampopoulos (see also Bruce et al. 1991). This absorption could be important for photochemical reactions.

At the longer wavelengths (greater than 1.0 μm), the refractive indices of BC also increase with wavelength. This is due to properties of the free electron and becomes important for the thermal emission properties of flame soot (Lee and Tien 1981). As mentioned above, the absolute magnitude of refractive indices varies greatly depending upon the type of fuel and specific combustion processes.

4. Comparison of predicted and measured absorption and discussion

a. Comparison to the TARFOX measurements

Using the values of the wavelength dependence of the refractive indices discussed above we computed the value of specific absorption from Eq. (1) and plotted

the relative values in Fig. 4. Also plotted is a simple λ^{-1} curve that represents a constant refractive index.

To describe the goodness of fit of the computed values to the data, we use two measures: reduced chi-squared χ_v^2 and the root-mean-square difference between computed and measured values. Here, χ_v^2 (e.g., Bevington and Robinson 1992) is proportional to the weighted sum of squares of the difference between each data point and the corresponding computed value; the weight for each difference is the inverse square of its measurement error bar. As noted by Bevington and Robinson (1992), if the calculated curve is a good approximation to the data, χ_v^2 should be approximately unity or less. Increasingly larger values indicate increasingly poorer fit. For the Chang and Charalampopoulos refractive indices χ_v^2 is 0.79, 1.97 for the constant refractive index (λ^{-1}), and 38.23 for the Lee and Tien refractive indices. Corresponding rms differences between computed and measured values are 0.085, 0.12, and 0.49.

As shown in Fig. 4 the λ^{-1} curve (the small particle limit with constant index of refraction) fits the data within nearly all the uncertainty bars ($\chi_v^2 = 1.97$). The wavelength dependence of the absorption coefficient using the Chang and Charalampopoulos refractive indices is very similar to that obtained using a constant index of refraction and in fact fits the data somewhat better ($\chi_v^2 = 0.79$). Use of the Lee and Tien refractive indices, however, overestimates the relative absorption at wavelengths longer than 0.6 μm and gives a very poor fit overall ($\chi_v^2 = 38$). However, the Lee and Tien refractive indices do capture the flattening out of the absorption at longer wavelengths.

Our results indicate that the simple constant index of refraction–small particle limit is a relatively good representation of the wavelength dependence of the aerosol absorption by BC observed in the TARFOX field campaign. This implies that the refractive indices of BC were relatively constant in agreement with much of the data on BC (such as Chang and Charalampopoulos 1990).

b. Comparison to other measurements of BC specific absorption

There is only a limited amount of data on the spectral absorption coefficient of BC particles (as compared to the amount of data on the specific absorption in the midvisible, 0.50–0.55 μm). Bruce et al. (1991) compared the available spectral data for the absorption and scattering properties of diesel soots. They found that there was an approximate λ^{-1} dependence at selected laser wavelengths from 0.488 μm to 0.857 μm .

Horvath et al. (1997) found a wavelength dependence of $\lambda^{-0.92}$ for the aerosol absorption from their studies in Santiago, Chile. The absorption was most likely due to BC particles since the measurements were made in an urban area.

Moosmuller et al. (1998) measured the absorption coefficient of atmospheric aerosols near Denver, Colorado, at 0.532 and 0.685 μm with a photoacoustic instrument. They compared their results with other measurements that indicated a wavelength dependence of the absorption of $\lambda^{-2.7}$. However, there were only a few measurements included and from different sources. Also shown in Fig. 4 is the $\lambda^{-2.7}$ curve. It clearly underestimates the TARFOX absorption for wavelengths greater than about 0.6 μm .¹

Dubovik et al. (1998) inferred the aerosol absorption optical depth of the atmosphere from sky radiance data for areas that have significant biomass burning. Their results show that for one year the wavelength dependence was λ^{-1} but for two other years the wavelength dependence was greater than λ^{-2} . Dubovik et al. (1998) argue that the cause of the increase in wavelength dependence of the absorption optical depth is most likely due to the wavelength dependence of the imaginary refractive indices (assuming the small particle limit is valid). However, as discussed above, the data from flame soots do not indicate that the imaginary index decreases enough from 0.3–1.0 μm to account for a λ^{-2} relationship. It is more likely that mineral (dusts) in the atmospheric aerosol caused the more rapid decrease in absorption with wavelength (see below). Dubovik et al. (1998) noted that iron oxide has refractive indices that

would cause such an increase in the wavelength dependence. Another possibility is that the biomass burning aerosol is sufficiently complicated (see Martins et al. 1998) that at times the λ^{-1} relationship is not applicable.

While combustion processes vary widely from place to place (e.g., diesel exhaust to biomass burning), it would appear that the small BC particles may have a common structure and perhaps a common wavelength dependence of the absorption. Such a result would be very useful for the estimate of the climatic impact of BC particles.

c. Implications for the single scattering albedo of the atmospheric aerosol

The single scattering albedo is a critical parameter for determining the effect of aerosols on the global temperature. It is defined as the ratio of the scattering to the extinction; that is,

$$\omega = \sigma/\beta,$$

where σ is the scattering coefficient and β is the extinction coefficient. Or expressed in terms of the absorption coefficient

$$\omega = 1 - (\kappa/\beta).$$

If there is a mixture of small BC (or BC-like material) particles and other nonabsorbing particles (such as sulfates), then the nonabsorbing particles will dominate the scattering and the BC, the absorption. If the extinction also follows a “power law,” then

$$\omega = 1 - (\kappa_o/\beta_o)(\lambda/\lambda_o)^{-1+\alpha}, \quad (2)$$

where α is the Ångström exponent.

For atmospheric aerosols that do not contain large particles (i.e., dust) α usually is greater than 1, so the single scattering albedo of the mixture should decrease with wavelength. This is because the scattering coefficient usually decreases with wavelength faster than the absorption coefficient (Bergstrom 1973). As the scattering coefficient goes to zero, the extinction coefficient equals the absorption coefficient and the single scattering albedo is zero. This result is also discussed by Dubovik et al. (1998).

However, for BC particle clusters, the single scattering albedo does not drop to zero at visible wavelengths. Bruce et al. (1991) show that the single scattering albedo for BC clusters is about 0.3 for 0.554 μm . This apparently is because the scattering coefficient of BC clusters does not follow the small particle limit (which is Rayleigh scattering). Berry and Percival’s fractal theory indicates that the scattering cross section for the cluster is larger than that for the sum of the spherules (unlike the absorption coefficient). However, Bruce et al. (1991) show that the single scattering albedo for carbon particles gets very small at 100 μm .

Estimates of the TARFOX aerosol single scattering albedo based on model calculations show a decrease

¹ Note added in proof: Bond (2001) investigated possible explanations for the spectral dependence of the absorption coefficient of carbon particles from coal combustion measured to be between λ^{-1} and $\lambda^{-2.5}$. She concluded that a spectrally dependent imaginary refractive index is the most plausible explanation.




FIG. 4. Normalized aerosol absorption coefficients as a function of wavelength. Also shown is a λ^{-1} curve and calculations using Eq. (1) with the Lee and Tein, and Chang and Charalampopoulos refractive indices. The chi-squared value for the Chang and Charalampopoulos refractive indices is 0.79 (rms error 0.085), 1.97 for the constant refractive index (rms error 0.12), and 38.23 for the Lee and Tien refractive indices (rms error 0.49).

with wavelength (Hignett et al. 1999; Russell et al. 1999b). Estimates of the effects of water growth on the particles do not change the decrease with wavelength [of course, the single scattering albedo increases with relative humidity due to the water taken up by the particles (see Hignett et al. 1999, Fig. 3)].

Recently, the Smoke, Clouds and Radiation-Brazil program measured the optical properties of the aerosol from biomass burning in Brazil (Kaufman et al. 1998). The biomass burning BC particles have a very complicated structure and are difficult to describe simply (see Martins et al. 1998, Fig. 5). However, Reid and Hobbs (1998) found that the single scattering albedo decreased with wavelength, as did Eck et al. (1998).

In contrast to BC, large particles (dust or sea salt) usually have an Ångström coefficient less than one [or relatively constant optical depth in the solar spectrum (Sokolik and Toon 1997)]. In addition, the small particle absorption limit of Eq. (1) is not valid for these large particles. However, it is useful to compare the wavelength dependence of the single scattering albedo of mineral aerosols and BC. Sokolik and Toon (1997) show single scattering albedos for various dust measurements and Sokolik and Toon (1999) present the refractive indices and radiative properties for a number of mineralogical components. Their results indicate that the single scattering albedo for dust increases with wavelength in the range of 0.3–1.0 μm . This is due to significant absorption in the UV and visible wavelengths less than

about 0.6 μm due to electronic bands and less absorption at longer solar wavelengths. Because the extinction coefficient is relatively constant with wavelength, the change in single scattering albedo is determined simply by the absorption coefficient spectral behavior.

If the difference in wavelength behavior between BC and dust is generally true, it means that the wavelength dependence of the single scattering albedo could be used to differentiate between dust absorption and BC absorption.

5. Conclusions

We have analyzed the wavelength dependence of the absorption coefficient measurements of aerosol collected during the TARFOX field campaign. The results show a decrease in the absorption of λ^{-1} that is consistent with the absorption of small particle BC. This result implies that the refractive indices of BC in this spectral region were relatively constant in agreement with much of the data on the refractive indices of BC.

We show that for atmospheric aerosol mixtures, where the absorption is due to BC and the extinction wavelength dependence is less than λ^{-1} , the single scattering albedo should decrease with wavelength in the region of 0.3–1.0 μm . This is in agreement with the limited data available. It contrasts with the increase in the single scattering albedo for mineralogical dusts and indicates the need for more careful measurements of the wave-

length dependence of the aerosol absorption and the single scattering albedo.

Acknowledgments. This research was conducted as part of the Global Aerosol Climatology Program (GACP) and the Tropospheric Aerosol Radiative Forcing Observational Experiment. TARFOX was part of the International Global Atmospheric Chemistry (IGAC) core project of the International Geosphere-Biosphere Programme (IGBP), and part of the IGAC Focus on Atmospheric Aerosols (FAA). GACP is supported by the Radiation Science Program Office of the U.S. National Aeronautics and Space Administration. Other support was provided by the Aerosol Program of the U.S. National Oceanographic and Atmospheric Administration Office of Global Programs.

REFERENCES

- Ackerman, A. S., O. B. Toon, D. E. Stevens, A. J. Heymsfield, V. Ramanathan, and E. J. Welton, 2000: Reduction of tropical cloudiness by soot. *Science*, **288**, 1042–1047.
- Ackerman, T. P., and O. B. Toon, 1981: Absorption of visible radiation in atmosphere containing mixtures of absorbing and nonabsorbing particles. *Appl. Opt.*, **20**, 3661–3668.
- Batten, C. E., 1985: Spectral optical constants of soots from polarized angular reflectance measurements. *Appl. Opt.*, **24**, 1193–1199.
- Bergstrom, R. W., 1972: Predictions of the spectral absorption and extinction coefficients of an urban aerosol model. *Atmos. Environ.*, **6**, 247–258.
- , 1973: Extinction and absorption coefficients of the atmospheric aerosol as a function of particle size. *Beitr. Phys. Atmos.*, **46**, 223–234.
- Berry, M. V., and I. C. Percival, 1986: Optics of fractal clusters such as smoke. *Opt. Acta*, **33**, 577–591.
- Bevington, P. R., and D. K. Robinson, 1992: *Data Reduction and Error Analysis for the Physical Sciences*. 2d ed. McGraw-Hill, 328 pp.
- Bodhaine, B. A., 1995: Aerosol absorption measurements at Barrow, Mauna Loa and the South Pole. *J. Geophys. Res.*, **100**, 8967–8975.
- Bohren, C. F., 1986: Applicability of effective-medium theories to problems of scattering and absorption by nonhomogeneous atmospheric particles. *J. Atmos. Sci.*, **43**, 469–475.
- , and D. R. Huffman, 1983: *Absorption and Scattering of Light by Small Particles*. John Wiley, 232 pp.
- Bond, T. C., 2001: Spectral dependence of visible light absorption by carbonaceous particles emitted from coal combustion. *Geophys. Res. Lett.*, **28**, 4075–4078.
- Bruce, C. W., T. F. Stromberg, K. P. Gurton, and J. B. Mozer, 1991: Trans-spectral absorption and scattering of electromagnetic radiation by diesel soot. *Appl. Opt.*, **30**, 1537–1546.
- Chamiedes, W. L., and Coauthors, 1999: Case study of the effects of atmospheric aerosols and regional haze on agriculture: An opportunity to enhance crop yields in China through emission controls? *Proc. Natl. Acad. Sci.*, **26**, 13 626–13 633.
- Chang, H., and T. T. Charalampopoulos, 1990: Determination of the wavelength dependence of refractive indices of flame soot. *Proc. Roy. Soc. London*, **430A**, 577–591.
- Clarke, A. D., 1982: Integrating sandwich: A new method of measurement of the light absorption coefficient for atmospheric particles. *Appl. Opt.*, **21**, 3011–3020.
- Colbeck, I., E. J. Hardman, and R. M. Harrison, 1989: Optical and dynamic properties of fractal clusters of carbonaceous smoke. *J. Aerosol Sci.*, **20**, 765–774.
- Dalzell, W. H., and A. F. Sarofim, 1969: Optical constants of soot and their application to heat-flux calculations. *J. Heat Transfer*, **91**, 100–104.
- Dubovik, O., B. N. Holben, Y. J. Kaufman, M. Yamasoe, A. Smirnov, D. Tanre, and I. Slutsker, 1998: Single scattering albedo of smoke retrieved from the sky radiance and solar transmittance measured from ground. *J. Geophys. Res.*, **103**, 31 903–31 923.
- Eck, T. F., B. N. Holben, I. Slutsker, and A. Setzer, 1998: Measurements of irradiance attenuation and estimation of aerosol single scattering albedo for biomass burning aerosol in Amazonia. *J. Geophys. Res.*, **103**, 31 865–31 878.
- Foot, J. S., and C. G. Kilsby, 1989: Aerosol absorption: An inter-comparison of techniques and spectral observations. *Atmos. Environ.*, **23**, 489–495.
- Fuller, K. A., W. C. Malm, and S. M. Kreidenweis, 1999: Effects of mixing on extinction by carbonaceous particles. *J. Geophys. Res.*, **104**, 15 941–15 954.
- Golitsyn, G. S., A. K. H. Shukurov, A. S. Ginzburg, A. G. Sutugin, and A. V. Andronova, 1988: Combined investigation of microphysical and optical properties of a smoke aerosol. *Atmos. Oceanic Phys.*, **24**, 163–168.
- Hansen, J., M. Sato, R. Ruedy, A. Lacis, and V. Oinas, 2000: Global warming in the 21st century: An alternative scenario. *Proc. Natl. Acad. Sci.*, **97**, 9875–9880.
- Heintzenberg, J., R. J. Charlson, A. D. Clarke, C. Lioussse, V. Ramanamamy, K. P. Shine, M. Wendish, and G. Helas, 1997: Measurements and modelling of aerosol single scattering albedo: Progress, problems and prospects. *Beitr. Phys. Atmos.*, **70**, 249–263.
- Hignett, P., J. P. Taylor, P. N. Francis, and M. D. Glew, 1999: Comparison of observed and modeled direct aerosol forcing during TARFOX. *J. Geophys. Res.*, **104**, 2279–2287.
- Horvath, H., 1993: Atmospheric light absorption—A review. *Atmos. Environ.*, **27A**, 319–326.
- , 1997a: Comparison of the light absorption coefficient and carbon measures for remote aerosols: An independent analysis of data from the IMPROVE network I and II: Discussion. *Atmos. Environ.*, **31**, 2885–2887.
- , 1997b: Systematic deviations of light absorption measurements by filter transmission methods. *J. Aerosol Sci.*, **28**, 533–536.
- , and G. Metzger, 1990: Experimental determination of the accuracy of light absorption measurement with the integrating plate method. *J. Aerosol Sci.*, **21**, 525–528.
- Jacobson, M. Z., 2000: A physically based treatment of elemental carbon optics: Implications for global direct forcing of aerosols. *Geophys. Res. Lett.*, **27**, 217–220.
- , 2001a: Global direct radiative forcing due to multicomponent anthropogenic and natural aerosols. *J. Geophys. Res.*, **106**, 1551–1568.
- , 2001b: Strong radiative heating due to the mixing state of black carbon in atmospheric aerosols. *Nature*, **409**, 695–697.
- Kaneyasu, N., and S. Murayama, 2000: High concentrations of black carbon over middle latitudes in the Pacific Ocean. *J. Geophys. Res.*, **105**, 19 881–19 890.
- Kaufman, Y. J., and Coauthors, 1998: Smoke, clouds, and radiation-Brazil (SCAR-B) experiment. *J. Geophys. Res.*, **103** (D24), 31 783–31 808.
- Lee, S. C., and C. L. Tien, 1981: Optical constants of soot in hydrocarbon flames. *Proc. 18th Int. Symp. on Combustion*, Pittsburgh, PA, The Combustion Institute.
- Lioussse, C., J. E. Penner, C. Chuang, J. J. Walton, H. Eddleman, and H. Cachier, 1996: A global three-dimensional model study of carbonaceous aerosol. *J. Geophys. Res.*, **101**, 19 411–19 432.
- Martins, J. V., P. Artaxa, C. Lioussse, J. S. Reid, P. V. Hobbs, and Y. J. Kaufman, 1998: Effects of black carbon content, particle size, and mixing on light absorption by aerosols from biomass burning in Brazil. *J. Geophys. Res.*, **103**, 32 042–32 050.
- Medalia, A. I., and L. W. Richards, 1972: Tinting strength of black carbon. *J. Colloid Interface Sci.*, **40**, 233–252.
- Moosmuller, H., W. P. Arnott, C. F. Rodgers, J. C. Chow, C. A. Frazier, L. E. Sherman, and D. L. Dietrich, 1998: Photoacoustic and filter

- measurements related to aerosol light absorption during the Northern Front Range Air Quality Study (Colorado 1996/1997). *J. Geophys. Res.*, **103**, 28 149–28 157.
- Novakov, T., D. A. Hegg, and P. V. Hobbs, 1997: Airborne measurements of carbonaceous aerosol on the east coast of the United States. *J. Geophys. Res.*, **102**, 30 023–30 030.
- , T. S. Bates, and P. K. Quinn, 2000: Shipboard measurements of concentrations and properties of carbonaceous aerosols during ACE-2. *Tellus*, **52B**, 228–238.
- Penner, J. E., 1986: Uncertainties in the smoke source term for “nuclear winter” studies. *Nature*, **324**, 222–226.
- Petzold, A., C. Kopp, and R. Niessner, 1997: The dependence of the specific attenuation cross-section on black carbon mass fraction and particle size. *Atmos. Environ.*, **31**, 661–672.
- Putaud, J.-P., and Coauthors, 2000: Chemical mass closure and assessment of the origin of the submicron aerosol in the marine boundary layer and the free troposphere at Tenerife during ACE-2. *Tellus*, **52B**, 141–168.
- Ramanathan, V., and Coauthors, 1996: Indian Ocean Experiment (INDOEX): A multi-agency proposal for a field experiment in the Indian Ocean. Scripps Institution of Oceanography Tech. Rep. 162, 82 pp.
- Redemann, J., P. B. Russell, and P. Hamill, 2001: Dependence of aerosol light absorption and single scattering albedo on ambient relative humidity for sulfate aerosols with black carbon cores. *J. Geophys. Res.*, **106**, 27 485–27 495.
- Reid, J. S., and P. V. Hobbs, 1998: Physical and optical properties of young smoke from individual biomass fires in Brazil. *J. Geophys. Res.*, **103**, 32 013–32 030.
- , —, C. Lioussé, J. V. Martins, R. E. Weiss, and T. F. Eck, 1998: Comparison of techniques for measuring shortwave absorption and black carbon content of aerosols from biomass burning in Brazil. *J. Geophys. Res.*, **103**, 32 031–32 040.
- Russell, P. B., P. V. Hobbs, and L. L. Stowe, 1999a: Aerosol properties and radiative effects in the United States east coast plume: An overview of the Tropospheric Aerosol Radiative Forcing Observational Experiment (TARFOX). *J. Geophys. Res.*, **104**, 2213–2222.
- , and Coauthors, 1999b: Aerosol-induced radiative flux changes off the United States mid-Atlantic coast: Comparison of values calculated from sunphotometer and in situ data with those measured by airborne pyranometer. *J. Geophys. Res.*, **104**, 2289–2307.
- Seinfeld, J. H., and S. N. Pandis, 1998: *Atmospheric Chemistry and Physics*. John Wiley and Sons, 1326 pp.
- Sokolik, I., and O. B. Toon, 1997: Direct radiative forcing by anthropogenic airborne mineral aerosols. *Nature*, **381**, 681–683.
- , and —, 1999: Incorporation of mineralogical composition into models of the radiative properties of mineral aerosol from UV to IR wavelengths. *J. Geophys. Res.*, **104**, 9423–9444.
- Stagg, B. J., and T. T. Charalampopoulos, 1993: Refractive indices of pyrolytic graphite, amorphous carbon and flame soot in the temperature range 25° to 600°C. *Combust. Flame*, **94**, 381–396.
- Turco, R. P., O. B. Toon, T. P. Ackerman, J. B. Pollack, and C. Sagan, 1991: Nuclear winter: Physics and physical mechanisms. *Annu. Rev. Earth Planet. Sci.*, **19**, 383–422.
- Twitty, J. T., and J. A. Weinman, 1971: Radiative properties of carbonaceous aerosol. *J. Appl. Meteor.*, **10**, 725–731.
- van de Hulst, H. C., 1957: *Light Scattering by Small Particles*. John Wiley, 470 pp.