Short-Term Solar Modulation of the Madden–Julian Climate Oscillation

LON L. HOOD

Lunar and Planetary Laboratory, The University of Arizona, Tucson, Arizona

(Manuscript received 6 September 2017, in final form 4 January 2018)

ABSTRACT

Normalized occurrence rates of daily Madden–Julian oscillation (MJO) events are calculated as a function of phase lag relative to peaks and minima in solar ultraviolet flux occurring on the solar rotational time scale (~ 27 days). All MJO phases and four solar maximum periods are considered (1979–83, 1989–93, 1999–2003, and 2011–15). Corresponding daily static stabilities in the tropical lower stratosphere (70–100 hPa) are calculated from ERA-Interim data and are averaged over the warm pool region. The statistical significance of occurrence-rate changes following UV peaks and minima is assessed using a Monte Carlo method. When MJO events with amplitudes greater than about 2 are considered during the December–May period (about 15% of those days), significant reductions of MJO occurrence rates and associated increases in static stability in the tropical lower stratosphere are obtained 1–7 days following solar UV peaks. Consistently, cross-correlation analyses of high-pass-filtered daily MJO amplitudes and solar UV flux during the same seasonal period produce significant negative correlations near and following solar UV peaks. Conversely, mean occurrence rates are increased and lower-stratospheric static stabilities are decreased following solar UV minima. The reductions (increases) in occurrence rate following solar UV peaks (minima) are largest when the stratospheric quasi-biennial oscillation is in its easterly phase. Little or no dependence of the solar modulation on the phase of El Niño–Southern Oscillation is obtained.

1. Introduction

Tropical deep convection is highly variable on sub-seasonal time scales and is further modulated at longer time scales by the annual cycle and El Niño–Southern Oscillation (ENSO; e.g., Wheeler and Kiladis 1999; Kiladis et al. 2001; Son et al. 2017). In addition, a number of previous observational and modeling studies have found evidence for a secondary influence of the stratospheric quasi-biennial zonal wind oscillation (QBO) on the interannual distribution of tropical convection (e.g., Giorgetta et al. 1999; Collimore et al. 2003; Liess and Geller 2012; Nie and Sobel 2015). For example, Liess and Geller (2012) analyzed 21.5 years of direct satellite cloud data as well as several cloud proxy datasets to show that convective cloud cover can be larger by as much as $51\% \pm 7\%$ of the annual mean over parts of the tropical western Pacific during the easterly QBO phase (QBOE) as compared to the westerly QBO phase (QBOW).

The Madden–Julian oscillation (MJO) is an eastward-propagating pattern of alternately intense and weak tropical convection and precipitation that accounts for a majority of tropical convective variability on intraseasonal time scales (Madden and Julian 1971, 1994; Kiladis et al. 2005; Zhang 2005). Initiated in the Indian Ocean, it propagates through the warm pool section of the Indo-Pacific region and has a recurrence period at a given location of 30–60 days. It has derivative effects on intraseasonal climate variability in many areas of the world and is important for extended-range weather forecasts (Zhang 2013). However, it remains poorly simulated in most climate models (Hung et al. 2013; DeMott et al. 2015; Ahn et al. 2017).

During the last few years, it has been found that the amplitude of the MJO is modulated by the QBO such that it is strengthened during northern winter under QBOE conditions relative to QBOW conditions (Yoo and Son 2016; Nishimoto and Yoden 2017; Son et al. 2017). This dependence on QBO phase is consistent with that found previously for the QBO influence on tropical convection in general. Marshall et al. (2017) have further demonstrated using retrospective forecasts that accounting for this observed QBO modulation results in improved prediction of the MJO and its convective consequences in the tropics with a lead time of about 8 days.

Supplemental information related to this paper is available at the Journals Online website: <https://doi.org/10.1175/JAS-D-17-0265.s1>.

Corresponding author: Lon L. Hood, lon@lpl.arizona.edu

A possible mechanism for the QBO influence on tropical convection and the MJO is a decrease in static stability in the tropical upper troposphere–lower stratosphere (UTLS) under QBOE conditions relative to QBOW conditions. The latter is caused at least initially by the QBO-induced mean meridional circulation, which produces equatorial upwelling increases and resulting temperature decreases during periods of easterly vertical wind shear (e.g., [Plumb and Bell 1982](#)). The temperature decreases increase the thermal lapse rate in the UTLS, decreasing the stability against convection. This initial influence on UTLS static stability would be augmented by positive feedbacks from below. As originally shown using model data by [Giorgetta et al. \(1999\)](#), one positive feedback involves higher and cooler (lower and warmer) cloud tops during QBOE (QBOW), which result in reduced (increased) outgoing infrared radiation, further cooling (heating) the tropical lower stratosphere and further reducing (increasing) static stabilities in the UTLS. Consistently, [Son et al. \(2017\)](#) show that near-tropopause cirrus frequency is larger during QBOE than during QBOW across a large part of the equatorial Indo-Pacific region.

Another possible positive feedback to enhance the QBO–MJO connection involves MJO influences on the strength of the residual meridional (Brewer–Dobson) circulation and its associated tropical upwelling rate. The latter is dominantly driven by wave drag in the extratropical stratosphere ([Holton et al. 1995](#)), although stratospheric and tropospheric diabatic heating may also contribute ([Plumb and Eluszkiewicz 1999](#)). Equatorial Pacific rainfall and latent heat anomalies, whether associated with ENSO or the MJO, can drive a Rossby wave train that propagates to higher latitudes ([Horel and Wallace 1981](#)). In addition to providing a teleconnection pathway along the Pacific–North America pattern ([Wallace and Gutzler 1981](#)), upward propagation of the waves into the stratosphere during boreal winter–spring has the effect of increasing the residual circulation and increasing the tropical upwelling rate. For example, [Calvo et al. \(2010\)](#) show that the residual circulation strengthens in the lowermost tropical stratosphere during warm ENSO events. If MJO amplitudes are larger during the easterly QBO phase, Rossby wave amplitudes may be increased, leading to increased tropical upwelling, which would further cool the tropical lowermost stratosphere. The reverse would occur under QBOW conditions. The latter feedback, if ultimately verified observationally and with models, may potentially explain why the QBO influence on the MJO is observed primarily during boreal winter.

As discussed further in [sections 2 and 3](#) below, it has most recently been proposed that interannual variability of the MJO in boreal winter is also influenced by tropical upwelling changes associated with the 11-yr

solar cycle ([Hood 2017](#)). In particular, based on past observations, larger-than-average MJO amplitudes are expected during the coming solar minimum if QBOE conditions (as monitored at 50 hPa) prevail during a given winter. However, only about 3.5 solar cycles of data are available since the beginning of continuous satellite measurements in 1979. Solar signals on this time scale can be difficult to separate from those caused by major volcanic eruptions and ENSO.

Here, results of an investigation of possible solar modulation of MJO amplitudes on the time scale of the solar rotation period (about 27 days) are reported. Statistical studies of solar rotational forcing have an advantage in that hundreds of 27-day “cycles” are available since 1979 and the time scale is much shorter than that of major volcanic eruptions and ENSO (e.g., [Hood 1986, 2004](#)). Partial motivation for this work is provided by a recent study of ERA-Interim temperature data that found evidence for a lagged cooling response of tropical tropospheric temperature to 27-day solar ultraviolet variations ([Hood 2016](#)). Some evidence that this response may be related to a reduction in the number of MJO events that propagate eastward into the central Pacific following peaks in short-term solar forcing was also obtained. The cooling response was therefore provisionally interpreted as due to reduced convection and associated latent heat release in the tropical troposphere.

The paper is organized as follows. In [section 2](#), possible mechanisms for solar influences on tropical deep convection and the MJO are discussed. A “top down” mechanism involving solar ultraviolet influences on the upper stratosphere with indirect effects on tropical upwelling rates and static stability in the lowermost stratosphere is selected for specific investigation. In [section 3](#), the adopted proxy solar ultraviolet flux data [spectral irradiance at 205 nm (hereafter F205)] used to define the solar forcing function and methods applied to edit and filter the F205 data during solar maximum periods are summarized. The adopted daily MJO amplitude data [outgoing longwave radiation (OLR)-based MJO index (OMI)], the 50-hPa QBO zonal wind index, and the Niño-3.4 ENSO index are also described. Finally, the methods used to estimate daily static stabilities in the tropical lowermost stratosphere using reanalysis temperature data are summarized. In [section 4](#), a cross-correlation analysis of high-pass-filtered daily OMI MJO amplitude data and F205 is conducted to show that, during the December–May season and for time periods when the MJO amplitude is relatively high, significant negative correlations occur near and following solar UV peaks. Comparisons are made with QBO and 11-yr solar effects on MJO occurrence rates during northern winter, which also indicate that the

stratospheric influence on the MJO is largest for higher MJO amplitudes. In [section 5](#), a linear regression analysis is first performed of lower-stratospheric static stability versus F205 to show that the annual-mean response to 27-day solar forcing is largest in the warm pool region. Then normalized occurrence rates of MJO events with amplitudes greater than a chosen threshold are calculated as a function of phase lag relative to peaks in short-term solar UV flux. The possibility that warm pool static stabilities in the lower stratosphere are increased in association with MJO occurrence-rate decreases following solar UV peaks is specifically investigated. The dependences of the apparent 27-day solar signals on season and ambient conditions in the stratosphere and troposphere (QBO and ENSO phase) are determined. Conclusions and discussion are given in [section 6](#).

2. Possible mechanisms for solar influences on tropical convection

While a variety of solar-related forcings could in principle influence tropospheric climate ([Gray et al. 2010](#)), two main classes of mechanisms have been discussed in recent literature for the 11-yr time scale. These are (i) top-down mechanisms, involving mainly solar UV-related heating of the tropical stratosphere with indirect effects on tropospheric circulation (e.g., [Haigh et al. 2005](#); [Simpson et al. 2009](#)) and (ii) “bottom up” mechanisms, involving direct total solar irradiance (TSI) heating increases at the surface followed by increased evaporation, convection, and latent heat release ([Meehl et al. 2008](#); [Zhou and Tung 2013](#)). Top-down mechanisms have the advantage that solar-cycle UV variations are relatively large (up to 6% near 200 nm compared to ~0.1% for TSI), while bottom-up mechanisms benefit from the fact that the solar energy change is deposited directly into the troposphere. It has been argued using general circulation model simulations with fixed sea surface temperatures and an imposed lower-stratospheric heating anomaly that the top-down mechanism alone is sufficient to produce the 11-yr tropospheric circulation changes that have been inferred from the limited observational record ([Haigh et al. 2005](#); see also [Shindell et al. 1999](#); [Matthes et al. 2004](#); [Balachandran et al. 1999](#)). But some observational analyses have found that the 11-yr warming response in the tropical lower stratosphere extends into the upper troposphere, suggesting that it could be caused in part by increased TSI heating at the surface, followed by convection and latent heat release at higher altitudes ([Zhou and Tung 2013](#)). Most recently, [Misios et al. \(2016\)](#) have found evidence in climate model data for a surface warming in the equatorial Pacific lagging

solar-cycle maxima that could be a consequence of the small 11-yr TSI variation.

However, current evidence reviewed in the introduction for a QBO–MJO connection and a lagged cooling response to 27-day solar forcing in the tropical troposphere provide new support for a top-down solar mechanism for influencing tropical tropospheric convection and the MJO. First, as suggested by [Yoo and Son \(2016\)](#) and [Nishimoto and Yoden \(2017\)](#), the observed influence of the QBO on MJO amplitudes most probably occurs via changes in lower-stratospheric temperature, which are driven at least initially by tropical upwelling changes originating above the tropopause. This supports the general feasibility of top-down forcing from the stratosphere. Second, as discussed further below, the top-down solar forcing mechanism also involves a modulation of tropical upwelling rates, which would potentially change lower-stratospheric temperatures and static stabilities in a manner similar to that observed for the QBO. Third, the most consistent correlations of tropical tropospheric temperature with 27-day solar forcing were obtained when UV irradiance was used as the solar forcing variable rather than alternatives such as TSI and galactic cosmic ray (GCR) flux [see Fig. S5 of [Hood \(2016\)](#)]. Nevertheless, as discussed in the introduction, positive feedbacks from below undoubtedly enhance the QBO–MJO connection and are likely to also play an important role in enhancing the solar-induced tropical tropospheric response. In this respect, top-down and bottom-up solar forcing are not entirely independent of one another even if direct TSI forcing of surface heating can be neglected.

As reviewed recently by [Mitchell et al. \(2015\)](#); see their Fig. 1), solar UV forcing on the 11-yr time scale can modify tropical upwelling rates (see also [Kodera and Kuroda 2002](#); [Matthes et al. 2004](#)). As outlined in Fig. 1, this can potentially result in influences on tropical deep convection and the MJO. Under solar maximum (SMAX) conditions, increased ozone production and UV heating in the tropical upper stratosphere positively perturb the midlatitude zonal wind with maximum amplitude near the time of solstice in the winter hemisphere (e.g., [Hood et al. 1993](#)). As described originally by [Kodera and Kuroda \(2002\)](#), this leads to a reduction of planetary-scale Rossby wave flux convergence and absorption ($\nabla \cdot \mathbf{F} > 0$), an associated additional positive perturbation of the midlatitude zonal-mean zonal wind ($\partial \bar{u} / \partial t > 0$) and a polar night jet that propagates poleward and downward. Observational evidence for a reduction of stratospheric planetary wave amplitudes at middle to high latitudes at solar maximum relative to solar minimum has been reported ([Powell and Xu 2011](#)). The strengthened zonal wind is accompanied by a

FIG. 1. A possible mechanism by which an increase in solar UV flux can result in a significant reduction of convective activity and temperature in the tropical troposphere. Heavy dark arrows indicate the flow of the top-down mechanism; thin arrows represent the solar-induced anomaly of the Brewer–Dobson circulation. See the text for details.

slowing of the residual meridional (Brewer–Dobson) circulation (thin arrows) via the Coriolis torque, which results in relative downwelling in the tropics ($\Delta \bar{w}^* < 0$). The relative downwelling at solar maximum is analogous to the tropical upwelling decrease under QBO conditions but extends to higher tropical latitudes. It leads to a combination of adiabatic heating and diabatic heating, the latter being due to an associated transport-induced ozone increase in the tropical lower stratosphere (Hood and Soukharev 2003). The net relative heating reduces the temperature gradient in the UTLS, lowering the tropopause altitude and increasing static stability, potentially leading to reduced convection in the upper troposphere. Associated reduced latent heat release would result in a cooling in the tropical troposphere (Hood 2016). The reverse would occur under solar minimum (SMIN) conditions ($\Delta \bar{w}^* > 0$), favoring increased convection as occurs under QBOE conditions. Consistent with this mechanism, some evidence that mean MJO amplitudes during boreal winter are especially large under QBOE/SMIN conditions and especially small under QBOW/SMAX conditions has been reported (Hood 2017; see Fig. 6 below). However, positive feedbacks analogous to those that apparently enhance the QBO/MJO connection are clearly needed to amplify the solar-induced MJO response to a detectable level.

The mechanism on the 11-yr time scale outlined in Fig. 1 has so far been only partly verified in observations and model simulations. The importance of including the solar-induced stratospheric ozone response when modeling solar effects on climate was first emphasized by Haigh (1994, 2003). The weakening of the Brewer–Dobson circulation and propagation of the zonal wind anomaly near solar

maxima was first simulated in a climate model by Matthes et al. (2004). The existence of a secondary, dynamically induced ozone and temperature response in the tropical lower stratosphere associated with changes in upwelling rates has been inferred from analyses of satellite ozone and reanalysis temperature data (Gray et al. 2010; Hood and Soukharev 2012). However, this has been difficult to confirm in observations because of the short record of accurate global satellite data and in climate models because of possible aliasing of the modeled solar signal by the effects of volcanic eruptions and sea surface temperature variations related to ENSO (Marsh and Garcia 2007; Chiodo et al. 2014; Hood et al. 2015; Kuchar et al. 2017). Furthermore, current models tend to underestimate the magnitude of the solar-induced temperature response near the stratopause and the polar night jet response relative to that which is calculated from meteorological reanalysis data (Mitchell et al. 2015). This may be due in part to a less-than-realistic representation of the solar-cycle ozone signal in models that do not include interactive ozone chemistry (Maycock et al. 2016, 2018). On the positive side, some models within the minority that do include chemistry produce a stronger solar-induced variation in the upper stratosphere that is more consistent with available observations (Hood et al. 2015). However, even these latter models are not able to verify a tropical lower-stratospheric response during the last few decades because of possible volcanic aliasing.

In the remainder of the paper, it will be assumed as a working hypothesis that the basic mechanism by which solar forcing impacts tropical convection and the MJO on the 27-day time scale is similar to that proposed for the 11-yr time scale in Fig. 1. Solar UV flux is therefore

adopted as the primary solar forcing variable in the next section. However, correlative analyses will also be conducted using TSI for comparison purposes in section 4. The MJO response to 27-day solar UV forcing is expected to occur as a weakening near UV peaks and as a strengthening near UV minima but with a finite time lag to account for the radiative lifetime in the upper stratosphere and the development of a secondary dynamical response.

A possible criticism of this hypothesized mechanism is that the chain of events outlined in Fig. 1 for the 11-yr time scale may not be fast enough to explain a solar rotational response with a lag of a week or less. However, several considerations suggest that such a response is feasible. First, wave-breaking events in the extratropical winter stratosphere (in the most extreme case, stratospheric polar warming events) produce observable coolings in the tropical upper and lower stratosphere almost immediately [within a few days; see, e.g., Figs. 4 and 7 of Chandra (1985) and Figs. 9 and 11 of Hood and Soukharev (2003)]. Therefore, if solar UV-induced perturbations of the upper stratosphere are able to modify Rossby wave breaking behavior shortly after a 27-day UV peak, then a rapid response of static stability in the tropical lower stratosphere is possible. Second, short-term solar UV forcing consists of a series of UV peaks and minima at ~ 27 -day intervals rather than a single isolated maximum. While the tropospheric convective response to a single isolated UV increase may be undetectable, a larger response to periodic forcing may occur if the convective system behaves in a manner analogous to that of a sinusoidally forced damped harmonic oscillator with a driving frequency near the resonance frequency. This is possible if the MJO, which has a preferred periodicity in the 30–60-day range, is involved in producing the tropospheric response. Observational evidence for the existence of a detectable dynamical response in the stratosphere on the 27-day time scale has been previously reported using several independent datasets (Hood and Jirikowic 1991; Hood 2003, 2016).

3. Data and methods

Following previous work, the F205, estimated according to the Naval Research Laboratory model, version 2 (NRL2; e.g., Lean 2000), is adopted here as an indicator of solar UV variability. The solar flux at this wavelength penetrates well into the upper stratosphere where UV-induced ozone production and radiative heating are important but is located short of the AI I ionization edge where UV variability remains relatively large (e.g., Fig. 3 of Heath and Schlesinger 1986). The NRL2 205-nm model solar fluxes are available from the Laboratory for Atmospheric and Space Physics (LASP)

FIG. 2. Time series of (a) daily OMI amplitude, (b) 50-hPa equatorial zonal wind (QBO index), (c) monthly N3.4 ENSO anomaly, and (d) daily solar spectral irradiance at 205 nm according to the NRL2 showing the four 5-yr time periods selected for detailed analysis. Solar cycles are also indicated by number.

at the University of Colorado, Boulder (http://lasp.colorado.edu/lisird/data/nrl2_files). Figure 2d plots the daily time series with brackets indicating four 5-yr intervals centered on solar maxima that are selected for detailed study in the remainder of the paper.

As reviewed by Yoo and Son (2016) and Hood (2017), the two main MJO indices are the real-time multivariate MJO (RMM) index (Wheeler and Hendon 2004) and the OMI (Kiladis et al. 2014). The OMI is based entirely on OLR data and requires prefiltering to eliminate variability on time scales less than 20 days and greater than 96 days. Although it is consequently not suitable for real-time weather prediction, it has the advantage that it more directly represents the convective component of MJO variability (via OLR). Therefore, as in previous work, the OMI is adopted here. Daily OMI amplitude and phase data are available from the National Oceanic and Atmospheric Administration's Earth System Research Laboratory (<http://www.esrl.noaa.gov/psd/mjo/mjoindex>). The OMI consists of a projection of 20–96-day-filtered satellite OLR data onto the daily spatial empirical orthogonal function (EOF) patterns of 30–96-day eastward-filtered

OLR (Kiladis et al. 2014). The daily OMI amplitude is defined as the square root of the sum of squares of the daily eigenvalues of the first two EOFs. The daily OMI amplitude over the period from 1 January 1979 to 26 April 2016 is plotted in Fig. 2a.

Also following previous work, the equatorial zonal wind at 50 hPa (u_{50}) is adopted as an index of QBO variability. When the QBO at this level is westerly (easterly), relative downwelling (upwelling) typically occurs in the lowermost stratosphere (70–100 hPa). It therefore provides a useful indicator of QBO-induced static stability changes in the UTLS that could affect upper-tropospheric convection. Direct monthly mean near-equatorial radiosonde measurements at a series of lower-stratospheric pressure levels have been compiled at the Freie Universität Berlin (FUB) and are available online (<http://www.geo.fu-berlin.de/en/met/ag/strat/produkte/qbo/index.html>). The monthly FUB data at 50 hPa are specifically used here and are plotted in Fig. 2b. Daily values in a given month are assumed to be equal to monthly means. Westerly (easterly) QBO is defined for simplicity as $u_{50} > 0$ (< 0).

Finally, the Niño-3.4 index (N3.4) is adopted as an indicator of the phase of ENSO. N3.4 is defined as the mean sea surface temperature within 5°S – 5°N , 120° – 170°W and is available online (<http://www.cpc.ncep.noaa.gov/data/indices>). Monthly mean N3.4 anomalies (deviations from long-term monthly means) are plotted in Fig. 2c. Anomaly values > 0 (< 0) are assumed to imply El Niño (La Niña) conditions. Daily values in a given month are assumed to be equal to monthly mean anomalies.

Figure 3 shows the result of filtering in the time domain the solar F205 data for the four 5-yr solar maximum intervals considered here. A similar time-domain filtering technique is applied to lower-stratospheric static stability data in section 5. Specifically, 5-day running means and 35-day running means were first calculated. Then deviations of the 5-day running means from 35-day running means were calculated. Periodicities shorter than ~ 5 days and longer than ~ 35 days are thereby suppressed. The dependence of the results on the choice of a 35-day upper bound is investigated below by carrying out separate analyses using deviations from 47- and 61-day running means.

Brackets in Fig. 3 indicate edited time periods when solar rotational UV variations were especially strong. These edited intervals are the same as considered by Hood (2016) but with the addition of data for the 2011–15 period. Limitation of the analysis to edited intervals should ideally increase the detectability of any atmospheric responses. A total of 136 peaks with deviation amplitudes $> 0.1 \text{ mW m}^{-2} \text{ nm}^{-1}$ and 128 minima with amplitudes $< -0.1 \text{ mW m}^{-2} \text{ nm}^{-1}$ were identified in the edited intervals for analysis in the next section. Tables S1 and S2 of the supporting online material (SOM) list

FIG. 3. Deviations from the 35-day running means of 5-day running-mean solar spectral irradiance at 205 nm according to the NRL2 (same data as in Fig. 2d). Brackets indicate periods of relatively strong UV variations, which are selected for edited analysis.

the dates and amplitudes of each selected peak and minimum.

To calculate daily static stabilities at individual grid points, four-times-daily ERA-Interim temperature data on a $1^{\circ} \times 1^{\circ}$ latitude–longitude grid at the 70- and 100-hPa levels were first downloaded from the European Centre for Medium-Range Weather Forecasts (<http://apps.ecmwf.int/datasets>). At each grid point and at both levels, daily mean atmospheric mass densities were then calculated from the ideal gas law, and the vertical separation between levels was estimated from the hydrostatic law. Static stability was calculated as the difference between the dry adiabatic lapse rate ($\sim 9.8 \text{ K km}^{-1}$) and the observed lapse rate. At the cold temperatures of the upper troposphere and lower stratosphere, the dry adiabatic lapse rate should be approximately equal to the moist adiabatic rate. As explained further in section 5, to simplify the comparison of MJO occurrence rates with static stability in the lower stratosphere, daily mean

FIG. 4. As in Fig. 3, but for OMI MJO amplitude deviations (same data as in Fig. 2a).

static stabilities are computed by averaging over the warm pool region, defined following Yoo and Son (2016) as extending from 10°S to 10°N and from 60°E to 180° . Vertical temperature gradients (dT/dz) are normally positive in the lowermost stratosphere, so the observed lapse rate ($-dT/dz$) is negative. Hence, mean static stabilities over the warm pool region are typically more than 12 K km^{-1} even during the winter season. While this indicates high stability against convection on the average, a lower-than-average value on a given day implies an increased probability that some areas within the region are convectively unstable. The reverse is true if the static stability is higher than average.

4. Cross-correlation analyses

As a first approach toward investigating whether a lagged solar signal is detectable in the OMI MJO amplitude data of Fig. 2a, the daily amplitude data were high-pass filtered in the time domain in the same manner as was done for the F205 data of Fig. 3, and then

FIG. 5. Cross-correlation functions of OMI MJO amplitude deviations from the 35-day running mean (Fig. 4) vs similar deviations of the solar flux at 205 nm according to the NRL2 (Fig. 3). Only days during the DJFMAM season with OMI amplitude greater than a given threshold are retained. Solid lines indicate use of data after editing as shown by brackets in Fig. 4 (261 days for OMI > 2.0 , 449 days for OMI > 1.75 , 661 days for OMI > 1.5 , 1253 days for OMI > 1.0 , and 2069 days for OMI > 0). Dashed lines indicate use of all available data (518 days for OMI > 2.0 , 821 days for OMI > 1.75 , 1171 days for OMI > 1.5 , 2155 days for OMI > 1.0 , and 3622 days for OMI > 0).

cross-correlation functions were calculated. As shown in Fig. 4, deviations from the 35-day running means of 5-day running-mean OMI MJO amplitude were calculated, and the same edited intervals with relatively strong UV variations identified in Fig. 3 were selected.

Figure 5 shows cross-correlation results for the December–May (DJFMAM) period using the edited and filtered data of Figs. 3 and 4 (solid lines) as well as the unedited data (dashed lines). A total of 3622 days of solar maximum data (2069 days after editing) are available for this seasonal period. Separate cross-correlation functions were calculated for days when the unfiltered OMI amplitude exceeded a given threshold at zero lag. As shown in the figure, no significant correlation at any phase lag is obtained when all OMI amplitudes are considered (Fig. 5e; 2069 days). However, as the minimum OMI threshold is increased, stronger correlations are found with maxima

and minima separated by about 27 days. An increasingly negative correlation coefficient R is obtained near and following UV peaks. For example, for $\text{OMI} > 2.0$, $R = -0.40$ at +1-day lag. Because of the 5-day smoothing, the number of observations is $261/5 \simeq 52$. Successive 5-day averages of either F205 or OMI deviations have serial correlation coefficients of less than 0.35, which leads to an effective sample size of $52 \times 0.78 = 41$ [Bretherton et al. 1999, their Eq. (31)]. The correlation is therefore significant at more than the 95% confidence level (e.g., Bevington 1969). Similarly, the lagged negative correlation for $\text{OMI} > 1.75$ is $R = -0.25$ at +2 days, and the number of observations is $449/5 \simeq 89$. The effective sample size is $89 \times 0.78 = 69$, and the correlation is again significant at the 95% confidence level. But for lower OMI amplitude thresholds, the correlations become insignificant in spite of the larger sample sizes.

Figures S1–S4 of the SOM show results similar to that of Fig. 5 but for each of the four 3-month seasons (DJF, MAM, JJA, and SON). For $\text{OMI} > 2.0$, $R = -0.28$ at 3-days lag using edited data for DJF (Fig. S1), while $R = -0.49$ at 0-days lag for MAM (Fig. S2). These correlations are not formally significant because of the smaller effective sample sizes. However, the approximate agreement of the lag of maximum negative correlation for the two time periods supports the existence of a similar signal in both seasons. For this reason, the DJF and MAM periods were combined to yield a statistically significant correlation as shown in Fig. 5.

On the other hand, the JJA and SON results in Figs. S3 and S4 are based on a reduced number of days with OMI amplitudes > 2.0 (74 days for JJA and 78 days for SON within the edited period; 100 days in JJA and 92 days for SON using the unedited data). This reflects the generally weaker activity of the MJO during this half of the year. As seen in the figures, cross-correlation results for JJA and SON at higher MJO amplitudes are inconsistent. In most of the remainder of the paper, attention will therefore be focused on the December–May seasonal period.

The correlative results of Fig. 5 are not very sensitive to the choice of a 35-day upper bound in the time-domain high-pass-filtering method used here. To demonstrate this, results of alternate analyses using deviations from 47- and 61-day running means are shown in Figs. S5 and S6. For example, the correlation using edited data for $\text{OMI} > 2.0$ at 1-day lag changes from -0.40 in Fig. 5 to -0.39 in Fig. S5 to -0.36 in Fig. S6. A more exact elimination of periods beyond a certain limit can be obtained using a digital bandpass filter in the frequency domain rather than using the time-domain method employed here. However, given the insensitivity shown in Figs. S5 and S6, it is unlikely that using a bandpass filter would yield very different results. Cross-spectral methods could also be applied

rather than the time-domain cross-correlation approach used here. However, these methods usually require the time series to be at least weakly stationary (having a consistent spectrum throughout). This may not be the case for a discontinuous time series composed of only data above a given OMI threshold.

Given the possibility reviewed in section 2 that bottom-up forcing involving direct penetration of total irradiance into the troposphere could contribute to the present results, the analysis of Fig. 5 has been repeated using daily TSI as estimated according to the NRL2. The NRL2 TSI data were also obtained from LASP (section 3) (http://lasp.colorado.edu/lisird/data/nrl2_files). Figure S7 shows the time series for the four solar maximum periods after calculating deviations of 5-day running means from 35-day running means as was done for the F205 data of Fig. 3. In addition to 27-day TSI increases, larger decreases are evident, which are presumably associated with the appearance of major sunspot groups on the solar disk. This is one difference of the TSI variability compared to UV variability, which is due almost entirely to emission from bright plage regions in the chromosphere near sunspots. Nevertheless, there is a positive correlation between the smoothed and filtered F205 time series of Fig. 3 and the TSI time series of Fig. S7 ($R \simeq 0.24$).

Figure S8 shows cross-correlation functions for filtered OMI amplitude versus filtered TSI considering the same edited intervals in Fig. 3 that were used for the analysis of Fig. 5. The correlations for the edited intervals (solid lines) are qualitatively similar to those of Fig. 5 but are generally weaker. For example, for $\text{OMI} > 2.0$, $R = -0.26$ at +3 days in Fig. S8, while $R = -0.4$ at +1 day in Fig. 5. Also, using the edited intervals rather than the unedited data gives a noticeably stronger correlation with F205 as the solar variable (Fig. 5) but produces almost no improvement when TSI is adopted (Fig. S8). Figure S9 shows similar results when the edited intervals of Fig. S7 are considered rather than those of Fig. 3. The correlation is slightly weaker ($R = -0.24$ at +3 days), and there is again no improvement for the edited results over the unedited results. Given that there is a modest positive correlation between TSI and F205, it is possible that the correlations shown in Figs. S8 and S9 are caused mainly by the UV forcing rather than the TSI forcing. In any case, there is no empirical evidence that the assumption of TSI forcing yields an improved correlation. Also, as reviewed in section 2, TSI-based mechanisms predict an increase in tropical convection with increasing TSI, in contrast to the apparent negative correlation following TSI peaks found in Figs. S8 and S9.

Finally, the correlative results of Fig. 5 indicate that stronger reductions of MJO amplitude near and following short-term solar UV peaks are found when the unfiltered

FIG. 6. Summary of QBO and 11-yr solar normalized occurrence-rate results for northern winter. (a) The two bars on the left show the percentages of days during DJF and during each QBO phase in 1979.0–2016.3 when the OMI amplitude exceeded unity [from Fig. 2a of Hood (2017)]. Error bars are the 95% confidence limits based on Monte Carlo simulations. The remaining bars show how these

MJO amplitude is relatively high. There is similar evidence that the stratospheric influence on the MJO during northern winter at longer time scales is strongest when the MJO amplitude is relatively high. Figure 6a shows normalized occurrence rates of MJO events during DJF and during the two phases of the QBO separated into different ranges of MJO OMI amplitude. As mentioned in section 2, the westerly phase (QBOW) is defined as existing for days when the monthly mean u_{50} (Fig. 2b) is >0 , while QBOE is defined for $u_{50} < 0$. The two bars on the left [taken from Fig. 2a of Hood (2017)] show that a larger fraction of days have an active MJO (defined as $OMI > 1$) during QBOE ($\sim 75\%$) than during QBOW ($\sim 56\%$). The remaining bars show that most of the difference occurs for days when $OMI > 2.0$ ($35\% \pm 7\%$ during QBOE vs $15\% \pm 10\%$ during QBOW).

Figures 6b and 6c show similar plots for the case when the two phases of the solar cycle are considered in addition to the QBO. SMAX conditions are assumed to prevail when the daily value of F_{205} exceeds $10.5 \text{ mW m}^{-2} \text{ nm}^{-1}$, while SMIN conditions are defined for F_{205} less than $10.1 \text{ mW m}^{-2} \text{ nm}^{-1}$ (Fig. 2d). In Fig. 6b, the two forcings are expected to be working together (i.e., during QBOE/SMIN conditions, both would produce relative upwelling in the tropical lower stratosphere, while during QBOW/SMAX conditions, both would be expected to produce relative downwelling). Under these conditions, based on 37 years of data, as indicated by the two bars on the left, about 90% of days during QBOE/SMIN conditions had an active MJO, while only about 50% of days during QBOW/SMAX conditions had $OMI > 1$. The remaining bars in Fig. 6b show that most of this large difference occurs for days when $OMI > 2.0$ ($57\% \pm 10\%$ of days during QBOE/SMIN conditions vs $10\% \pm 11\%$ during QBOW/SMAX conditions). Figure 6c shows a plot for the case when the two forcings are assumed to be opposed. Under these conditions, the occurrence rate under QBOE/SMAX conditions is only marginally larger than that during QBOW/SMIN conditions, and there is no statistically significant difference for $OMI > 2$.

5. MJO occurrence rates relative to solar UV extrema

According to the 11-yr mechanism outlined in Fig. 1 and considered here also for the 27-day time scale

←

percentages depend on MJO amplitude. (b) As in (a), but during the QBOE/SMIN and QBOW/SMAX conditions. (c) As in (a) and (b), but during the QBOE/SMAX and QBOW/SMIN conditions. See the text.

FIG. 7. Static stability response estimated by linear regression in the lowermost stratosphere (70–100 hPa) to an increase in F205 of $0.3 \text{ mW m}^{-2} \text{ nm}^{-1}$ as a function of phase lag. All edited data (about 4500 days) were considered. Heavy dark lines enclose regions that are formally significant at the 2σ (95% confidence) level.

(at least for a resonant response as discussed near the end of section 2), reductions in MJO occurrence rates are to be expected following solar UV peaks in conjunction with increases in static stability in the lowermost stratosphere. The opposite is expected following 27-day solar UV minima. While the mechanism is hypothesized to be valid for tropical convection in general, it is possible that the MJO, which has more vertically extended, large-scale convection than more localized convective systems (Fig. 16 of Madden and Julian 1972), is more sensitive to forcing from the upper boundary of the troposphere. Downward penetration of the convective response into the middle and lower troposphere may be facilitated by positive feedbacks such as the cloud radiative and Rossby wave amplification mechanisms described in the introduction. To investigate this hypothesis, normalized occurrence rates of MJO events with amplitudes greater than a given threshold and corresponding static stabilities in the lowermost stratosphere were calculated and are presented in this section. All calculations were done as a function of phase lag relative to short-term solar UV peaks and minima.

In previous work (e.g., Yoo and Son 2016; Hood 2017), static stabilities in the tropical lowermost stratosphere have been estimated using methods described in section 2 with results averaged over the warm pool region (10°S – 10°N , 60°E – 180°). To investigate whether averaging over the warm pool region is a reasonable approach for the present application, the static stability response to 27-day solar UV forcing has been estimated by carrying out a linear regression analysis as follows: The F205 and static stability time series at individual $3^{\circ} \times 3^{\circ}$ grid points were first filtered in the time domain as described in section 3 so that periodicities outside of the 5–35-day window were suppressed. To allow a determination of the response at a given phase lag relative to the UV peak, the static stability deviation time series were displaced forward or backward in time. Only solar and linear trend independent variables (basis functions) were considered so that static stability deviations at a given grid point, $ss'(t)$, were assumed to be related to daily F205 deviations by $ss'(t) = \beta_{\text{solar}}F205'(t) + \beta_{\text{trend}}t + r(t)$, where t is time (days), β_{solar} and β_{trend} are regression coefficients, and $r(t)$ is the residual noise term. The value of β_{trend} is normally very small but was calculated for completeness.

Autocorrelation of the regression residuals (data minus statistical model) was accounted for using a Cochrane–Orcutt (prewhitening) transformation (Cochrane and Orcutt 1949; Tiao et al. 1990; Garny et al. 2007). Specifically, both the static stability deviation time series and the filtered solar and linear trend basis functions were adjusted to ensure that the residuals

FIG. 8. (a) Normalized occurrence rates (%) of daily MJO amplitudes ≥ 2 during the DJFMAM season at a series of lags relative to dates when the 5–35-day-filtered solar flux at 205 nm reached a pronounced peak (percentage of 66–70 peak days during 1979–83, 1989–93, 1999–2003, and 2011–15). Error bars are 95% confidence limits based on Monte Carlo simulations. All MJO phases and both QBO phases are considered. (b) Corresponding mean static stabilities and one standard deviation limits in the tropical lowermost stratosphere (70–100 hPa) averaged over the warm pool region (see the text). The horizontal dotted lines are the long-term-mean occurrence rate and long-term-mean static stability for MJO amplitudes ≥ 2.0 during DJFMAM under solar maximum conditions. (c),(d) As in (a) and (b), respectively, but relative to dates when the filtered solar flux reached a pronounced minimum (62–65 minimum days).

are approximately white noise. This was done by first determining the autoregression coefficient ρ of the residuals, assuming that they can be represented as a first-order autoregressive process. The observed time series at each grid point and the two basis functions were then transformed as follows: $x(t) \rightarrow x(t) - \rho x(t-1)$, where $x(t)$ is a value at time t and $x(t-1)$ is the value at the previous observation time. If the residuals were significantly autocorrelated, the net effect is a reduction of the regression coefficient amplitudes and an increase of the standard error estimates.

Solar regression coefficient results for the annual-mean response using all edited data (about 4500 days) are shown in Fig. 7. It is seen that a maximum response occurs within the longitude range of the warm pool region about 4 days after the UV peak. This static stability response is unlikely to be a direct consequence of the UV forcing alone, which would be expected to produce a more zonally symmetric response. Rather, it is more likely that it is a result of amplification by feedbacks from below, which are only initiated by the stratospheric

solar forcing. As shown in Fig. S10 of the SOM, most of this annual-mean response, calculated using all days regardless of MJO amplitude, occurs during the June–November (JJASON) half of the year rather than during DJFMAM when the best correlations of filtered F205 versus filtered OMI amplitude are found. However, as shown in Fig. S11, the static stability response during DJFMAM becomes larger in the western part of the warm pool region when only days with strong MJO amplitudes are considered. Specifically, it becomes stronger in the Indian Ocean sector while becoming weaker in the Pacific to the west of the date line. Based on these results and to be consistent with previous work, the static stability amplitudes calculated in this section will mainly be averaged over the warm pool region as defined above. However, alternate analyses using other averaging sectors will also be performed to allow an assessment of the sensitivity of the results to this choice.

At a given phase lag, the normalized occurrence rate is defined as the number of UV peaks or minima when the OMI amplitude exceeds the chosen threshold

FIG. 9. (a),(b) As in Figs. 8a and 8b, respectively, but only during the easterly phase of the QBO (percentage of 59–60 peak days during 1979–83, 1989–93, 1999–2003, and 2011–15). All seasons and MJO phases are considered. (c),(d) As in (a) and (b), respectively, but relative to dates when the filtered solar flux reached a pronounced minimum (58–59 minimum days).

divided by the total number of strong UV peaks or minima considered in the analysis. As noted in the previous section, a total of 136 strong peaks and 128 strong minima were identified during the four 5-yr periods centered on solar maxima. Based on the correlative results of the previous section, calculations were mainly done for the DJFMAM period when correlations are most consistent and there are enough strong peaks or minima for an effective evaluation of statistical significance (especially when considering separate phases of the QBO and ENSO). As described previously (Hood 2017), a Monte Carlo procedure is applied to estimate 95% confidence limits on the occurrence rates. Specifically, the calculations are repeated a large number of times, each time replacing the original MJO amplitude time series with an artificial series whose start date is randomly reselected. Two standard deviation (2σ) limits from these calculations then represent the approximate 95% confidence bounds, which are assumed to apply also to the observed occurrence rates. In the subsequent figures, error limits are shown at lags where occurrence rates are especially large or small or at lags where error estimates are representative of those at nearby lags. Similar criteria are applied to choose lags where one standard

deviation (1σ ; 68% confidence) limits on mean static stabilities are shown.

Figure 8 shows results for OMI MJO amplitudes ≥ 2 relative to solar UV peaks and minima during DJFMAM. Depending on phase lag, 66–70 peak days and 61–65 minimum days were available during DJFMAM within the edited periods of Fig. 3. Normalized occurrence rates and 95% confidence limits are shown in Figs. 8a and 3c, while mean 70–100-hPa static stabilities averaged over the warm pool region (also calculated only for days with MJO amplitudes ≥ 2) and 1σ error limits are shown in Figs. 8b and 8d. The horizontal dotted lines in Figs. 8a and 8c are the annual-mean occurrence rates ($\sim 15\%$ during DJFMAM) of MJO events with amplitudes ≥ 2 , while those in Figs. 8b and 8d are the DJFMAM-mean static stabilities when OMI > 2.0 ($\sim 12.25 \text{ K km}^{-1}$). Near and following the time of solar UV peaks under solar maximum conditions, there is a reduction of occurrence rate from ~ 11 of 68 or $\sim 16\%$ on average before a UV peak to ~ 5 of 68 or $\sim 7\%$ several days after a UV peak, which is more than 2σ below the long-term mean. As shown in Fig. 8b, corresponding mean static stabilities increase by more than one standard deviation following UV peaks as compared to before UV peaks. As seen in Figs. 8c and 8d,

FIG. 10. (a),(b) As in Figs. 9a and 9b, respectively, but only during DJFMAM when the QBO was in its easterly phase ($u_{50} < 0$; percentage of 25–28 peak days during 1979–83, 1989–93, 1999–2003, and 2011–15). (c),(d) As in (a) and (b), respectively, but only for minimum days during DJFMAM when the QBO was in its easterly phase (percentage of 26–28 minimum days).

the behavior is qualitatively opposite following solar UV minima: Occurrence rates increase from ~ 7 of 64 or $\sim 11\%$ before minima to ~ 12 of 64 or $\sim 19\%$ after minima. Static stabilities are also relatively low after minima.

As shown in Figs. S12 and S13 of the SOM, the apparent short-term solar modulation of MJO amplitudes and static stabilities seen in Fig. 8 becomes progressively less detectable as the MJO amplitude threshold is decreased from 2 to 1.75 to 1.5. Also, during the JJASON seasonal period, the solar modulation is undetectable even for MJO amplitudes ≥ 2 (Fig. S14). During JJASON, mean warm pool static stabilities for MJO amplitudes ≥ 2 are higher ($\sim 13.5 \text{ K km}^{-1}$), and mean MJO occurrence rates are lower ($\sim 6\%$). Mean static stabilities vary with phase lag but are generally high during this seasonal period. There is no discernible effect on MJO occurrence rates, which are already low. Both of these characteristics (dependence on MJO amplitude threshold and dependence on season) are consistent with the correlative results of section 4.

As noted above, linear regression analysis indicates that the lower-stratospheric static stability response to 27-day solar forcing has a distribution that differs somewhat from the usual warm pool region during DJFMAM. Specifically, according to Fig. S11 for $\text{OMI} > 2.0$, the maximum response occurs primarily in an equatorial zone extending from about 50°W to 100°E . The analysis of Fig. 8 was therefore repeated using this modified averaging region with results shown in Fig. S15. As seen there, the mean static stabilities following a solar UV peak using the modified averaging region (Fig. S15b) are modestly improved over that shown in Fig. 8b; that is, the maximum value occurs at 2-days lag, in better agreement with the occurrence-rate minimum in Fig. S15a. However, the stabilities following a UV minimum in Fig. S15d are not greatly improved, and there is still no statistically significant change before and after solar UV peaks and minima. For this reason and in order to allow direct comparisons with previous work (e.g., Yoo and Son 2016; Hood 2017), the more

FIG. 11. (a),(b) As in Figs. 8a,b, 9a,b, and 10a,b, but only for peak days when ENSO was in its warm phase ($N3.4 > 0$; percentage of 81–82 peak days during 1979–83, 1989–93, 1999–2003, and 2011–15). All seasons, MJO phases, and both QBO phases were considered. (c),(d) As in (a) and (b), respectively, but only for peak days when ENSO was in its cold phase ($N3.4 < 0$; percentage of 53–54 peak days).

conventional warm pool averaging region (10°S – 10°N , 60°E – 180°) is mainly used in this paper.

Figure 9 shows results in the same format as Fig. 8 but considering only days during the QBOE ($u50 < 0$) regardless of season. For this case, depending on phase lag, 59–60 peak days and 58–59 minimum days are available for analysis. Similar to the DJFMAM results of Fig. 5, warm pool static stabilities tend to increase following solar UV peak days and tend to decrease following UV minimum days. Specifically, MJO occurrence rates for amplitudes ≥ 2 decrease from ~ 10 of 60 or $\sim 17\%$ before UV peaks to ~ 4 of 60 or $\sim 7\%$ following UV peaks, which is more than 2σ below the long-term mean. Corresponding occurrence rates increase from 6 or 7 of 59 or $\sim 11\%$ before minima to ~ 11 of 59 or $\sim 19\%$ following minima. In contrast, as shown in Figs. S16a and 16b of the SOM, under QBOW conditions, while warm pool static stabilities for MJO amplitudes ≥ 2 increase somewhat after UV peaks, there is no discernible effect on MJO occurrence rates, which are already less than 10%. Likewise, no significant effect is seen following UV minima (Figs. S16c,d).

In view of the positive evidence for short-term solar modulation of the MJO during DJFMAM but not

during JJASON and during QBOE but not during QBOW, it may be expected that the strongest modulation would occur during DJFMAM when the QBO is also in its easterly phase. As shown in Fig. 10, this may be the case. Under these conditions, long-term-mean MJO occurrence rates for $\text{OMI} > 2$ are especially large ($\sim 21\%$), and long-term-mean warm pool static stabilities are unusually low ($\sim 11.7 \text{ K km}^{-1}$). There are 25–28 peak days and 26–28 minimum days available for analysis during the edited periods of Fig. 3. Mean occurrence rates decrease from ~ 8 of 27 or $\sim 30\%$ during the week prior to a UV peak to ~ 2 of 27 or $\sim 7\%$ within a few days after a UV peak. Error bars are large because of the smaller number of qualifying peaks, but the reduction following UV peaks is more than 2σ below the mean according to the Monte Carlo simulations. Mean warm pool static stabilities generally increase during the week after a UV peak relative to the week prior, but the difference is not formally significant. Conversely, as shown in Fig. 10c, occurrence rates increase from about 3 of 27 or $\sim 10\%$ per week before a UV minimum to about 8 of 27 or $\sim 30\%$ per week after a UV minimum. Static stabilities in Fig. 10d show only a slight decline following a UV minimum, however.

Finally, Fig. 11 shows MJO occurrence-rate and warm pool static stability results when only UV peak days during the warm and cold phases of ENSO are considered ($N3.4 > 0$ and $N3.4 < 0$, respectively). Depending on phase lag, 81–82 peak days during the warm phase and 53–54 peak days during the cold phase remain within the edited periods, regardless of season and QBO phase. Occurrence-rate results show no statistically significant difference between the two ENSO phases. Static stabilities tend to be larger following UV peaks, and there is a corresponding small reduction in the occurrence rate during both phases. Overall, there is no evidence for a significant ENSO influence on the short-term solar modulation of the MJO.

6. Conclusions and discussion

Both the lagged correlation results of section 4 and the normalized occurrence-rate results of section 5 give consistent evidence for a statistically significant modulation of relatively strong MJO events by solar ultraviolet variations occurring on the solar rotational time scale. The modulation is such that the number of MJO events with amplitudes greater than about 2 is suppressed following UV peaks and is increased following UV minima. The modulation is detectable mainly during the DJFMAM seasonal period and is stronger during the easterly phase of the QBO. According to Fig. 10, the strongest modulation may occur under both DJFMAM and QBOE conditions. No significant difference of the short-term solar modulation of the MJO is obtained under different ENSO conditions (Fig. 11). These results on the 27-day time scale are qualitatively consistent with the proposed QBO/solar modulation of the boreal-winter MJO on the 11-yr time scale (Hood 2017).

The normalized occurrence-rate results of Figs. 8–11 do not require that the total number of MJO events is modulated by solar forcing, only that the amplitudes of the stronger events are reduced (increased) following UV peaks (minima). For example, although not shown here, under DJFMAM/QBOE conditions, the number of MJO events with amplitudes in the 1.5–2 range increases following UV peaks at about the same lags as when the number of events with amplitudes ≥ 2 is being suppressed. This suggests that the same MJO events whose amplitudes are being suppressed are now being counted in the 1.5–2 category.

The static stability results shown in the same figures provide some qualitative support for the hypothesis that solar-induced changes in static stability in the tropical UTLS are the initiating mechanism for the apparent 27-day solar forcing of the MJO. For example, the mean static stability within the warm pool region, for days with

MJO amplitudes ≥ 2 , increases following UV peaks in approximate conjunction with the observed decrease in MJO occurrence rates. However, these static stability increases are not formally significant, and there is no clear indication that they precede or coincide exactly with decreases in MJO occurrence rate. If solar-induced static stability changes are indeed the initiating mechanism, then the final static stability response must be heavily influenced by positive feedbacks from below. As shown in section 5 (Figs. 7, S10, and S11), while the lagged annual-mean static stability response to 27-day solar UV forcing maximizes in the warm pool region, a significant response in this region during the DJFMAM seasonal period only occurs if days with higher MJO amplitudes are considered. The latter increase is concentrated in the Indian Ocean region where individual MJO events originate. Further work is therefore needed to demonstrate a robust causal link between static stability changes in the lower stratosphere and changes in MJO amplitude. It may alternatively be hypothesized, for example, that stratospheric influences on temperature near the tropopause indirectly modify the thermodynamic efficiency and therefore the potential intensity of deep convective systems (e.g., Emanuel et al. 2013).

The evidence reported here for short-term solar modulation of the MJO is obtained mainly for OMI amplitudes ≥ 2 . As shown in Fig. 6, QBO/solar forcing of the MJO on longer time scales also appears to occur mainly for OMI amplitudes ≥ 2 . If so, then the stratospheric influence on the MJO in general may be limited to strong (high OMI amplitude) MJO events. This could be partly because positive feedback mechanisms are most effective for strong MJO events and/or partly that the initial forcing from above is more effective for strong MJO events. As shown in the original work of Madden and Julian (1972, their Figs. 15 and 16), the tropopause is higher and colder above and to the east of the main MJO convective center, which peaks over the warm pool region. Presumably, the strongest MJO events are associated with the highest and coldest tropopause anomalies. Since top-down forcing (e.g., decreased temperatures and static stabilities under QBOE conditions or following solar UV minima) is expected to be effective mainly in the lowermost stratosphere, stronger MJO events with their higher tropopause altitudes may be more favorable for facilitating the solar and QBO influences. Further observational analyses and model studies are needed to evaluate this hypothesis.

Acknowledgments. Investigation of the existence of short-term solar UV radiation effects on the occurrence rate of MJO events and on static stability in the tropical lower stratosphere is supported under a 1.5-yr pilot

study (Grant 1643160) from the Climate and Large-Scale Dynamics branch of the National Science Foundation. Helpful discussions with Dr. George Kiladis of the National Oceanic and Atmospheric Administration and with Dr. Thomas Galarneau and Malori Redman of the Hydrology and Atmospheric Sciences Department at The University of Arizona are appreciated. Thanks also to two anonymous reviewers for taking the time to read the manuscript and provide useful critical reviews.

REFERENCES

- Ahn, M.-S., D. Kim, K. Sperber, I.-S. Kang, E. Maloney, D. Waliser, and H. Hendon, 2017: MJO simulation in CMIP5 climate models: MJO skill metrics and process-oriented diagnosis. *Climate Dyn.*, **49**, 4023–4045, <https://doi.org/10.1007/s00382-017-3558-4>.
- Balachandran, N. K., D. Rind, P. Lonergan, and D. T. Shindell, 1999: Effects of solar cycle variability on the lower stratosphere and the troposphere. *J. Geophys. Res.*, **104**, 27 321–27 339, <https://doi.org/10.1029/1999JD900924>.
- Bevington, P. R., 1969: *Data Reduction and Error Analysis for the Physical Sciences*. McGraw-Hill, 336 pp.
- Bretherton, C. S., M. Widmann, V. P. Dymnikov, J. M. Wallace, and I. Bladé, 1999: The effective number of spatial degrees of freedom of a time-varying field. *J. Climate*, **12**, 1990–2009, [https://doi.org/10.1175/1520-0442\(1999\)012<1990:TENOSD>2.0.CO;2](https://doi.org/10.1175/1520-0442(1999)012<1990:TENOSD>2.0.CO;2).
- Calvo, N., R. Garcia, W. Randel, and D. Marsh, 2010: Dynamical mechanism for the increase in tropical upwelling in the lowermost tropical stratosphere during warm ENSO events. *J. Atmos. Sci.*, **67**, 2331–2340, <https://doi.org/10.1175/2010JAS3433.1>.
- Chandra, S., 1985: Solar-induced oscillations in the stratosphere: A myth or reality? *J. Geophys. Res.*, **90**, 2331–2339, <https://doi.org/10.1029/JD090iD01p02331>.
- Chiodo, G., D. R. Marsh, R. Garcia-Herrera, N. Calvo, and J. Garcia, 2014: On the detection of the solar signal in the tropical stratosphere. *Atmos. Chem. Phys.*, **14**, 5251–5269, <https://doi.org/10.5194/acp-14-5251-2014>.
- Cochrane, D., and D. H. Orcutt, 1949: Application of least squares regression to relationships containing auto-correlated error terms. *J. Amer. Stat. Assoc.*, **44**, 32–61, <https://doi.org/10.1080/01621459.1949.10483290>.
- Collimore, C. C., D. W. Martin, M. H. Hitchman, A. Huesmann, and D. E. Waliser, 2003: On the relationship between the QBO and tropical deep convection. *J. Climate*, **16**, 2552–2568, [https://doi.org/10.1175/1520-0442\(2003\)016<2552:OTRBTO>2.0.CO;2](https://doi.org/10.1175/1520-0442(2003)016<2552:OTRBTO>2.0.CO;2).
- DeMott, C. A., N. P. Klingaman, and S. J. Woolnough, 2015: Atmosphere-ocean coupled processes in the Madden-Julian oscillation. *Rev. Geophys.*, **53**, 1099–1154, <https://doi.org/10.1002/2014RG000478>.
- Emanuel, K., S. Solomon, D. Folini, S. Davis, and C. Cagnazzo, 2013: Influence of tropical tropopause layer cooling on Atlantic hurricane activity. *J. Climate*, **26**, 2288–2301, <https://doi.org/10.1175/JCLI-D-12-00242.1>.
- Garny, H., G. E. Bodeker, and M. Dameris, 2007: Trends and variability in stratospheric mixing: 1979–2005. *Atmos. Chem. Phys.*, **7**, 5611–5624, <https://doi.org/10.5194/acp-7-5611-2007>.
- Giorgetta, M. A., L. Bengtsson, and K. Arpe, 1999: An investigation of QBO signals in the East Asian and Indian monsoon in GCM experiments. *Climate Dyn.*, **15**, 435–450, <https://doi.org/10.1007/s003820050292>.
- Gray, L. J., and Coauthors, 2010: Solar influences on climate. *Rev. Geophys.*, **48**, RG4001, <https://doi.org/10.1029/2009RG000282>.
- Haigh, J. D., 1994: The role of stratospheric ozone in modulating the solar radiative forcing of climate. *Nature*, **370**, 544–546, <https://doi.org/10.1038/370544a0>.
- , 2003: The effects of solar variability on the Earth's climate. *Philos. Trans. Roy. Soc. London*, **361A**, 95–111, <https://doi.org/10.1098/rsta.2002.1111>.
- , M. Blackburn, and R. Day, 2005: The response of tropospheric circulation to perturbations in lower-stratospheric temperature. *J. Climate*, **18**, 3672–3685, <https://doi.org/10.1175/JCLI3472.1>.
- Heath, D. F., and B. M. Schlesinger, 1986: The Mg 280-nm doublet as a monitor of changes in solar ultraviolet irradiance. *J. Geophys. Res.*, **91**, 8672–8682, <https://doi.org/10.1029/JD091iD08p08672>.
- Holton, J. R., P. H. Haynes, M. E. McIntyre, A. R. Douglass, R. B. Rood, and L. Pfister, 1995: Stratosphere-troposphere exchange. *Rev. Geophys.*, **33**, 403–439, <https://doi.org/10.1029/95RG02097>.
- Hood, L. L., 1986: Coupled stratospheric ozone and temperature responses to short-term changes in solar ultraviolet flux: An analysis of Nimbus 7 SBUV and SAMS data. *J. Geophys. Res.*, **91**, 5264–5276, <https://doi.org/10.1029/JD091iD04p05264>.
- , 2003: Thermal response of the tropical tropopause region to solar ultraviolet variations. *Geophys. Res. Lett.*, **30**, 2215, <https://doi.org/10.1029/2003GL018364>.
- , 2004: Effects of solar UV variability on the stratosphere. *Solar Variability and Its Effect on Climate*, *Geophys. Monogr.*, Vol. 141, Amer. Geophys. Union, 283–304.
- , 2016: Lagged response of tropical tropospheric temperature to solar ultraviolet variations on intraseasonal time scales. *Geophys. Res. Lett.*, **43**, 4066–4075, <https://doi.org/10.1002/2016GL068855>.
- , 2017: QBO/solar modulation of the boreal winter Madden-Julian oscillation: A prediction for the coming solar minimum. *Geophys. Res. Lett.*, **44**, 3849–3857, <https://doi.org/10.1002/2017GL072832>.
- , and J. Jirikowic, 1991: Stratospheric dynamical effects of solar ultraviolet variations: Evidence from zonal mean ozone and temperature data. *J. Geophys. Res.*, **96**, 7565–7577, <https://doi.org/10.1029/91JD00228>.
- , and B. E. Soukharev, 2003: Quasi-decadal variability of the tropical lower stratosphere: The role of extratropical wave forcing. *J. Atmos. Sci.*, **60**, 2389–2403, [https://doi.org/10.1175/1520-0469\(2003\)060<2389:QVOTTL>2.0.CO;2](https://doi.org/10.1175/1520-0469(2003)060<2389:QVOTTL>2.0.CO;2).
- , and —, 2012: The lower-stratospheric response to 11-yr solar forcing: Coupling to the troposphere–ocean response. *J. Atmos. Sci.*, **69**, 1841–1864, <https://doi.org/10.1175/JAS-D-11-086.1>.
- , J. L. Jirikowic, and J. P. McCormack, 1993: The stratospheric quasi-decadal variation: Influence of long-term solar ultraviolet variations. *J. Atmos. Sci.*, **50**, 3941–3958, [https://doi.org/10.1175/1520-0469\(1993\)050<3941:QDVOTS>2.0.CO;2](https://doi.org/10.1175/1520-0469(1993)050<3941:QDVOTS>2.0.CO;2).
- , and Coauthors, 2015: Solar signals in CMIP-5 simulations: The ozone response. *Quart. J. Roy. Meteor. Soc.*, **141**, 2670–2689, <https://doi.org/10.1002/qj.2553>.
- Horel, J. D., and J. M. Wallace, 1981: Planetary-scale atmospheric phenomena associated with the Southern Oscillation. *Mon. Wea. Rev.*, **109**, 813–829, [https://doi.org/10.1175/1520-0493\(1981\)109<0813:PSAPAW>2.0.CO;2](https://doi.org/10.1175/1520-0493(1981)109<0813:PSAPAW>2.0.CO;2).
- Hung, M.-P., J.-L. Lin, W. Wang, D. Kim, T. Shinoda, and S. J. Weaver, 2013: MJO and convectively coupled equatorial waves simulated by CMIP5 climate models. *J. Climate*, **26**, 6185–6214, <https://doi.org/10.1175/JCLI-D-12-00541.1>.

- Kiladis, G. N., K. H. Straub, G. C. Reid, and K. S. Gage, 2001: Aspects of interannual and intraseasonal variability of the tropopause and lower stratosphere. *Quart. J. Roy. Meteor. Soc.*, **127**, 1961–1983, <https://doi.org/10.1002/qj.49712757606>.
- , —, and P. T. Haertel, 2005: Zonal and vertical structure of the Madden–Julian oscillation. *J. Atmos. Sci.*, **62**, 2790–2809, <https://doi.org/10.1175/JAS3520.1>.
- , J. Dias, K. H. Straub, M. C. Wheeler, S. N. Tulich, K. Kikuchi, K. M. Weickmann, and M. J. Ventrice, 2014: A comparison of OLR and circulation-based indices for tracking the MJO. *Mon. Wea. Rev.*, **142**, 1697–1715, <https://doi.org/10.1175/MWR-D-13-00301.1>.
- Kodera, K., and Y. Kuroda, 2002: Dynamical response to the solar cycle. *J. Geophys. Res.*, **107**, 4749, <https://doi.org/10.1029/2002JD002224>.
- Kuchar, A., W. Ball, E. Rozanov, A. Stenke, L. Revell, J. Miksovsky, P. Pisoft, and T. Peter, 2017: On the aliasing of the solar cycle in the lower stratospheric tropical temperature. *J. Geophys. Res. Atmos.*, **122**, 9076–9093, <https://doi.org/10.1002/2017JD026948>.
- Lean, J., 2000: A decadal solar effect in the evolution of the sun's spectral irradiance since the Maunder Minimum. *Geophys. Res. Lett.*, **27**, 2425–2428, <https://doi.org/10.1029/2000GL000043>.
- Liess, S., and M. A. Geller, 2012: On the relationship between QBO and distribution of tropical deep convection. *J. Geophys. Res.*, **117**, D03108, <https://doi.org/10.1029/2011JD016317>.
- Madden, R. A., and P. R. Julian, 1971: Detection of a 40–50 day oscillation in the zonal wind in the tropical Pacific. *J. Atmos. Sci.*, **28**, 702–708, [https://doi.org/10.1175/1520-0469\(1971\)028<0702:DOADOI>2.0.CO;2](https://doi.org/10.1175/1520-0469(1971)028<0702:DOADOI>2.0.CO;2).
- , and —, 1972: Description of global-scale circulation cells in the tropics with a 40–50 day period. *J. Atmos. Sci.*, **29**, 1109–1123, [https://doi.org/10.1175/1520-0469\(1972\)029<1109:DOGSCC>2.0.CO;2](https://doi.org/10.1175/1520-0469(1972)029<1109:DOGSCC>2.0.CO;2).
- , and —, 1994: Observations of the 40–50-day tropical oscillation—A review. *Mon. Wea. Rev.*, **122**, 814–837, [https://doi.org/10.1175/1520-0493\(1994\)122<0814:OOTDIO>2.0.CO;2](https://doi.org/10.1175/1520-0493(1994)122<0814:OOTDIO>2.0.CO;2).
- Marsh, D. R., and R. Garcia, 2007: Attribution of decadal variability in lower-stratospheric tropical ozone. *Geophys. Res. Lett.*, **34**, L21807, <https://doi.org/10.1029/2007GL030935>.
- Marshall, A. G., H. H. Hendon, S.-W. Son, and Y. Lim, 2017: Impact of the quasi-biennial oscillation on predictability of the Madden–Julian oscillation. *Climate Dyn.*, **49**, 1365–1377, <https://doi.org/10.1007/s00382-016-3392-0>.
- Matthes, K., U. Langematz, L. L. Gray, K. Kodera, and K. Labitzke, 2004: Improved 11-year solar signal in the Freie Universität Berlin Climate Middle Atmosphere Model (FUB-CMAM). *J. Geophys. Res.*, **109**, D06101, <https://doi.org/10.1029/2003JD004012>.
- Maycock, A. C., K. Matthes, S. Tegtmeier, R. Thiéblemont, and L. Hood, 2016: The representation of solar cycle signals in stratospheric ozone—Part 1: A comparison of recently updated satellite observations. *Atmos. Chem. Phys.*, **16**, 10 021–10 043, <https://doi.org/10.5194/acp-16-10021-2016>.
- , and Coauthors, 2018: The representation of solar cycle signals in stratospheric ozone. Part II: Analysis of global models. *Atmos. Chem. Phys. Discuss.*, <https://doi.org/10.5194/acp-2017-477>, in press.
- Meehl, G., J. Arblaster, G. Branstator, and H. van Loon, 2008: A coupled air–sea response mechanism to solar forcing in the Pacific region. *J. Climate*, **21**, 2883–2897, <https://doi.org/10.1175/2007JCLI1776.1>.
- Misios, S., and Coauthors, 2016: Solar signals in CMIP-5 simulations: Effects of atmosphere–ocean coupling. *Quart. J. Roy. Meteor. Soc.*, **142**, 928–941, <https://doi.org/10.1002/qj.2695>.
- Mitchell, D. M., and Coauthors, 2015: Solar signals in CMIP-5 simulations: The stratospheric pathway. *Quart. J. Roy. Meteor. Soc.*, **141**, 2390–2403, <https://doi.org/10.1002/qj.2530>.
- Nie, J., and A. H. Sobel, 2015: Responses of tropical deep convection to the QBO: Cloud-resolving simulations. *J. Atmos. Sci.*, **72**, 3625–3638, <https://doi.org/10.1175/JAS-D-15-0035.1>.
- Nishimoto, E., and S. Yoden, 2017: Influence of the stratospheric quasi-biennial oscillation on the Madden–Julian oscillation during austral summer. *J. Atmos. Sci.*, **74**, 1105–1125, <https://doi.org/10.1175/JAS-D-16-0205.1>.
- Plumb, R. A., and R. C. Bell, 1982: A model of the quasi-biennial oscillation on an equatorial beta-plane. *Quart. J. Roy. Meteor. Soc.*, **108**, 335–352, <https://doi.org/10.1002/qj.49710845604>.
- , and J. Eluszkiewicz, 1999: The Brewer–Dobson circulation: Dynamics of the tropical upwelling. *J. Atmos. Sci.*, **56**, 868–890, [https://doi.org/10.1175/1520-0469\(1999\)056<0868:TBDCDO>2.0.CO;2](https://doi.org/10.1175/1520-0469(1999)056<0868:TBDCDO>2.0.CO;2).
- Powell, A. M., Jr., and J. Xu, 2011: Possible solar forcing of interannual and decadal stratospheric planetary wave variability in the Northern Hemisphere: An observational study. *J. Atmos. Sol-Terr. Phys.*, **73**, 825–838, <https://doi.org/10.1016/j.jastp.2011.02.001>.
- Shindell, D., D. Rind, N. Balachandran, J. Lean, and P. Lonergan, 1999: Solar cycle variability, ozone, and climate. *Science*, **284**, 305–308, <https://doi.org/10.1126/science.284.5412.305>.
- Simpson, I. R., M. Blackburn, and J. D. Haigh, 2009: The role of eddies in driving the tropospheric response to stratospheric heating perturbations. *J. Atmos. Sci.*, **66**, 1347–1365, <https://doi.org/10.1175/2008JAS2758.1>.
- Son, S.-W., Y. Lim, C. Yoo, H. Hendon, and J. Kim, 2017: Stratospheric control of the Madden–Julian oscillation. *J. Climate*, **30**, 1909–1922, <https://doi.org/10.1175/JCLI-D-16-0620.1>.
- Tiao, G., and Coauthors, 1990: Effects of autocorrelation and temporal sampling schemes on estimates of trend and spatial correlation. *J. Geophys. Res.*, **95**, 20 507–20 517, <https://doi.org/10.1029/JD095iD12p20507>.
- Wallace, J. M., and D. S. Gutzler, 1981: Teleconnections in the geopotential height field during the Northern Hemisphere winter. *Mon. Wea. Rev.*, **109**, 784–812, [https://doi.org/10.1175/1520-0493\(1981\)109<0784:TITGHF>2.0.CO;2](https://doi.org/10.1175/1520-0493(1981)109<0784:TITGHF>2.0.CO;2).
- Wheeler, M., and G. N. Kiladis, 1999: Convectively coupled equatorial waves: Analysis of clouds and temperature in the wavenumber–frequency domain. *J. Atmos. Sci.*, **56**, 374–399, [https://doi.org/10.1175/1520-0469\(1999\)056<0374:CCEWAO>2.0.CO;2](https://doi.org/10.1175/1520-0469(1999)056<0374:CCEWAO>2.0.CO;2).
- , and H. H. Hendon, 2004: An all-season real-time multivariate MJO index: Development of an index for monitoring and prediction. *Mon. Wea. Rev.*, **132**, 1917–1932, [https://doi.org/10.1175/1520-0493\(2004\)132<1917:AARMMI>2.0.CO;2](https://doi.org/10.1175/1520-0493(2004)132<1917:AARMMI>2.0.CO;2).
- Yoo, C., and S.-W. Son, 2016: Modulation of the boreal wintertime Madden–Julian oscillation by the stratospheric quasi-biennial oscillation. *Geophys. Res. Lett.*, **43**, 1392–1398, <https://doi.org/10.1002/2016GL067762>.
- Zhang, C., 2005: Madden–Julian oscillation. *Rev. Geophys.*, **43**, RG2003, <https://doi.org/10.1029/2004RG000158>.
- , 2013: Madden–Julian oscillation: Bridging weather and climate. *Bull. Amer. Meteor. Soc.*, **94**, 1849–1870, <https://doi.org/10.1175/BAMS-D-12-00026.1>.
- Zhou, J., and K.-K. Tung, 2013: Observed tropospheric temperature response to 11-yr solar cycle and what it reveals about mechanisms. *J. Atmos. Sci.*, **70**, 9–14, <https://doi.org/10.1175/JAS-D-12-0214.1>.