Eddy-Mediated Hadley Cell Expansion due to Axisymmetric Angular Momentum Adjustment to Greenhouse Gas Forcings

NICHOLAS A. DAVIS^a AND THOMAS BIRNER^b

^a *Atmospheric Chemistry Observations and Modeling Laboratory, National Center for Atmospheric Research, Boulder, Colorado*

^b *Meteorological Institute, Ludwig-Maximilians-University, Munich, Germany*

(Manuscript received 21 May 2020, in final form 20 January 2021)

ABSTRACT: The poleward expansion of the Hadley cells is one of the most robust modeled responses to increasing greenhouse gas concentrations. There are many proposed mechanisms for expansion, and most are consistent with modeled changes in thermodynamics, dynamics, and clouds. The adjustment of the eddies and the mean flow to greenhouse gas forcings, and to one another, complicates any effort toward a deeper understanding. Here we modify the Gray Radiation and Moist Aquaplanet (GRANDMA) model to uncouple the eddy and mean flow responses to forcings. When eddy forcings are held constant, the purely axisymmetric response of the Hadley cell to a greenhouse gas–like forcing is an intensification and poleward tilting of the cell with height in response to an axisymmetric increase in angular momentum in the subtropics. The angular momentum increase drastically alters the circulation response compared to axisymmetric theories, which by nature neglect this adjustment. Model simulations and an eddy diffusivity framework demonstrate that the axisymmetric increase in subtropical angular momentum—the direct manifestation of the radiative–convective equilibrium temperature response—drives a poleward shift of the eddy stresses which leads to Hadley cell expansion. Prescribing the eddy response to the greenhouse gas–like forcing shows that eddies damp, rather than drive, changes in angular momentum, moist static energy transport, and momentum transport. Expansion is not driven by changes in baroclinic instability, as would otherwise be diagnosed from the fully coupled simulation. These modeling results caution any assessment of mechanisms for circulation change within the fully coupled wave–mean flow system.

KEYWORDS: Hadley circulation; Jets; Climate change; General circulation models

1. Introduction

The poleward expansion and weakening of the Hadley cells is a robust response to greenhouse gas forcings (Frierson et al. 2007b; Medeiros et al. 2015; Grise and Polvani 2016; Davis et al. 2016; Staten et al. 2018) and idealized heating perturbations (Butler et al. 2010; Wang et al. 2012; Tandon et al. 2013) in climate model simulations. Concurrent with the expansion and weakening are increases in the meridional temperature gradient in the upper troposphere/lower stratosphere, decreases in the meridional temperature gradient at the surface (Adam et al. 2014), increases in tropical static stability (Shaw 2019), a poleward shift of the eddy-driven jet and its associated eddy fluxes of momentum and heat (Butler et al. 2010), poleward shifts in stratocumulus clouds and precipitation (Zelinka et al. 2018), drying in the subtropical troposphere (Tivig et al. 2020), and poleward shifts in surface pressure maxima and surface wind stress (Choi et al. 2014; Davis and Birner 2017; Schmidt and Grise 2017). In greenhouse gas forcing experiments these climate system responses are often unified and scale with climate sensitivity in the annual mean (Grise and Polvani 2016; Davis et al. 2016; Grise and Davis 2020). Clouds (Lipat et al. 2017), temperatures, and eddy fluxes (Davis and Birner 2016) are also associated with model biases in Hadley cell extent and intensity.

Many of the changes in the circulation over the observed record appear more consistent with multidecadal variability than forced change, complicating efforts to interpret and evaluate climate model projections (Staten et al. 2019; Grise et al. 2019).

A number of metrics for the extent of the tropical belt, a superset of metrics for the extent of the Hadley cells, have been used to infer historical changes in the Hadley cells and relate these to modeled expansion. Metrics related to temperature such as the subtropical jet and tropopause break latitudes seem to be decoupled from variability in the Hadley cell extent (Solomon et al. 2016; Davis and Birner 2017; Waugh et al. 2018), though a correlation emerges on longer time scales (Menzel et al. 2019). In other cases the variability in the metrics tend to concentrate in zonally asymmetric centers of action rather than in a unified zonally symmetric structure, such as the ridge in Northern Hemisphere sea level pressure (Schmidt and Grise 2017). The Hadley cells themselves are by construction zonally symmetric, but the climatology, variability, and long-term trends of the tropical mean circulation are zonally asymmetric, even in the Southern Hemisphere (Staten et al. 2019; Hoskins et al. 2020).

Just as there is a zoo of metrics for the extent of the tropical belt, there is a zoo of mechanisms for Hadley cell expansion (Shaw 2019; Staten et al. 2020). The Held (2000) scaling imagines the Hadley cell edge as a transition between steady tropical overturning and eddy baroclinic zones, using angular momentum conservation and the two-layer model for baroclinic instability as heuristic constraints. Its prediction that the Hadley cell extent scales with static stability is consistent with both idealized and comprehensive model simulations (Frierson et al. 2007b;

Supplemental information related to this paper is available at the Journals Online website: <https://doi.org/10.1175/jas-d-20-0149.s1>.

Corresponding author: Nicholas A. Davis, nadavis@ucar.edu

Lu et al. 2008; Kang and Lu 2012). Subtropical static stability adjusts rapidly in concert with Hadley cell expansion and the poleward shift of the eddies in abrupt forcing experiments (Chemke and Polvani 2019), so this scaling is certainly plausible even if the Hadley cell does not conserve angular momentum.

However, a separate hypothesis is that tropical upper-tropospheric warming, equivalent to increased tropospheric static stability, shifts the meridional potential vorticity gradient and its associated eddy potential vorticity fluxes poleward (Butler et al. 2011). Competing and complementary perspectives on how the eddies respond to upper-tropospheric warming and lower-stratospheric cooling and how these changes may drive expansion include supercriticality (Walker and Schneider 2006; Körtj and Schneider 2008; Levine and Schneider 2015), eddy diffusivity (Lu et al. 2014), eddy length scales (Rivière 2011), and absolutely vorticity (Kidston and Vallis 2012; Lorenz 2014). Expansion may also be driven from the extratropics through changes in eddy phase speeds (Chen and Held 2007; Chen et al. 2008), eddy length scales (Kidston et al. 2010, 2011), and poleward shifts of the eddy-driven jet in response to stratospheric cooling (Sigmond et al. 2004; Wu et al. 2013). Poleward storm track shifts and Hadley cell expansion can be related to changes in thermodynamic gradients through moist available potential energy and moist static energy, as well (Mbengue and Schneider 2013; Shaw and Voigt 2016; Mbengue and Schneider 2017, 2018; Mooring and Shaw 2020). Many of these mechanisms operate locally within the subtropics, where forcings project strongly onto Hadley cell expansion (Shaw and Tan 2018; Shaw 2019).

Nearly all of these arguments are consistent with the modeled response to forcings. One approach to make progress has been to isolate specific aspects of the forcing (Shaw 2019). Even in these targeted experiments, the eddy and mean flow responses are coupled and difficult to parse. Our starting point here is to uncouple the eddies from the mean flow within a general circulation model to gain mechanistic insights into the circulation response to changes in the atmospheric basic state in isolation.

2. The axisymmetric response to forcings as a change in the basic state

Even though it neglects eddies, the Held and Hou (1980) model of the Hadley cell is often used as a starting point to understand its essential dynamics. It imagines an energetically closed Hadley cell in an atmosphere with vanishing momentum stresses. The circulation fluxes heat and angular momentum from the tropics to the subtropics to homogenize angular momentum, satisfying Hide's theorem. Conservation of energy results in the "equal-area" constraint in which the integral of the difference between the steady-state and radiative-convective equilibrium potential temperatures must be zero. This leads to a scaling that predicts the extent of the Hadley cell as proportional to the radiative-convective equilibrium meridional temperature gradient.

By construction the steady-state meridional temperature gradient is predetermined because it must satisfy angular momentum conservation, so the radiative-convective equilibrium temperature fully determines the width of the circulation.¹ If

the radiative-convective equilibrium temperature gradient increases, the radiative-convective equilibrium temperature will intersect the steady-state temperature curve further poleward, leading to an expansion of the circulation.

In the upper troposphere of Earth's atmosphere, the steady-state meridional temperature gradient and angular momentum increase in the subtropics in response to increasing greenhouse gas concentrations. The Held and Hou (1980) model cannot support this fundamental adjustment in angular momentum. The adjustment is only possible because Earth's Hadley cell is not at the limit of angular momentum conservation due to eddy stresses (Schneider 2006), which are analogous to viscosity in the Held and Hou (1980) model (Adam and Harnik 2013) and are the proximate driver of the circulation (Davis and Birner 2019).

Suppose we relax the constraint of angular momentum conservation by introducing a heuristic parameter α , which parameterizes the eddy stress on the circulation through $u = \alpha u_M$, where u_M is the angular momentum conserving wind [Eq. (8) in Held and Hou (1980)] and $0 < \alpha \leq 1$. When $\alpha = 1$, the circulation conserves angular momentum, but if $\alpha < 1$ the circulation deviates from angular momentum conservation. It is straightforward to show that this leads to the following modified Held and Hou (1980) scaling for the Hadley cell extent ϕ_{HC} :

$$\phi_{HC} = \left(\frac{5gH\Delta_H}{3\alpha\Omega^2 a^2} \right)^{1/2} = \phi_{HH} \alpha^{-1/2}, \quad (1)$$

where g is the gravitational acceleration, H is the rigid lid height, Δ_H is the fractional change in potential temperature from the equator to the pole, Ω is the angular velocity of Earth, a is the radius of Earth, and ϕ_{HH} is the original Held and Hou (1980) scaling. The modified scaling is simply the original Held and Hou (1980) scaling multiplied by $\alpha^{-1/2}$. Eddy stresses ($\alpha < 1$) act to widen the cell (relative to $\alpha = 1$) by weakening the steady-state temperature gradient, which forces the steady-state temperature to intersect the radiative-convective equilibrium temperature further poleward.

a. Conceptual framework for the eddy-mean flow response

For the response to a greenhouse gas forcing, consider the time-mean angular momentum budget in the zonal mean for the free atmosphere,

$$[\mathbf{v}] \cdot \nabla[M] = [\tau_{\text{eddy}}], \quad (2)$$

where $[\mathbf{v}] = ([v], [\omega])$ is the zonal mean meridional and vertical wind vector, $[M] = a \cos(\phi) \{ [u] + \Omega a \cos(\phi) \}$ is the zonal mean angular momentum, $[\tau_{\text{eddy}}]$ the zonal mean eddy

¹ The other parameters in the scaling, such as gravity, rotation rate, and planetary radius, are invariant for a temperature forcing. While the rigid lid height H is often interpreted as the tropopause height, variations in H inversely modify the static stability through Eq. (2) of Held and Hou (1980), so H scales both the fluid depth and the thermodynamic profile. See also Walker and Schneider (2005) for a discussion of an unrealistic aspect of the rigid lid.

FIG. 1. Schematic of the (a) changes in basic state temperature and (b) angular momentum in response to increased greenhouse gas concentrations, (c) the axisymmetric circulation response at equilibrium, (d) the coupled circulation response at equilibrium, and (e) the eddy stress response. Solid black contours with arrows indicate the meridional circulation, while gray contours indicate the zonal wind response in (a), (c), and (d). The climatological Hadley cell edge is indicated by the dotted pink line in (a), (c), and (d), while in (c) the angular momentum contours are indicated by the solid pink lines.

angular momentum stress, which is negative within the Hadley cell, and square brackets indicate the zonal mean.

Suppose there is a global increase in greenhouse gas concentrations, which cools the lower stratosphere and warms the troposphere (see, e.g., Bony et al. 2013; Grise and Polvani 2017) (Fig. 1a). Let us first consider the axisymmetric response by holding $[\tau_{\text{eddy}}]$ constant. Note that “axisymmetric” here specifically refers to the circulation response in the absence of any eddy response, as opposed to the zonal mean response when the eddies are allowed to adjust.

Angular momentum increases in the subtropical upper troposphere at the poleward edge of the Hadley cell consistent with thermal wind adjustment to the increased meridional temperature gradient [increased Δ_H in Eq. (1)]. The increase in angular momentum reduces the meridional angular momentum gradient equatorward of the Hadley cell edge, but increases the gradient poleward of the Hadley cell edge (Fig. 1b). From Eq. (2), $[v]$ must become more positive equatorward of the Hadley cell edge and more negative poleward of the Hadley cell edge to maintain a constant value of angular momentum advection, in light of the changes in the angular momentum gradient (Fig. 1c). This results in an intensification of the Hadley and Ferrel cells.

The circulation becomes closer to angular momentum conservation [increased α in Eq. (1)] because the angular momentum increases at the poleward edge of the Hadley cell. From the perspective of a parcel, while the eddy stress within the region of subsidence is constant, the circulation is faster so that the integrated stress is weaker *and* the parcel begins from a state of elevated angular momentum in the upper troposphere (Fig. 1c). Therefore, the circulation’s angular momentum upon reaching the frictional boundary layer will be higher. As the angular momentum response increases with height above the frictional boundary layer, the circulation must reach the surface further equatorward compared to the control climate. Importantly, since the changes in α and Δ_H oppose one another, we do not necessarily expect any change in the Hadley cell width. This is unlike the unmodified Held and Hou (1980) scaling, which based on the increase in Δ_H predicts expansion.

Now, consider the coupled response in which $[\tau_{\text{eddy}}]$ can change. Eddy diffusivity arguments typically link the zonal mean eddy flux of potential vorticity to the zonal mean potential vorticity gradient (Butler et al. 2011; Lu et al. 2014). If barotropic dynamics are a reasonable approximation, the relationship simplifies to absolute vorticity. Note that the absolute vorticity is the negative of the meridional gradient of angular momentum,

$$(f + [\zeta]) = -\frac{1}{a^2 \cos(\phi)} \frac{\partial [M]}{\partial \phi}, \quad (3)$$

where f is the Coriolis parameter, $[\zeta]$ is the relative vorticity, ϕ is latitude, and p is pressure.

Supposing the axisymmetric response is the response of the basic state to the forcing, the diffusive eddy angular momentum flux and its resulting stress will shift poleward as the meridional gradient in basic state angular momentum shifts poleward (Figs. 1b,d,e). By Eq. (2), this will induce a poleward expansion of the circulation (Fig. 1d; imagine the difference of the two curves in Fig. 1e to see $\Delta[\tau_{\text{eddy}}]$ and therefore $\Delta[v]$). The reduced eddy stress in the tropics will also weaken the Hadley cell.

Formally, our hypotheses are as follows:

- 1) Through radiative–convective adjustment, an increase in the meridional temperature gradient due to upper-tropospheric warming and lower-stratospheric cooling projects an increase in angular momentum onto the subtropical jet.
- 2) Absent any changes to the eddies, this results in an (axisymmetric) intensification and poleward tilting of the Hadley cell.
- 3) The axisymmetric increase in the angular momentum gradient poleward of the Hadley cell edge and decrease equatorward of the Hadley cell edge signals a change in the zonal mean basic state and shifts the diffusive eddy stresses poleward, which produces a poleward expansion and weakening of the Hadley cell. We do not expect diffusivity to remain constant, but instead expect diffusivity changes to be secondary to the change in the basic state.

Subtropical winds and the meridional temperature gradient take longer to fully respond to abrupt forcings than the poleward shift of the eddy-driven jet and the Hadley cell edge, so an understandable criticism is that this hypothesis erroneously posits that these changes are the cause of the circulation shifts (Grise and Polvani 2017; Menzel et al. 2019). However, there are rapid responses in temperature and zonal wind separate from their long-term adjustment [see Fig. 3e of Grise and Polvani (2017), Fig. 4 of Menzel et al. (2019), and Fig. 7 of Chemke and Polvani (2019)]. Perhaps more importantly, we do not expect the axisymmetric increase in subtropical angular momentum to be fully realized in the coupled eddy–mean flow response. On the contrary, we expect it to be almost entirely diffused away by the eddies, but its dynamical forcing to remain to sustain their poleward shift (Fig. 1d). We argue the axisymmetric response is the change in the zonal mean basic state that is the *direct manifestation* of the radiative–convective equilibrium temperature response which communicates the radiative forcing to the circulation.

In this study, we will use the modeling framework developed in Davis and Birner (2019) to decouple the eddy and mean flow responses to forcings in the Gray Radiation and Moist Aquaplanet (GRANDMA) model, an extension of the Kushner and Polvani (2004), Sun et al. (2013), Lu et al. (2014), and Singh and Kuang (2016) techniques pioneered in similar idealized models. GRANDMA includes interactive gray radiation such that we simply prescribe increases in the longwave optical depth analogous to increases in well-mixed greenhouse gases, rather than

assuming temperature changes or heating perturbations that may bake in the responses of the circulation. We will show that our model experiments generally support this hypothetical framework, and demonstrate that increases in static stability, insofar as changes to baroclinic instability are concerned, do not drive an expansion of the Hadley cells.

3. Model, experiments, and methods

The idealized gray radiation aquaplanet model is described in Frierson et al. (2006) and based on the Geophysical Fluid Dynamics Laboratory dry dynamical core. “GRANDMA” refers to the configuration described in Davis and Birner (2019) with an idealized ocean heat flux scheme, an ozone-like shortwave absorption scheme, and a fixed eddy-forcing scheme. Source code modifications with instructions are permanently archived at Zenodo (Davis 2020). Radiation is modeled with bulk upward and downward longwave and shortwave fluxes with absorption specified by an emissivity that is constant across all wavelengths. We apply annual mean-like insolation as in Frierson et al. (2006) with nonzero insolation as the poles. GRANDMA includes shortwave absorption in the stratosphere meant to mimic the effect of ozone as described in Davis and Birner (2019). For the tropical and polar longwave surface optical depths we use values of 6 and 2, respectively, and use a value of 0.1 for the linear optical depth scaling f_l . See the appendix for a discussion of why this parameter should not be higher than 0.1, and why shortwave absorption by stratospheric ozone is essential to modeling the poleward shift of the eddy-driven jet.

GRANDMA models latent heating effects through large-scale condensation and a Betts–Miller convection scheme which adjusts profiles to a relative humidity of 80% on a 6-h time scale (Frierson 2007). Water vapor is advected with a finite volume parabolic scheme and initiates large-scale condensation should a grid box become supersaturated. Precipitation and reevaporation is instantaneous so that surface precipitation only occurs if the column below the level where precipitation initiates is saturated.

Surface fluxes are parameterized with bulk formulas, and vertical diffusion within the boundary layer is modeled with a simplified Monin–Obukhov similarity theory. The ocean has a fixed albedo of 0.3 and a mixed layer depth of 2.3 m such that the ocean and atmosphere have the same total heat capacity. We employ the Levine and Schneider (2011) ocean heat flux scheme which idealizes the ocean’s tropical meridional circulation as a poleward-flowing layer at the surface in Ekman balance within the zonal mean tropical easterly wind stress and an equatorward-flowing adiabatic layer below.

GRANDMA solves the vorticity–divergence form of the primitive equations on the sphere using the spectral method at T42 spectral resolution, equivalent to a 2.81° spatial resolution (with comparable results at T84 but not T21 spectral resolutions). We run the model with 40 vertical hybrid-sigma levels, with the transition to pure pressure between 300 and 200 hPa. Simulations are run for 3000 days, with the final 1000 days used in our analysis.

FIG. 2. Zonal mean climate of the control (a) coupled, (b) axisymmetric, and (c) fixed dynamics simulations. Temperatures shaded every 10 K, winds in black contours every 10 m s^{-1} (positive, solid) and 5 m s^{-1} (negative, dashed), and MMS every $30 \times 10^9 \text{ kg s}^{-1}$ in magenta contours (negative, dashed). The Hadley cell edge latitude is indicated by the asterisk, and the tropopause is indicated by the bolded black contour.

GRANDMA’s climate resembles Earth’s (Fig. 2) with well-separated eddy-driven, subtropical, and polar stratospheric jets due to the ozone scheme (Davis and Birner 2019) and a Hadley cell close to its observed intensity due to the ocean heat flux scheme (Levine and Schneider 2011).

The eddy-driven jet latitude is defined as the latitude of maximum 850-hPa zonal mean zonal wind, with a climatological value of 48° , while the latitude of the Hadley cell edge is defined as the latitude where the vertically integrated mean meridional streamfunction (MMS) vanishes poleward of its tropical maximum, with a climatological value of 29° . Here the streamfunction is defined in the conventional way as

$$\Psi = \frac{2\pi \cos(\phi)}{g} \int_0^p [v] dp. \quad (4)$$

Both metrics are calculated using the TropD software diagnostics package (Adam et al. 2018). Hadley cell intensity is assessed from the maximum streamfunction value equatorward of the edge latitude.

a. Experiments

We make extensive use of the fixed eddy-forcing axisymmetric framework developed in Davis and Birner (2019). In the fixed eddy-forcing framework, eddy fluxes of heat, moisture,

and momentum, as well as eddy impacts on surface fluxes, latent heating, and mass are calculated from a coupled simulation and prescribed as time mean, zonal mean tendencies to an axisymmetric simulation. The result is a zonal mean climate that is an analog of the coupled simulation, without explicitly modeling the eddies. There are smaller differences between the fixed eddy-forcing simulation and the coupled simulation than there were in Davis and Birner (2019), owing to our more comprehensive treatment of wind and temperature tendencies within the Simmons and Burridge (1981) vertical differencing scheme (Fig. S1 in the online supplemental material). We also use atmospheric nudging to hold the zonal mean circulation constant, applied as a Newtonian relaxation of the zonal mean with a 2-h time scale.

We assess one forcing, a doubling of the optical depth, and analyze the changes between the control (“CTRL” in figure legends) and the perturbed (“ $2\times\tau$ ” in figure legends) climates—see Table 1 for an overview of each simulation. The “3D $2\times\tau$ ” simulation models the coupled response of the eddies and mean flow to the doubling of the optical depth. In contrast, the “AXI $2\times\tau$ ” simulation holds the eddy forcings to their 3D CTRL values and resolves only the axisymmetric response to a doubling of the optical depth. In the “FIXED $2\times\tau$ ” simulation we assess the radiative–convective equilibrium response by fixing the eddy forcings, meridional wind, and surface pressure to their 3D

TABLE 1. Model parameters for the simulations. “Free” indicates the aspect of the circulation is free to evolve. “Fixed” indicates a fixed eddy-forcing (axisymmetric) simulation, and “nudged” indicates a simulation with zonal mean nudging. See the text for details.

Experiment	Optical depth	Eddy forcings	Zonal mean meridional wind, surface pressure	Zonal mean zonal wind, temperature
3D CTRL	1×	Free	Free	Free
AXI CTRL	1×	Fixed to 3D CTRL	Free	Free
FIXED CTRL	1×	Fixed to 3D CTRL	Nudged to 3D CTRL	Free
3D 2× τ	2×	Free	Free	Free
AXI 2× τ	2×	Fixed to 3D CTRL	Free	Free
FIXED 2× τ	2×	Fixed to 3D CTRL	Nudged to 3D CTRL	Free
AXI 2× τ eddies	1×	Fixed to 3D 2× τ	Free	Free
3D 2× τ AXI-nudge	1×	Free	Free	Nudged to AXI 2× τ

CTRL values. There are minor differences in the climatology between the 3D CTRL and FIXED CTRL simulations in the extratropical upper troposphere (Fig. S2).

Unlike the aforementioned 2× τ simulations that assess the responses to the doubling of the optical depth, we perform two experiments to assess the circulation response to the changes in the eddy forcings and the zonal mean basic state induced by a doubling of the optical depth. In the “AXI 2× τ eddies” simulation we set the eddy forcings to their values in the 3D 2× τ simulation, while in the “3D 2× τ AXI-nudge” simulation we set the zonal mean zonal wind and temperature to their values in the AXI 2× τ simulation. In neither simulation do we apply a doubling of the optical depth.

We will refer to the difference between the 3D 2× τ and 3D CTRL experiments as the “coupled” response, the difference between the AXI 2× τ and AXI CTRL experiments as the “axisymmetric” response, the difference between the AXI 2× τ eddies and AXI CTRL experiments as the response “induced” by the eddies, and the difference between the 3D 2× τ AXI-nudge and 3D CTRL experiments as the response “induced” by the changes in the basic state.

4. Response to a doubling of the optical depth

GRANDMA’s response to a doubling of the optical depth mimics all of the salient features of the canonical greenhouse gas response, including tropical upper-tropospheric amplification, polar amplification, and stratospheric cooling (Fig. 3a). With a global-average surface temperature warming of 12 K, the climate sensitivity to a doubling of the optical depth is about 1.5 times that of the abrupt 4×CO₂ equilibrium response in comprehensive models (Zelinka et al. 2020).

We emphasize this model has no ice–albedo or cloud–radiative feedbacks, so it seems likely that the polar amplification is due to the Planck feedback (Henry and Merlis 2019). When the longwave optical depth is increased it instantaneously reduces the outgoing longwave radiation at the top of the atmosphere, and for the sake of argument, assume this perturbation is equal at all latitudes. As the linearized response of longwave emission to an imposed radiative forcing scales as T^3 ($\Delta F = 4\sigma T_0^3 \Delta T$, where T_0 is the unperturbed temperature), the colder pole must warm more than the warmer tropics to balance the same radiative perturbation.

For unperturbed surface temperatures of 250 and 300 K, the Planck feedback predicts the pole should warm 1.75 times more than the equator, which is slightly less than what is realized (Fig. 3).

The axisymmetric response resembles the coupled response (Fig. 3b), but with clear differences in the tropical upper troposphere and at the pole. Eddies shift polar amplification (Merlis and Henry 2018) and stratospheric cooling poleward, and either directly or through feedbacks onto the mean flow, diffuse tropical upper-tropospheric warming further into the subtropics. There are only minor differences between the fixed dynamics and axisymmetric responses (Fig. 3c), confirming that the axisymmetric response is essentially the radiative–convective equilibrium response (hypothesis 1).

In response to a doubling of the optical depth, both the Hadley cell edge (Fig. 4a) and eddy-driven jet (Fig. 5a) shift poleward at a 2:3 ratio, consistent with more comprehensive models in the Southern Hemisphere (Vaugh et al. 2018). There is also an upward extension and weakening of the Hadley cell, a poleward shift of the Ferrel cell, and an upward shift and intensification of the subtropical jet—all canonical circulation responses to greenhouse gas forcings. The subtropical jet latitude does not change, but its meridional shift is generally model dependent because it the residual of its more fundamental upward shift and intensification (Shepherd and McLandress 2011; Davis and Birner 2017; Vaugh et al. 2018).

As hypothesized, the axisymmetric response is an intensification and poleward tilting of the Hadley cell (Fig. 4b; hypothesis 2). In the tropical upper troposphere the flow becomes closer to angular momentum conservation due to the increase in angular momentum in the subtropics (Fig. 5b). As a result of this reduction in the meridional gradient of angular momentum in the tropics, the Hadley cell intensifies in the upper troposphere equatorward of the maximum zonal wind increase, such that the Hadley cell strength increases by 30% [Figs. 4b, 1c; Eq. (2)]. Similarly, the Ferrel cell intensifies poleward of the maximum zonal wind increase where the meridional gradient of angular momentum increases. Above 600 hPa the Hadley cell contracts as the subsidence orients more along angular momentum contours (Fig. 1c). However, above 200 hPa the circulation expands because the maximum zonal wind response tilts poleward with height away from the climatological Hadley cell edge. The cancellation between

FIG. 3. Zonal mean temperature response to a doubling of the longwave optical depth in the (a) coupled, (b) axisymmetric, and (c) fixed dynamics simulations. Temperature response shaded every 2 K, control temperatures in black contours every 10 K, and tropopause indicated by the bolded black contour. Hadley cell edge latitude indicated by the asterisk.

contraction and expansion results in no net response in the vertically integrated Hadley cell edge.

Eddies induce Hadley cell expansion over most of the troposphere, contraction in the upper troposphere, and a weakening of more 50% (Fig. 4c). Held and Soden (2006) show that the convective mass flux weakens in warmer climates as static stability increases fractionally more than latent heating. This has been occasionally invoked to explain the weakening of the Hadley cells, but given that the atmosphere has a meridional gradient in angular momentum, any meridional circulation response must satisfy the balance between advection and internal momentum stresses (Choi et al. 2014; Davis and Birner 2019). Our results here confirm those in Feldl and Bordoni (2016), who showed that Hadley cell weakening is mostly driven by changes in the meridional eddy energy transport. The weakening of the convective mass flux in warmer climates is primarily associated with the zonal Walker circulation (Vecchi and Soden 2007), which does not face these angular momentum constraints.

Eddies also drive a weakening and poleward shift of the subtropical jet (Fig. 5c). In general the eddy-induced response is opposite to the axisymmetric response throughout most of the troposphere, which itself resembles the coupled response in the upper troposphere. On the other hand, the poleward shift of the eddy-driven jet and the weakening of the polar westerlies are driven by the eddies. The weakening of the polar westerlies is probably due to the eddies shifting high-latitude warming onto the pole (Fig. 3).

5. Response of the vertically integrated momentum budget

We now examine the responses in the vertically integrated momentum budget (Fig. 6) and moist static energy transport (Fig. 7) to develop a dynamical perspective of these circulation responses. The vertically integrated steady-state momentum budget (with units of Pa) is given by

$$0 = -\frac{\partial}{\partial \phi} \langle [u][v] \rangle - \frac{\partial}{\partial \phi} \langle [u'v'] \rangle - [\tau_{\text{drag}}] - \langle [D] \rangle, \quad (5)$$

where τ_{drag} is the surface drag, D is the hyperdiffusive damping, and primes denote deviations from the zonal mean. Angle brackets indicate the vertical integral defined as

$$\langle X \rangle = \frac{1}{g} \int_0^{p_s} X dp, \quad (6)$$

where p_s is the surface pressure, taken as 1000 hPa. In assuming a steady state, mass conservation requires the Coriolis torque term to vanish after vertical integration.

The Hadley cell fluxes momentum from the tropics to the subtropics (Fig. 6a), but the dominant global balance is between the eddy stress and surface drag (Figs. 6b,c). As the eddy-driven jet is nearly barotropic, there is little momentum transport by the Ferrel cell (Fig. 6a).

FIG. 4. Zonal mean (a) coupled MMS response to a doubling of the longwave optical depth, (b) axisymmetric MMS response to a doubling of the longwave optical depth, and (c) axisymmetric MMS response to the eddy changes from (a), but without a doubling of the longwave optical depth. MMS response shown in shading on a logarithmic scale, and control MMS in black contours every $30 \times 10^9 \text{ kg s}^{-1}$ (negative, dashed; zero contour dotted). Also shown is the Hadley cell expansion (ΔHC) and change in Hadley cell intensity ($\Delta\Psi$) for each simulation. Hadley cell edge latitude indicated by the asterisk.

The response to a doubling of the optical depth is characterized by a poleward shift of the eddy stresses and surface drag throughout the subtropics and midlatitudes, with a minor contraction away from the pole (Figs. 6b,c). There is also an amplification and poleward shift of Hadley cell momentum transport, which can be mostly explained by the axisymmetric response (Fig. 6a). As there is no change to the eddy stresses, the axisymmetric response is equilibrated by changes in surface drag and numerical damping (Figs. 6c,d).

While the midlatitude surface drag response is expectedly consistent with changes induced by the eddies (Fig. 6c), the response of the mean momentum flux convergence to the eddies is opposite to the coupled response in the tropics and subtropics and is balanced entirely by numerical damping (Figs. 6a,d).

6. Response of the vertically integrated moist static energy flux

The vertically integrated moist static energy flux with units of watts is given by

$$\langle [w_e] \rangle = c_p \langle [vT] \rangle + g \langle [vz] \rangle + L_v \langle [vq] \rangle, \quad (7)$$

where $s_e = c_p T + gz + L_v q$ is the moist static energy, $c_p = 1004 \text{ J kg}^{-1} \text{ K}$ is the specific heat of dry air, z is the geopotential height, and $L_v = 2.5 \times 10^6 \text{ J kg}^{-1}$ is the latent energy of vaporization. The sum of the first two terms on the right-hand side of Eq. (7) is the dry static energy flux, while the third term is the latent energy flux. The moist static energy flux is a fundamental energy transport in moist atmospheres, as the dry static and latent energy fluxes compensate across different hydroclimates (Frierson et al. 2007a).

The climatological moist static energy and dry static energy fluxes are poleward everywhere, though near the equator they are relatively weak due to compensation by the ocean heat flux (Figs. 7a,b). Water vapor partial pressure is an exponential function of temperature, which implies that latent energy transport will be dominated by the surface circulation. Equatorward transport in the tropics is driven by the Hadley cell, while poleward transport in the extratropics is driven by downgradient eddy mixing and the Ferrel cell (Fig. 7c).

Moist static energy transport increases and shifts poleward in response to a doubling of the optical depth (Fig. 7a), driven by an increase and poleward shift in latent energy transport (Fig. 7c) that is damped by a decrease in dry static energy transport (Fig. 7b). A reduction in the poleward moist static energy flux in the tropics is likewise driven by an enhanced

FIG. 5. Zonal mean (a) coupled zonal wind response to a doubling of the longwave optical depth, (b) axisymmetric zonal wind response to a doubling of the longwave optical depth, and (c) axisymmetric zonal wind response to the eddy changes from (a), but without a doubling of the longwave optical depth. Zonal wind response in shading on a logarithmic scale, and control zonal wind in black contours every 10 m s^{-1} (positive, solid) and 5 m s^{-1} (negative, dashed), with the zero contour dotted. Also shown is the change in the eddy-driven and subtropical jet latitudes (ΔEDJ and ΔSTJ , respectively) for each simulation.

equatorward latent energy flux and damped by an increase in the dry static energy flux.

The increased equatorward latent energy flux is consistent with the axisymmetric response and damped by the eddies (Fig. 7c). However, the increased poleward latent energy flux in the extratropics in the coupled response is stronger than in the axisymmetric response, suggesting a positive eddy-mean flow feedback on extratropical moisture transport. While the coupled response of the moist static energy flux is consistent with the axisymmetric response in the tropics, it resembles neither the axisymmetric nor eddy-driven responses in the extratropics (Fig. 7a). The coupled, axisymmetric, and eddy-driven responses in the dry static energy flux are all consistent with one another (Fig. 7b).

7. Angular momentum and the poleward shift of the eddies

We will turn to an angular momentum framework to untangle the apparent contradiction that while Hadley cell expansion must be associated with a poleward shift of the eddy stresses, eddies damp the axisymmetric thermodynamic and dynamic responses in the tropics and subtropics. Angular momentum decreases monotonically toward the poles as

required for inertial stability (Fig. 8a) and its meridional gradient peaks near 45° due to planetary rotation (Fig. 8b).

An axisymmetric increase in subtropical and decrease in midlatitude angular momentum (the latter due to polar amplification) weakens the angular momentum gradient within the Hadley cell and strengthens it poleward of the Hadley cell edge (dashed line, Figs. 8a,b). The reduction in the angular momentum gradient in the tropics is associated with a strengthening of the meridional overturning in the tropics (Fig. 8c, dashed line), whereas the overturning weakens at the edge of the circulation where the gradient strengthens, consistent with the balance in Eq. (2). This axisymmetric streamfunction response is out of phase with the coupled response (Fig. 8c, solid line), despite the similarity of their momentum and moist static energy flux responses (Figs. 6, 7).

Eddies induce a decrease in angular momentum in the subtropics, damping the axisymmetric response and strengthening the subtropical angular momentum gradient equatorward of 25° (dotted line, Figs. 8a,b). This leads to Hadley cell expansion and weakening (Fig. 8c). In the tropics and subtropics, the axisymmetric and eddy-induced responses in the streamfunction are not additive. Extratropical responses in angular momentum are generally consistent with the eddy-driven response.

FIG. 6. Vertically integrated (a) mean momentum flux convergence, (b) eddy momentum flux convergence, (c) surface drag, and (d) numerical damping in the control and perturbation simulations. See text for details on the definition of budget terms. The Hadley cell edge latitude is indicated by the asterisk. Note the control values are divided by 3 in (b) and (c).

Notice that the eddy stress anomalies are in phase with the axisymmetric and eddy-induced angular momentum responses (Figs. 8a,d), as would be the case if eddies diffuse angular momentum. Can the increase in subtropical angular momentum explain the poleward shift of the eddy stresses, and therefore the poleward expansion of the Hadley cell?

Eddies diffusing angular momentum

We can rigorously test hypothesis 3 by modeling the vertically integrated eddy stress through a diffusive closure based on vertically integrated angular momentum,²

$$\frac{\partial}{\partial \phi} [\langle [u'v'] \rangle \cos(\phi)] = - \frac{\partial}{\partial \phi} \left(D \frac{\partial}{\partial \phi} \langle [M] \rangle \right), \quad (8)$$

where D is the eddy angular momentum diffusivity. The resulting angular momentum equation [Eq. (2)] then takes

² We also examined eddy phase speed spectra in the upper troposphere and tested critical line, absolute vorticity, and jet shift arguments (Figs. S3, S4).

the form of the steady-state advection–diffusion equation. We invert D from 10 nonoverlapping 100-day samples in 3D CTRL and average D over all samples. We then apply D to the vertically integrated angular momentum gradient in the 3D $2 \times \tau$, AXI $2 \times \tau$, and AXI $2 \times \tau$ eddies experiments and compare the resulting diffusive eddy stresses to the actual change in the eddy stress in the 3D $2 \times \tau$ simulation.

The diffusive response to the coupled response in angular momentum is inconsistent with the actual eddy stress response (Fig. 9, dash-dotted and solid, respectively). However, the diffusive response to the axisymmetric increase in subtropical angular momentum drives a poleward shift in the eddy stress equal to half of its total shift (Fig. 9, dashed). It is larger in magnitude but consistent in structure and sign with the coupled response out to nearly 35° . While it is clear that eddy feedbacks will act to shape the axisymmetric response into the coupled response, this result is consistent with hypothesis 3: the initial impulse for a poleward shift of the subtropical eddy stresses is the increase in angular momentum in the axisymmetric response.

The decrease in subtropical angular momentum induced by the eddies results in a diffusive response that is opposite to the

FIG. 7. Vertically integrated (a) moist static energy flux, (b) dry static energy flux, and (c) latent energy flux in the control and perturbation experiments. See text for details on the definition of budget terms. The Hadley cell edge latitude is indicated by the asterisk. Note the control values are divided by 3 in (a) and (b).

coupled and axisymmetric responses in the tropics and subtropics, producing a 6° equatorward shift of the eddy stresses (Fig. 9, dotted line). Insofar as the eddies are diffusive, eddy feedbacks on angular momentum do not drive but instead damp the poleward shift of the eddy stresses. None of the diffusive responses can explain the eddy stress response in the midlatitudes. This could implicate an adjustment in the eddy diffusivity itself (Lu et al. 2014) or advection feedbacks (Chen et al. 2020), although the mechanisms are unclear.

Crucially, when the zonal mean zonal wind and temperature are set to the axisymmetric $2\times\tau$ response, the tropical and subtropical eddy stress responds in a similar fashion as predicted by the diffusive model with a poleward shift nearly equal to its poleward shift in the coupled response (Fig. 9, red dashed line). This validates the control diffusivity framework and demonstrates that the eddies cannot modify their behavior without feeding back onto the axisymmetric response. While diffusivity is expected to change as part of eddy-mean flow adjustment, we consider this to be secondary as the constant diffusivity assumption is sufficient to predict the poleward shift. Further, in the 3D $2\times\tau$ AXI-nudge simulation the Hadley cell expands poleward by 2.1° , compared to 1.3° in the coupled response, which suggests that the eddy feedbacks onto the axisymmetric angular momentum response work to damp Hadley cell expansion.

If one thinks of the coupled response as representing the sum of the axisymmetric (dashed) and eddy-induced (dotted) responses, then the coupled response appears to be a small residual of two large responses. However, the diffusive framework always leads to damping, so the eddy-induced response is by construction guaranteed to be opposite in sign to the axisymmetric response. Where the diffusive framework is valid, the sign of the coupled response is still given by the sign of the axisymmetric response.

8. Conclusions and discussion

An increase in the basic state subtropical angular momentum is the proximate driver of the poleward shift of the eddy stresses and the Hadley cell edge in our model simulations. In response to tropical upper-tropospheric warming and lower-stratospheric cooling, the zonal mean zonal wind and hence angular momentum increase at the edge of the Hadley cell through thermal wind adjustment (hypothesis 1). The angular momentum increase shifts the basic state absolute vorticity gradient poleward, and with it, the diffusive eddy momentum stresses (hypothesis 3). Changes in the tropical and subtropical moist static energy flux, momentum transport, and angular momentum are consistent with this axisymmetric response. Because the eddies damp these changes, we infer that the eddies primarily act to mediate and do not fundamentally

FIG. 8. Vertically averaged (a) angular momentum and (b) angular momentum gradient, (c) vertically integrated mean meridional streamfunction, and (d) vertically averaged eddy angular momentum stress. The coupled response to a doubling of the longwave optical depth, axisymmetric response to a doubling of the longwave optical depth, and axisymmetric response to the eddy changes in response to a doubling of the optical depth are shown in solid, dashed, and dotted. Control values are divided by 5 in (a) and (b) and divided by 2 in (c) and (d). The Hadley cell edge latitude is indicated by the asterisk.

drive changes in the tropical and subtropical circulation. Hadley cell expansion only occurs when the eddy angular momentum stresses shift poleward, though. Changes in the (axisymmetric) basic state angular momentum are the solution to this paradox. Scaling theories that assume angular momentum conservation neglect this important process, as by construction the angular momentum gradient is invariant.

Butler et al. (2011) similarly hypothesize that eddy fluxes respond diffusively to changes in the potential vorticity gradient and drive a poleward shift of the eddy potential vorticity flux. However, the coupled angular momentum response does not predict a poleward shift of the eddy stresses (Fig. 9). It is the embedded axisymmetric response, or the change in the zonal mean basic state which is never seen in the coupled response, that predicts this poleward shift. This is not surprising, though, as we expect the eddies to diffuse any perturbation to the basic state (Fig. 8a). If the coupled response predicted the poleward shift of the eddies, it would imply the eddies never feed back onto the mean flow, which is clearly not the case (Lu et al. 2014). Instead, it seems that the coupled response maintains roughly the same meridional shift in the eddy stress as the shift induced by the axisymmetric response (Fig. 9). Changes in eddy diffusivity may be

a part of the adjustment that propagates the basic state angular momentum changes further poleward and equatorward, leading to enhanced angular momentum everywhere except the pole (Figs. 8a,d).

Because the embedded axisymmetric response cannot be uncovered directly in coupled simulations or in nature, comparisons to previous results are not straightforward. Surprisingly, Shaw and Tan (2018) have recently shown that when they applied a CO_2 forcing only to tropical latitudes equatorward of 20° the Hadley cell did not expand. It is unclear what the change in the axisymmetric basic state looks like in these experiments, but the relative balance between upper-tropospheric warming and lower-stratospheric cooling (Fig. 3) may be important for anchoring the increase in angular momentum onto the subtropical jet to guarantee a robust poleward circulation shift. Further experiments are needed for a definitive answer.

Lu et al. (2014) find evidence that the axisymmetric response to upper-tropospheric heating should induce an equatorward shift of the eddy stresses, which would seem at odds with our results. However, the decrease in subtropical angular momentum in their experiments without eddy feedbacks is essentially opposite to the increase in subtropical angular momentum in response

FIG. 9. Vertically integrated (shading) eddy angular momentum stress, (solid black) its response to a doubling of the optical depth, (dashed red) its response to the axisymmetric response, and its responses using the control eddy angular momentum diffusivity applied to the vertically integrated angular momentum of the (dash-dotted black) coupled, (dashed black) axisymmetric, and (dotted black) eddy-driven responses. See text for details.

to increased longwave optical depth [cf. Fig. 1 of Lu et al. (2014) to Fig. 5 here]. This is likely due to the midtropospheric warming appearing as a vertical strip in the subtropics in their simulation [Fig. 1 of Lu et al. (2014)]. It seems that the meridional circulation advects the imposed upper-tropospheric heating down toward the surface which weakens the local meridional temperature gradient. This is unlike the axisymmetric response to increased longwave optical depth, in which the meridional temperature gradient increases throughout the tropics and subtropics (Fig. 3).

However, the initial radiative heating perturbation driving tropical upper-tropospheric warming is lower-tropospheric heating and upper-tropospheric cooling (Fig. 10). Prescribed upper-tropospheric heating bakes the combined moist convective and Hadley cell adjustment into the forcing itself. This damps their adjustment and contribution to the change in the basic state. Idealized heating experiments that emphasize surface heating, such as those examined in Tandon et al. (2013), may be more relevant for studying the response to increasing greenhouse gas concentrations.

As eddy diffusivity relates the eddy stress to the local angular momentum gradient [Eq. (12)], the location of the forcing is irrelevant so long as it projects onto the subtropical jet. Stronger extratropical midtropospheric warming in the Northern Hemisphere results in a weaker increase in angular momentum than in the Southern Hemisphere, which might explain the weaker Northern Hemisphere expansion (Grise and Polvani 2014; Watt-Meyer et al. 2019). In addition to the greenhouse gas problem, the angular momentum diffusivity framework may

also have some promise in explaining the atmospheric response to the narrow heating of El Niño–Southern Oscillation (Lu et al. 2008; Tandon et al. 2013).

Baroclinic instability as a mechanism for the poleward shift

Is the often-invoked Held (2000) scaling theory consistent with the dynamics we have identified here? The Held (2000) scaling posits that the poleward flow in the Hadley cell conserves angular momentum and terminates at the latitude where the zonal mean zonal wind becomes baroclinically unstable. The instability is assessed using the Phillips' two-layer criterion, and the scaling can be modified for varying initial latitudes (Kang and Lu 2012).

We can assess its relevance by examining the Phillips' instability criterion and the Eady growth rate. The Eady model assumes a continuously stratified Boussinesq atmosphere on an f plane, while the Phillips' model assumes a two-layer shallow water atmosphere on a β plane. If the Held (2000) theory captures the essence of the response to forcings, we should expect to see Eady growth rates and the Phillips' instability parameter decrease in the subtropics in the axisymmetric response. Either would indicate that changes in baroclinic instability—driven by the change in the basic state—could explain the poleward expansion of the Hadley cell. The Eady growth rate is defined as in Lindzen and Farrell (1980) as

$$\sigma = 0.31g[\rho] \frac{f}{N} \frac{\partial[u]}{\partial p}, \quad (9)$$

where N is the buoyancy frequency and $[\rho]$ is the density, while the Phillips' criterion is defined as in Held (2000) as

$$p_{\text{crit}} = \beta - \frac{f^2}{H_0} \frac{([u]_1 - [u]_2)}{g\Delta\theta}, \quad (10)$$

where H_0 is the layer thickness, $[u]_1$ and $[u]_2$ the upper- and lower-layer winds, and $\Delta\theta$ the fractional change in potential temperature in the vertical. Instability occurs should p_{crit} become negative. We split the troposphere into two layers, from 1000 to 600 hPa and from 600 to 200 hPa, and take H_0 as 5500 m (equal to half the 200-hPa geopotential height). The Eady growth rate is averaged from 1000 to 200 hPa. Time-averaged values are used to calculate these parameters so that they represent the background climate. In both instability models, vertical shear acts in favor of instability and static stability opposes instability.

The latitude of the maximum Eady growth rate and the latitude where the Phillips' instability criterion is satisfied both shift poleward by 2° in the coupled response (Fig. 11, solid lines). However, they shift equatorward by 8.3° and 3.4° in the axisymmetric response (Fig. 11, dashed lines). This shift is dominated by the shear term (cf. Fig. 11 to Figs. 5, 8), which by construction cannot change in the original Held (2000) scaling because of the assumption of angular momentum conservation.

Instead, it is the reduction in subtropical vertical shear induced by the eddies that drives a poleward shift in the latitude of the maximum Eady growth rate and the Phillips' criterion for

FIG. 10. Change in the radiative heating rate (K day^{-1}) in the (a) coupled and (b) axisymmetric responses to a doubling of the optical depth.

instability (Fig. 11, dotted lines; see also Fig. 5). The poleward shift of the baroclinic instability criteria is a result, rather than a cause, of the poleward shift of the circulation. Perhaps the reason the Held (2000) scaling appears successful is that the increase in subtropical static stability is not uniform in latitude (Figs. 3a,b) so that it describes an increase in subtropical

angular momentum through thermal wind adjustment just as easily as it describes an increase in stratification (Figs. 5a,b).

Our broader conclusion is that a diagnostic analysis of the response of the global circulation to any given forcing can give a false impression of causality and even a false sense of the fundamental mechanisms driving the response. This

FIG. 11. Change in (a) 1000–200-hPa average Eady growth rate and (b) Phillips' instability criterion. The change in latitude of the maximum Eady growth rate and latitude where the Phillips' criterion is first satisfied are displayed in the legend, and the Hadley cell edge latitude is indicated by the asterisk.

FIG. A1. Climatological zonal wind contoured every 5 m s⁻¹ (positive, solid; negative, dashed; zero contour, dotted) and response to a doubling of the optical depth (shading, logarithmic scale) in (top) simulations with ozone absorption and the ocean heat flux schemes and in (bottom) simulations without ozone absorption or ocean heat flux schemes. The τ indicates the equatorial and polar surface optical depths, while fl indicates the linear optical depth scale. The eddy-driven jet latitude and its response to a doubling of the optical depth are indicated in the upper left of each panel. See text for details.

challenge is not limited to atmospheric dynamics. It pervades all of the Earth sciences and the study of any coupled system. A modeling approach that decouples the system can provide more direct insight into how it couples. If extended to more comprehensive models, this approach could bring more clarity to our understanding of the response of the Earth system to anthropogenic and natural forcings.

Acknowledgments. The authors thank R. R. Garcia and W. J. Randel for their comments on an early version of this manuscript, and thank T. A. Shaw, E. P. Gerber, and one anonymous reviewer for the time and effort they took to help us improve this manuscript. This material is based upon work supported by the National Center for Atmospheric Research, which is a major facility sponsored by the National Science Foundation under Cooperative Agreement 1852977. Computing and data storage resources, including the Cheyenne supercomputer (doi:10.5065/

D6RX99HX), were provided by the Computational and Information Systems Laboratory (CISL) at NCAR.

Data availability statement. Modifications for the source code for GRANDMA are archived at <https://zenodo.org/record/3744495#.XpM1hnp7lQY>. Please contact the corresponding author with any questions. Model output is available upon request.

APPENDIX

Importance of Shortwave Absorption by Stratospheric Ozone

Dwyer and O’Gorman (2017), Lachmy and Shaw (2018), and Tan et al. (2019) note that the gray radiation model erroneously shifts the eddy-driven jet equatorward in response

FIG. A2. As in Fig. A1, but for equatorial and polar optical depths of 6 and 2 and a linear optical depth factor of 0.1, (a) with ozone-like shortwave absorption and the ocean heat flux schemes, (b) with ozone-like shortwave absorption but without the ocean heat flux scheme, and (c) with the ocean heat flux scheme but without ozone-like shortwave absorption.

to a warming climate. This is primarily due to the lack of an ozone-like shortwave absorption scheme in the stratosphere, and to a lesser extent the choice to run the model with a linear optical depth parameter of 0.2 (Fig. A1).

The zonal wind response to a doubling of the optical depth (Figs. 5, A1) is generally robust across variations of the surface longwave optical depth, but shifts equatorward for higher values of the linear optical depth parameter (cf. Figs. A1e–d, and Figs. A1a–c). At the same optical depth parameters, running the model with an ocean heat flux and ozone absorption scheme results in a 1.5° poleward shift in response to a doubling of the optical depth (Fig. A1b), versus a 1.5° equatorward shift with neither (Fig. A1e). We can confirm this is due to the ozone scheme, though the ocean heat flux scheme also has an impact (Fig. A2).

In particular, a distinct stratospheric polar jet shifts the zonal wind deceleration toward the pole by broadening the climatological upper-tropospheric jet (cf. Figs. A1b–e). If the climatological zonal winds over the pole are already weak, eddies with fast phase speeds may not be able to shift poleward, especially as the winds decelerate in response to polar amplification. However, if the climatological zonal winds are stronger, even with polar deceleration the fast eddies can still shift poleward.

We were unable to develop a stratospheric shortwave optical depth distribution that could shift the stratospheric jet toward the pole when the linear optical depth factor is set to 0.2 (Fig. A1c), even when we increased the shortwave optical depth at the pole by an order of magnitude. At a

linear optical depth factor of 0.2, the stratosphere is simply too effective at longwave cooling.

The implicit choice to use a linear optical depth factor of 0.2 is to decrease the otherwise unnaturally long radiative time scale of the stratosphere. However, we believe it is more physically appropriate to include shortwave absorption in the stratosphere that mimics ozone absorption. It has the dual benefits of decreasing the radiative time scale in a way consistent with the real atmosphere and producing a more realistic poleward shift of the eddy-driven jet.

REFERENCES

- Adam, O., and N. Harnik, 2013: Idealized annually averaged macro-turbulent Hadley circulation in a shallow-water model. *J. Atmos. Sci.*, **70**, 284–302, <https://doi.org/10.1175/JAS-D-12-072.1>.
- , T. Schneider, and N. Harnik, 2014: Role of changes in mean temperatures versus temperature gradients in the recent widening of the Hadley circulation. *J. Climate*, **27**, 7450–7461, <https://doi.org/10.1175/JCLI-D-14-00140.1>.
- , and Coauthors, 2018: The TropD software package (v1): Standardized methods for calculating tropical-width diagnostics. *Geosci. Model Dev.*, **11**, 4339–4357, <https://doi.org/10.5194/gmd-11-4339-2018>.
- Bony, S., G. Bellon, D. Klocke, S. Sherwood, S. Fermepin, and S. Denvil, 2013: Robust direct effect of carbon dioxide on tropical circulation and regional precipitation. *Nat. Geosci.*, **6**, 447–451, <https://doi.org/10.1038/ngeo1799>.
- Butler, A. H., D. W. J. Thompson, and R. Heikes, 2010: The steady-state atmospheric circulation response to climate

- change-like thermal forcings in a simple general circulation model. *J. Climate*, **23**, 3474–3496, <https://doi.org/10.1175/2010JCLI3228.1>.
- , —, and T. Birner, 2011: Isentropic slopes, downgradient eddy fluxes, and the extratropical atmospheric circulation response to tropical tropospheric heating. *J. Atmos. Sci.*, **68**, 2292–2305, <https://doi.org/10.1175/JAS-D-10-05025.1>.
- Chemke, R., and L. M. Polvani, 2019: Exploiting the abrupt $4 \times \text{CO}_2$ scenario to elucidate tropical expansion mechanisms. *J. Climate*, **32**, 859–875, <https://doi.org/10.1175/JCLI-D-18-0330.1>.
- Chen, G., and I. M. Held, 2007: Phase speed spectra and the recent poleward shift of Southern Hemisphere surface westerlies. *Geophys. Res. Lett.*, **34**, L21805, <https://doi.org/10.1029/2007GL031200>.
- , J. Lu, and D. M. W. Frierson, 2008: Phase speed spectra and the latitude of surface westerlies: Interannual variability and global warming trend. *J. Climate*, **21**, 5942–5959, <https://doi.org/10.1175/2008JCLI2306.1>.
- , P. Zhang, and J. Lu, 2020: Sensitivity of the latitude of the westerly jet stream to climate forcing. *Geophys. Res. Lett.*, **47**, e2019GL086563, <https://doi.org/10.1029/2019GL086563>.
- Choi, J., S.-W. Son, J. Lu, and S.-K. Min, 2014: Further observational evidence of Hadley cell widening in the Southern Hemisphere. *Geophys. Res. Lett.*, **41**, 2590–2597, <https://doi.org/10.1002/2014GL059426>.
- Davis, N., 2020: Source code modifications for the GFDL Grey Radiation Aquaplanet Model (v1.0). Zenodo, <https://doi.org/10.5281/zenodo.3744495>.
- , and T. Birner, 2016: Climate model biases in the width of the tropical belt. *J. Climate*, **29**, 1935–1954, <https://doi.org/10.1175/JCLI-D-15-0336.1>.
- , and —, 2017: On the discrepancies in tropical belt expansion between reanalyses and climate models and among tropical belt width metrics. *J. Climate*, **30**, 1211–1231, <https://doi.org/10.1175/JCLI-D-16-0371.1>.
- , and —, 2019: Eddy influences on the Hadley circulation. *J. Adv. Model. Earth Syst.*, **11**, 1563–1581, <https://doi.org/10.1029/2018MS001554>.
- , D. J. Seidel, T. Birner, S. M. Davis, and S. Tilmes, 2016: Changes in the width of the tropical belt due to simple radiative forcing changes in the GeoMIP simulations. *Atmos. Chem. Phys.*, **16**, 10083–10095, <https://doi.org/10.5194/acp-16-10083-2016>.
- Dwyer, J. G., and P. A. O’Gorman, 2017: Moist formulations of the Eliassen–Palm flux and their connection to the surface westerlies. *J. Atmos. Sci.*, **74**, 513–530, <https://doi.org/10.1175/JAS-D-16-0111.1>.
- Feldl, N., and S. Bordoni, 2016: Characterizing the Hadley circulation response through regional climate feedbacks. *J. Climate*, **29**, 613–622, <https://doi.org/10.1175/JCLI-D-15-0424.1>.
- Frierson, D. M. W., 2007: The dynamics of idealized convection schemes and their effect on the zonally averaged tropical circulation. *J. Atmos. Sci.*, **64**, 1959–1976, <https://doi.org/10.1175/JAS3935.1>.
- , I. M. Held, and P. Zurita-Gotor, 2006: A gray-radiation aquaplanet moist GCM. Part I: Static stability and eddy scale. *J. Atmos. Sci.*, **63**, 2548–2566, <https://doi.org/10.1175/JAS3753.1>.
- , —, and —, 2007a: A gray-radiation aquaplanet moist GCM. Part II: Energy transports in altered climates. *J. Atmos. Sci.*, **64**, 1680–1693, <https://doi.org/10.1175/JAS3913.1>.
- , J. Lu, and G. Chen, 2007b: Width of the Hadley cell in simple and comprehensive general circulation models. *Geophys. Res. Lett.*, **34**, L18804, <https://doi.org/10.1029/2007GL031115>.
- Grise, K. M., and L. M. Polvani, 2014: The response of midlatitude jets to increased CO_2 : Distinguishing the roles of sea surface temperature and direct radiative forcing. *Geophys. Res. Lett.*, **41**, 6863–6871, <https://doi.org/10.1002/2014GL061638>.
- , and —, 2016: Is climate sensitivity related to dynamical sensitivity? *J. Geophys. Res. Atmos.*, **121**, 5159–5176, <https://doi.org/10.1002/2015JD024687>.
- , and —, 2017: Understanding the time scales of the tropospheric circulation response to abrupt CO_2 forcing in the Southern Hemisphere: Seasonality and the role of the stratosphere. *J. Climate*, **30**, 8497–8515, <https://doi.org/10.1175/JCLI-D-16-0849.1>.
- , and S. M. Davis, 2020: Hadley cell expansion in CMIP6 models. *Atmos. Chem. Phys.*, **20**, 5249–5268, <https://doi.org/10.5194/acp-20-5249-2020>.
- , and Coauthors, 2019: Recent tropical expansion: Natural variability or forced response? *J. Climate*, **32**, 1551–1571, <https://doi.org/10.1175/JCLI-D-18-0444.1>.
- Held, I. M., 2000: The general circulation of the atmosphere. Woods Hole Oceanographic Institute, 70 pp., https://www.gfdl.noaa.gov/wp-content/uploads/files/user_files/ih/lectures/woods_hole.pdf.
- , and A. Y. Hou, 1980: Nonlinear axially symmetric circulations in a nearly inviscid atmosphere. *J. Atmos. Sci.*, **37**, 515–533, [https://doi.org/10.1175/1520-0469\(1980\)037<0515:NASCIA>2.0.CO;2](https://doi.org/10.1175/1520-0469(1980)037<0515:NASCIA>2.0.CO;2).
- , and B. J. Soden, 2006: Robust responses of the hydrological cycle to global warming. *J. Climate*, **19**, 5686–5699, <https://doi.org/10.1175/JCLI3990.1>.
- Henry, M., and T. M. Merlis, 2019: The role of the nonlinearity of the Stefan–Boltzmann law on the structure of radiatively forced temperature change. *J. Climate*, **32**, 335–348, <https://doi.org/10.1175/JCLI-D-17-0603.1>.
- Hoskins, B. J., G.-Y. Yang, and R. M. Fonseca, 2020: The detailed dynamics of the June–August Hadley cell. *Quart. J. Roy. Meteor. Soc.*, **146**, 557–575, <https://doi.org/10.1002/qj.3702>.
- Kang, S. M., and J. Lu, 2012: Expansion of the Hadley cell under global warming: Winter versus summer. *J. Climate*, **25**, 8387–8393, <https://doi.org/10.1175/JCLI-D-12-00323.1>.
- Kidston, J., and G. K. Vallis, 2012: The relationship between the speed and the latitude of an eddy-driven jet in a stirred barotropic model. *J. Atmos. Sci.*, **69**, 3251–3263, <https://doi.org/10.1175/JAS-D-11-0300.1>.
- , S. M. Dean, J. A. Renwick, and G. K. Vallis, 2010: A robust increase in the eddy length scale in the simulation of future climates. *Geophys. Res. Lett.*, **37**, L03806, <https://doi.org/10.1029/2009GL041615>.
- , G. K. Vallis, S. M. Dean, and J. A. Renwick, 2011: Can the increase in the eddy length scale under global warming cause the poleward shift of the jet streams? *J. Climate*, **24**, 3764–3780, <https://doi.org/10.1175/2010JCLI3738.1>.
- Korty, R. L., and T. Schneider, 2008: Extent of Hadley circulations in dry atmospheres. *Geophys. Res. Lett.*, **35**, L23803, <https://doi.org/10.1029/2008GL035847>.
- Kushner, P. J., and L. M. Polvani, 2004: Stratosphere-troposphere coupling in a relatively simple AGCM: The role of eddies. *J. Climate*, **17**, 629–639, [https://doi.org/10.1175/1520-0442\(2004\)017<0629:SCIARS>2.0.CO;2](https://doi.org/10.1175/1520-0442(2004)017<0629:SCIARS>2.0.CO;2).
- Lachmy, O., and T. Shaw, 2018: Connecting the energy and momentum flux response to climate change using the Eliassen–Palm relation. *J. Climate*, **31**, 7401–7416, <https://doi.org/10.1175/JCLI-D-17-0792.1>.

- Levine, X. J., and T. Schneider, 2011: Response of the Hadley circulation to climate change in an aquaplanet GCM coupled to a simple representation of ocean heat transport. *J. Atmos. Sci.*, **68**, 769–783, <https://doi.org/10.1175/2010JAS3553.1>.
- , and —, 2015: Baroclinic eddies and the extent of the Hadley circulation: An idealized GCM study. *J. Atmos. Sci.*, **72**, 2744–2761, <https://doi.org/10.1175/JAS-D-14-0152.1>.
- Lindzen, R. S., and B. Farrell, 1980: A simple approximate result for the maximum growth rate of baroclinic instabilities. *J. Atmos. Sci.*, **37**, 1648–1654, [https://doi.org/10.1175/1520-0469\(1980\)037<1648:ASARFT>2.0.CO;2](https://doi.org/10.1175/1520-0469(1980)037<1648:ASARFT>2.0.CO;2).
- Lipat, B. R., G. Tselioudis, K. M. Grise, and L. M. Polvani, 2017: CMIP5 models' shortwave cloud radiative response and climate sensitivity linked to the climatological Hadley cell extent. *Geophys. Res. Lett.*, **44**, 5739–5748, <https://doi.org/10.1002/2017GL073151>.
- Lorenz, D. J., 2014: Understanding midlatitude jet variability and change using Rossby wave chromatography: Poleward-shifted jets in response to external forcing. *J. Atmos. Sci.*, **71**, 2370–2389, <https://doi.org/10.1175/JAS-D-13-0200.1>.
- Lu, J., G. Chen, and D. M. W. Frierson, 2008: Response of the zonal mean atmospheric circulation to El Niño versus global warming. *J. Climate*, **21**, 5835–5851, <https://doi.org/10.1175/2008JCLI2200.1>.
- , L. Sun, Y. Wu, and G. Chen, 2014: The role of subtropical irreversible PV mixing in the zonal mean circulation response to global warming-like thermal forcing. *J. Climate*, **27**, 2297–2316, <https://doi.org/10.1175/JCLI-D-13-00372.1>.
- Mbengue, C., and T. Schneider, 2013: Storm track shifts under climate change: What can be learned from large-scale dry dynamics. *J. Climate*, **26**, 9923–9930, <https://doi.org/10.1175/JCLI-D-13-00404.1>.
- , and —, 2017: Storm-track shifts under climate change: Toward a mechanistic understanding using baroclinic mean available potential energy. *J. Atmos. Sci.*, **74**, 93–110, <https://doi.org/10.1175/JAS-D-15-0267.1>.
- , and —, 2018: Linking Hadley circulation and storm tracks in a conceptual model of the atmospheric energy balance. *J. Atmos. Sci.*, **75**, 841–856, <https://doi.org/10.1175/JAS-D-17-0098.1>.
- Medeiros, B., B. Stevens, and S. Bony, 2015: Using aquaplanets to understand the robust responses of comprehensive climate models to forcing. *Climate Dyn.*, **44**, 1957–1977, <https://doi.org/10.1007/s00382-014-2138-0>.
- Menzel, M. E., D. Waugh, and K. Grise, 2019: Disconnect between Hadley cell and subtropical jet variability and response to increased CO₂. *Geophys. Res. Lett.*, **46**, 7045–7053, <https://doi.org/10.1029/2019GL083345>.
- Merlis, T. M., and M. Henry, 2018: Simple estimates of polar amplification in moist diffusive energy balance models. *J. Climate*, **31**, 5811–5824, <https://doi.org/10.1175/JCLI-D-17-0578.1>.
- Mooring, T. A., and T. A. Shaw, 2020: Atmospheric diffusivity: A new energetic framework for understanding the midlatitude circulation response to climate change. *J. Geophys. Res. Atmos.*, **125**, e2019JD031206, <https://doi.org/10.1029/2019JD031206>.
- Rivière, G., 2011: A dynamical interpretation of the poleward shift of the jet streams in global warming scenarios. *J. Atmos. Sci.*, **68**, 1253–1272, <https://doi.org/10.1175/2011JAS3641.1>.
- Schmidt, D. F., and K. M. Grise, 2017: The response of local precipitation and sea level pressure to Hadley cell expansion. *Geophys. Res. Lett.*, **44**, 10 573–10 582, <https://doi.org/10.1002/2017GL075380>.
- Schneider, T., 2006: The general circulation of the atmosphere. *Ann. Rev. Earth Planet. Sci.*, **34**, 655–688, <https://doi.org/10.1146/annurev.earth.34.031405.125144>.
- Shaw, T. A., 2019: Mechanisms of future predicted changes in the zonal mean mid-latitude circulation. *Curr. Climate Change Rep.*, **5**, 345–357, <https://doi.org/10.1007/s40641-019-00145-8>.
- , and A. Voigt, 2016: What can moist thermodynamics tell us about circulation shifts in response to uniform warming? *Geophys. Res. Lett.*, **43**, 4566–4575, <https://doi.org/10.1002/2016GL068712>.
- , and Z. Tan, 2018: Testing latitudinally dependent explanations of the circulation response to increased CO₂ using aquaplanet models. *Geophys. Res. Lett.*, **45**, 9861–9869, <https://doi.org/10.1029/2018GL078974>.
- Shepherd, T. G., and C. McLandress, 2011: A robust mechanism for strengthening of the Brewer–Dobson circulation in response to climate change: Critical-layer control of subtropical wave breaking. *J. Atmos. Sci.*, **68**, 784–797, <https://doi.org/10.1175/2010JAS3608.1>.
- Sigmond, M., P. C. Siegmund, E. Manzini, and H. Kelder, 2004: A simulation of the separate climate effects of middle-atmospheric and tropospheric CO₂ doubling. *J. Climate*, **17**, 2352–2367, [https://doi.org/10.1175/1520-0442\(2004\)017\(2352:ASOTSC\)2.0.CO;2](https://doi.org/10.1175/1520-0442(2004)017(2352:ASOTSC)2.0.CO;2).
- Simmons, A. J., and D. M. Burridge, 1981: An energy and angular-momentum conserving vertical finite-difference scheme and hybrid vertical coordinates. *Mon. Wea. Rev.*, **109**, 758–766, [https://doi.org/10.1175/1520-0493\(1981\)109%3C0758:AEAAMC%3E2.0.CO;2](https://doi.org/10.1175/1520-0493(1981)109%3C0758:AEAAMC%3E2.0.CO;2).
- Singh, M. S., and Z. Kuang, 2016: Exploring the role of eddy momentum fluxes in determining the characteristics of the equinoctial Hadley circulation: Fixed-SST simulations. *J. Atmos. Sci.*, **73**, 2427–2444, <https://doi.org/10.1175/JAS-D-15-0212.1>.
- Solomon, A., L. M. Polvani, D. W. Waugh, and S. M. Davis, 2016: Contrasting upper and lower atmospheric metrics of tropical expansion in the Southern Hemisphere. *Geophys. Res. Lett.*, **43**, 10 496–10 503, <https://doi.org/10.1002/2016GL070917>.
- Staten, P. W., J. Lu, K. M. Grise, S. M. Davis, and T. Birner, 2018: Re-examining tropical expansion. *Nat. Climate Change*, **8**, 768–775, <https://doi.org/10.1038/s41558-018-0246-2>.
- , K. M. Grise, S. M. Davis, K. Karneuska, and N. Davis, 2019: Regional widening of tropical overturning: Forced change, natural variability, and recent trends. *J. Geophys. Res. Atmos.*, **124**, 6104–6119, <https://doi.org/10.1029/2018JD030100>.
- , and Coauthors, 2020: Tropical widening: From global variations to regional impacts. *Bull. Amer. Meteor. Soc.*, **101**, E897–E904, <https://doi.org/10.1175/BAMS-D-19-0047.1>.
- Sun, L., G. Chen, and J. Lu, 2013: Sensitivities and mechanisms of the zonal mean atmospheric circulation response to tropical warming. *J. Atmos. Sci.*, **70**, 2487–2504, <https://doi.org/10.1175/JAS-D-12-0298.1>.
- Tan, Z., O. Lachmy, and T. A. Shaw, 2019: The sensitivity of the jet stream response to climate change to radiative assumptions. *J. Adv. Model. Earth Syst.*, **11**, 934–956, <https://doi.org/10.1029/2018MS001492>.
- Tandon, N. F., E. P. Gerber, A. H. Sobel, and L. M. Polvani, 2013: Understanding Hadley cell expansion versus contraction: Insights from simplified models and implications for recent observations. *J. Climate*, **26**, 4304–4321, <https://doi.org/10.1175/JCLI-D-12-00598.1>.
- Tivig, M., V. Grutzun, V. O. John, and S. A. Buehler, 2020: Trends in upper-tropospheric humidity: Expansion of the

- subtropical dry zones? *J. Climate*, **33**, 2149–2161, <https://doi.org/10.1175/JCLI-D-19-0046.1>.
- Vecchi, G. A., and B. J. Soden, 2007: Global warming and the weakening of the tropical circulation. *J. Climate*, **20**, 4316–4340, <https://doi.org/10.1175/JCLI4258.1>.
- Walker, C. C., and T. Schneider, 2005: Response of idealized Hadley circulations to seasonally varying heating. *Geophys. Res. Lett.*, **32**, L06813, <https://doi.org/10.1029/2004GL022304>.
- , and —, 2006: Eddy influences on Hadley circulations: Simulations with an idealized GCM. *J. Atmos. Sci.*, **63**, 3333–3350, <https://doi.org/10.1175/JAS3821.1>.
- Wang, S., E. P. Gerber, and L. M. Polvani, 2012: Abrupt circulation responses to tropical upper-tropospheric warming in a relatively simple stratosphere-resolving AGCM. *J. Climate*, **25**, 4097–4115, <https://doi.org/10.1175/JCLI-D-11-00166.1>.
- Watt-Meyer, O., D. M. W. Frierson, and Q. Fu, 2019: Hemispheric asymmetry of tropical expansion under CO₂ forcing. *Geophys. Res. Lett.*, **46**, 9231–9240, <https://doi.org/10.1029/2019GL083695>.
- Waugh, D. W., and Coauthors, 2018: Revisiting the relationship among metrics of tropical expansion. *J. Climate*, **31**, 7565–7581, <https://doi.org/10.1175/JCLI-D-18-0108.1>.
- Wu, Y., R. Seager, T. A. Shaw, M. Ting, and N. Naik, 2013: Atmospheric circulation response to an instantaneous doubling of carbon dioxide. Part II: Atmospheric transient adjustment and its dynamics. *J. Climate*, **26**, 918–935, <https://doi.org/10.1175/JCLI-D-12-00104.1>.
- Zelinka, M. D., K. M. Grise, S. A. Klein, C. Zhou, A. M. DeAngelis, and M. W. Christensen, 2018: Drivers of the low-cloud response to poleward jet shifts in the North Pacific in observations and models. *J. Climate*, **31**, 7925–7947, <https://doi.org/10.1175/JCLI-D-18-0114.1>.
- , T. A. Myers, D. T. McCoy, S. Po-Chedley, P. M. Caldwell, P. Ceppi, S. A. Klein, and K. E. Taylor, 2020: Causes of higher climate sensitivity in CMIP6 models. *Geophys. Res. Lett.*, **47**, e2019GL085782, <https://doi.org/10.1029/2019GL085782>.