New NASA Missions Focus on Terrestrial Forcing of the Space Environment

T. J. IMMEL AND R. W. EASTES

FIG. 1. Simulated airglow emissions as observed by GOLD. (left) Disk image of the 135.6-nm emission, showing dayglow, nightglow, and aurora. Landmass boundaries are shown from geostationary orbit at 47.5°W longitude, where GOLD will be stationed. (right) Spectral image in wavelength and position along the slit (shown as white rectangle on the disk image). (Figure courtesy S. C. Solomon, NCAR/HAO.)

In late 2019, two ground-breaking NASA missions to explore the conditions in near-Earth space environment are expected to both be on orbit and operational. These missions are designed to observe changes in the state of the neutral upper

atmosphere and ionosphere (at 100–500-km altitude) in response to forcing from the sun and solar wind, as well as the terrestrial atmosphere below. Together these observatories will provide a comprehensive view of the near-Earth space environment, the conditions of which can be broadly described as space weather, and explore the degree to which the lower and middle atmosphere exert a significant, possibly controlling, influence. These missions provide complementary capabilities for quantifying the drivers of the near-Earth space environment and its responses at low and middle latitudes. The comprehensive view provided by these missions will lead to a much improved understanding of space weather, and inform the international effort to improve prediction of conditions in Earth's space environment, both in terms of its neutral and ionized components.

AFFILIATIONS: IMMEL—Space Science Laboratory, University of California, Berkeley, Berkeley, California; EASTES—Laboratory for Atmospheric and Space Physics, University of Colorado Boulder, Boulder, Colorado

CORRESPONDING AUTHOR: T. J. Immel, immel@ssl.berkeley.edu

DOI:10.1175/BAMS-D-19-0066.1

©2019 American Meteorological Society

For information regarding reuse of this content and general copyright information, consult the [AMS Copyright Policy](#).

TWO NEW SPACE MISSIONS. The Global-Scale Observations of the Limb and Disk Imager (GOLD; Fig. 1) (Eastes et al. 2017) was launched in January 2018. This NASA mission of opportunity is flying a pair of identical, imaging spectrometers on *SES-14*, a communications satellite in geostationary orbit over South America at 47.5°W. GOLD makes spectrally and spatially resolved scans of the far-ultraviolet (FUV) emissions (132–162 nm) from the upper atmosphere and ionosphere. Earth's daytime disk and limb are observed with a 30-min cadence. From these observations the first full-disk images of both the neutral temperatures and composition in the upper atmosphere are retrieved. Simultaneous imaging of both quantities during the dayside observations provide a never-before-seen view of the short-term changes that occur across the American continents and the Atlantic Ocean. At night, GOLD measures the intensity of ultraviolet emissions of atomic oxygen, from which the maximum ionospheric density can be directly retrieved.

The Ionospheric Connection Explorer (ICON; Fig. 2) mission (Immel et al. 2018) is prepared for a 2019 launch and will retrieve a comprehensive set of characteristics of the neutral upper atmosphere and ionosphere. Implementing an innovative heterodyne imaging technique (Harlander et al. 2017), ICON retrieves wind profiles in the upper atmosphere using a Michelson interferometer. In daytime, these wind profiles extend from the edge of space at 95 km to the ionospheric peak above 300 km. Simultaneously, ICON makes in situ plasma density and 3D velocity measurements (at 575 km, the satellite's target altitude) using an ion velocity meter. By the design of the mission, these measurements are often connected by Earth's magnetic field, where near the magnetic equator a magnetic field line at the spacecraft can be followed directly down to the region of the remote measurement. Given the high electrical conductivity in this region, the electric potential that is, in part, generated by the dynamo action of the

FIG. 2. Key observations during a pass of ICON over the Pacific. Products are provided from the same physical region within a 7-min window. The cardinal wind and temperature profiles come from interferometric observations from positions A and D. The neutral and ion composition profiles are provided at position B. Electric fields that originate from the region of interest are measured continuously (green line) and specifically measured at position C. Earth's magnetic field is described by arched blue lines, illustrating the connection between the in situ and remote measurements.

remotely observed wind is measured at the spacecraft by measuring the plasma velocity field. In this way, the relationship between the neutral and plasma velocity fields can be evaluated for the first time. In addition, height profiles of plasma density in the daytime are retrieved from daytime measurements by an extreme ultraviolet spectrometer. ICON also carries an FUV imager for retrieval of atmospheric composition and ionospheric density profiles in day and night, respectively, providing a high-resolution local complement to the large-scale imaging by GOLD.

The combination of local and global observations from these two missions provides data that can enable new assimilative and physics-based approaches for predictions of the conditions in Earth's "space weather." Together the two missions represent a corollary to that used by the meteorological community, combining global-scale contextual observations from geosynchronous orbit with critical, focused in situ observations at smaller scales.

Improvement in the reliability and accuracy of space weather forecasts is enabled by increasing the number and breadth of key state variables measured and made available for analysis and assimilation. Space weather conditions are known to respond strongly to variability in solar UV irradiance, solar energetic particle flux, and bulk plasma emanating from the sun; the last of these reaching Earth in a magnetized solar wind. Knowledge of the solar radiation and solar wind output is a national interest that is currently fulfilled by NOAA's space assets including Deep Space Climate Observatory (DSCOVR; Burt and Smith 2012) with planned continuation follow-on missions. However, it is now clear that the transfer of energy and momentum from the troposphere and stratosphere upward into Earth's upper atmosphere is likely a defining process in space weather (Immel et al. 2006; Hagan et al. 2007; Liu et al. 2013). Together, ICON and GOLD provide more than a dozen related products to help resolve the mystery of day to day variations in space weather that are apparently not related to the currently measured solar wind key parameters and therefore not predictable at this time.

TERRESTRIAL ATMOSPHERIC DRIVERS OF SPACE WEATHER.

There are several atmospheric phenomena that are now thought to be of key importance when considering the connection of the terrestrial atmosphere to space weather. Examples include atmospheric tides energized in the lower atmosphere that propagate to much higher altitudes. Several

upward-propagating atmospheric tides with origins in the troposphere grow to their largest amplitudes in the 90–120-km range before breaking or dissipating (Hagan and Forbes 2002). Tropospheric forcing, therefore, extends to and well above the boundary of space and into the plasma of the lower ionosphere. The largest of these tides is energized by absorption of solar infrared radiation by water vapor and latent heat released by cloud formation. These tides can therefore vary over short time periods as weather patterns change, varying more rapidly than does the relatively steady diurnal tide generated in the thermosphere by the absorption of solar extreme UV radiation. In addition, gravity and acoustic waves originating from tropospheric sources also propagate into the region, with some reaching the ionospheric peak near 300–400 km. Perhaps most spectacular are global-scale disruptions of the ionosphere that have been attributed to large changes in the planetary wave spectrum in the polar stratosphere [see Goncharenko et al. (2010) for an excellent example]. The changes in the ionosphere observed during major stratospheric warming events, with both enhancements and reductions of plasma density as large as a factor of 2, over locations at low or middle latitudes, are of a magnitude comparable to extreme space weather events driven by the solar wind.

CONCLUSIONS. Although measurements and models of the space environment have advanced significantly in recent decades, the needs for prediction still exceed current capabilities. The ICON and GOLD missions will provide unprecedented measurements that will advance our understanding of the behavior of the near-Earth space environment and open avenues for improvement in scientific capability in space weather prediction. As each mission seeks to discern the influence of terrestrial drivers on Earth's upper atmosphere, combining these new observations with those from other NASA missions, U.S. agencies, and international partners holds promise for even more comprehensive views of the system. The ICON and GOLD missions were designed separately, but carry naturally synergistic capabilities that are soon to be realized.

FOR FURTHER READING

Burt, J., and B. Smith, 2012: The Deep Space Climate Observatory: The DSCOVR Mission. 2012 IEEE Aerospace Conference, Big Sky, MT, IEEE, 1–13, <https://doi.org/10.1109/AERO.2012.6187025>.

Eastes, R. W., and Coauthors, 2017: The Global-Scale Observations of the Limb and Disk (GOLD) Mission. *Space Sci. Rev.*, **212**, 383–408, <https://doi.org/10.1007/s11214-017-0392-2>.

Goncharenko, L. P., J. L. Chau, H.-L. Liu, and A. J. Coster, 2010: Unexpected connections between the stratosphere and ionosphere. *Geophys. Res. Lett.*, **37**, L10101, <https://doi.org/10.1029/2010GL043125>.

Hagan, M. E., and J. M. Forbes, 2002: Migrating and nonmigrating diurnal tides in the middle and upper atmosphere excited by tropospheric latent heat release. *J. Geophys. Res.*, **107**, 4754, <https://doi.org/10.1029/2001JD001236>.

—, A. Maute, R. G. Roble, A. D. Richmond, T. J. Immel, and S. L. England, 2007: The effects of deep tropical clouds on the earth's ionosphere. *Geophys. Res. Lett.*, **34**, L20109, <https://doi.org/10.1029/2007GL030142>.

Harlander, J. M., C. R. Englert, C. M. Brown, K. D. Marr, I. J. Miller, V. Zastera, B. W. Bach, and S. B. Mende,

2017: Michelson Interferometer for Global High-Resolution Thermospheric Imaging (MIGHTI): Monolithic interferometer design and test. *Space Sci. Rev.*, **212**, 601, <https://doi.org/10.1007/s11214-017-0374-4>.

Immel, T. J., and Coauthors, 2006: The control of equatorial ionospheric morphology by atmospheric tides. *Geophys. Res. Lett.*, **33**, 15108, <https://doi.org/10.1029/2006GL026161>.

—, and Coauthors, 2018: The Ionospheric Connection Explorer Mission: Mission goals and design. *Space Sci. Rev.*, **214**, 13, <https://doi.org/10.1007/s11214-017-0449-2>.

Liu, H.-L., V. A. Yudin, and R. G. Roble, 2013: Day-to-day ionospheric variability due to lower atmosphere perturbations. *Geophys. Res. Lett.*, **40**, 665–670, <https://doi.org/10.1002/grl.50125>.

Richmond, A. D., 1989: Modeling the ionosphere wind dynamo: A review. *Pure Appl. Geophys.*, **131**, 413–435, <https://doi.org/10.1007/BF00876837>.

Interpret and Forecast Weather with The WeatherCycler

Purchase for your home or classroom

The WeatherCycler
For Interpreting and Forecasting Your Weather
American Meteorological Society Education Program
ametsoc.org/ametsedu

INSTRUCTIONS:
Consult the latest television or newspaper weather map and pull the slide until A or B points to the station best showing your location in the HIGH or LOW affecting your weather. The Surface Weather Map and the windows above it describe your weather. If Point A indicates your location in the HIGH or LOW, use the windows below if Point B indicates your position.

To forecast your weather, pull the slide out slowly to represent movement of HIGH or LOW. Your changing map location shows how your weather will develop. The next plotted position shows what your weather may be in about 12 hours. Check your forecasts by following the instructions on back.

COMMON WEATHER MAP SYMBOLS

ISOBARS give form to regions and LOWS and connect places with the same air pressure.

SHADED AREAS indicate precipitation—rain or snow.

Center of LOW

WEATHER STATION: Indicates direction from which wind is blowing. **Observed WIND SPEED:** Feathers are fitted: 1 to 10 knots, 11 to 20 knots, 21 to 30 knots, 31 to 40 knots, 41 to 50 knots, 51 to 60 knots, 61 to 70 knots, 71 to 80 knots, 81 to 90 knots, 91 to 100 knots, 101 to 110 knots, 111 to 120 knots, 121 to 130 knots, 131 to 140 knots, 141 to 150 knots, 151 to 160 knots, 161 to 170 knots, 171 to 180 knots, 181 to 190 knots, 191 to 200 knots, 201 to 210 knots, 211 to 220 knots, 221 to 230 knots, 231 to 240 knots, 241 to 250 knots, 251 to 260 knots, 261 to 270 knots, 271 to 280 knots, 281 to 290 knots, 291 to 300 knots, 301 to 310 knots, 311 to 320 knots, 321 to 330 knots, 331 to 340 knots, 341 to 350 knots, 351 to 360 knots, 361 to 370 knots, 371 to 380 knots, 381 to 390 knots, 391 to 400 knots, 401 to 410 knots, 411 to 420 knots, 421 to 430 knots, 431 to 440 knots, 441 to 450 knots, 451 to 460 knots, 461 to 470 knots, 471 to 480 knots, 481 to 490 knots, 491 to 500 knots, 501 to 510 knots, 511 to 520 knots, 521 to 530 knots, 531 to 540 knots, 541 to 550 knots, 551 to 560 knots, 561 to 570 knots, 571 to 580 knots, 581 to 590 knots, 591 to 600 knots, 601 to 610 knots, 611 to 620 knots, 621 to 630 knots, 631 to 640 knots, 641 to 650 knots, 651 to 660 knots, 661 to 670 knots, 671 to 680 knots, 681 to 690 knots, 691 to 700 knots, 701 to 710 knots, 711 to 720 knots, 721 to 730 knots, 731 to 740 knots, 741 to 750 knots, 751 to 760 knots, 761 to 770 knots, 771 to 780 knots, 781 to 790 knots, 791 to 800 knots, 801 to 810 knots, 811 to 820 knots, 821 to 830 knots, 831 to 840 knots, 841 to 850 knots, 851 to 860 knots, 861 to 870 knots, 871 to 880 knots, 881 to 890 knots, 891 to 900 knots, 901 to 910 knots, 911 to 920 knots, 921 to 930 knots, 931 to 940 knots, 941 to 950 knots, 951 to 960 knots, 961 to 970 knots, 971 to 980 knots, 981 to 990 knots, 991 to 1000 knots.

EXAMPLE: Northeast wind at 15 knots. Overcast.

FRONTS form the boundary between neighboring air masses (HIGHS), important and rapid weather changes occur across fronts. Symbols indicate kind of front and direction of movement.

Fronts: Cold Front, Warm Front, Stationary Front.

Weather at A: High, Low, Clouds, Air Pressure and Temperature Changes, Side View of Atmosphere.

Surface Weather Map: Shows weather systems passing by.

Weather at B: High, Low, Clouds, Air Pressure and Temperature Changes, Side View of Atmosphere.

A durable 8 1/2" x 11" slide chart that uniquely displays map views and vertical cross sections through Highs, Lows, and fronts—the broad-scale systems and features that dominate our weather. Sky views and changes in temperature and pressure complete the comprehensive chart depictions. By simply pulling the chart insert, the weather systems are put into their typical movements from west to east. Weather changes are shown for your location as the weather systems pass by. Accompanying four-page instruction booklet describes the basic causes of weather and explains step-by-step how to use The WeatherCycler.

\$9.99 INDIVIDUAL KIT
\$4 S&H

\$169.95 CLASSROOM KIT
+\$45 S&H (15 student, 1 instructor)

To Order (check, PO, credit card)

Phone: 1-800-824-0405
Fax: 1-800-258-1176

Learn more: ametsoc.org/EducationMaterials

Mail: American Meteorological Society Education Program
1200 New York Ave., NW, Suite 500,
Washington, DC 20005-3928

Questions: amsedu@ametsoc.org