OUR SKILL IN MODELING MOUNTAIN RAIN AND SNOW IS BYPASSING THE SKILL OF OUR OBSERVATIONAL NETWORKS

JESSICA LUNDQUIST, MIMI HUGHES, ETHAN GUTMANN, AND SARAH KAPNICK

In mountainous areas, high-resolution atmospheric models can represent total annual precipitation better than the collective network of precipitation gauges.

We have now crossed a threshold where, for many mountain ranges, well-configured high-resolution atmospheric models are better able to represent range-wide total annual precipitation than the collective network of precipitation gauges: that is, observations. The prior sentence is disturbing. If we even assign some truth to the statement “Models

are better than observations,” where does that lead us? If two models are “better than observations” but disagree with each other, which one should we trust more? What do we mean by “better,” and what counts as an “observation”? Generalities are dangerous, and for which models, which observations, and which times and locations might this statement be true? How do we identify these specificities in a way that allows us to move forward, scientifically and objectively, in a situation where the *truth* is difficult to discern?

Here, we review recent research that collectively suggests, at least for the mid- to northern latitudes, that modeled precipitation has crossed a threshold in range-wide accuracy relative to observation-based precipitation datasets in complex terrain. We carefully examine the basis for this conclusion, which often consists of multiple indirect observations. We then propose that crossing this threshold requires a fundamental shift in how hydrologists and atmospheric scientists interact. In the past, gridded precipitation datasets that interpolated between existing observations, such as PRISM (Daly et al.

AFFILIATIONS: LUNDQUIST—Civil and Environmental Engineering, University of Washington, Seattle, Washington; HUGHES—NOAA/Earth Sciences Research Laboratory/Physical Sciences Division, Boulder, Colorado; GUTMANN—National Center for Atmospheric Research, Boulder, Colorado; KAPNICK—NOAA/Geophysical Fluid Dynamics Laboratory, Princeton, New Jersey

CORRESPONDING AUTHOR: Jessica Lundquist, jdland@uw.edu

The abstract for this article can be found in this issue, following the table of contents.

DOI:10.1175/BAMS-D-19-0001.1

In final form 22 July 2019

©2019 American Meteorological Society

For information regarding reuse of this content and general copyright information, consult the [AMS Copyright Policy](#).

2008), provided a stable medium of both the best available input to a hydrologic model and the best available benchmark for an atmospheric model (Fig. 1). These gridded datasets are often referred to as observations, when in truth they are based on a statistical model interpolating between point measurements. To improve atmospheric model performance beyond the quality of a gridded precipitation field, it is necessary to explore other measurements, such as streamflow or snow on the ground. Similarly, for hydrologists to make better use of atmospheric model output, they need to understand how those models work, which requires greater knowledge of atmospheric science to understand what such models can and cannot be relied on for. Both steps require greater integration across disciplines to move forward.

REVIEW OF TRADITIONAL DISCIPLINARY KNOWLEDGE OF OROGRAPHIC PRECIPITATION. *Why we care.* Accurate precipitation estimates are essential for hydrologic predictions, ecological assessments, and infrastructure management. Hydrologists must translate atmospheric forecasts or point gauge measurements into precipitation amounts for the entire basin of interest. The importance of mountain precipitation to floods and summer water supplies has driven extensive studies and additional measurements in the fields of both hydrology and meteorology (Dettinger et al. 2004; Galewsky and Sobel 2005; Heggli and Rauber 1988; Jeton et al. 1996; Marwitz 1983; Pandey et al. 1999; Parish 1982; Ralph et al. 2005; Reeves et al. 2008; Smith et al. 2010). Despite our best efforts, many factors conspire to limit our ability to measure mountain

FIG. 1. Gridded precipitation products, such as PRISM (shown here for three river basins in California), have long served as the intermediary between hydrologic science and atmospheric science. As model output improves beyond the skill of our observational network and interpolation techniques, more complex and interdisciplinary evaluations are required.

precipitation, including blocked radar signals, spatial details too fine for satellites or sparse gauge networks, and precipitation gauges either capped by snow or missing snow blown over the top of the orifice. Direct measurements of precipitation, particularly snowfall, in complex terrain are frequently scarce relative to the spatial heterogeneity they are expected to measure and may be unreliable (Derin et al. 2016; Milly and Dunne 2002; Ohara et al. 2011; Rasmussen et al. 2012).

Hydrologist's approach to orographic precipitation.

Consistent and reliable estimates of precipitation are essential to hydrologic models (Clark and Slater 2006; Mizukami and Smith 2012). In situ data are considered the gold standard, more accurate and precise than either radar- or satellite-based estimates. The standard practice in mountain hydrologic modeling is to distribute precipitation across a basin by some interpolation scheme (Clark and Slater 2006; Daly et al. 1994; Hay and Clark 2003). In flat terrain, radar signals aid interpolation (Nelson et al. 2016), but in complex terrain, radar signals are generally blocked, and with the exception of a high-quality radar system in Switzerland (e.g., Panziera et al. 2018), radars frequently report less than 50% of observed precipitation (Ralph et al. 2014; Trappero et al. 2009; Westrick et al. 1999; Young et al. 1999; Zhang et al. 2012). Interpolation can be done either by allowing the number of stations and spatial patterns to change through time (Clark and Slater 2006), or based on a climatological pattern assumed to remain fixed proportional to a base station [e.g., Daly et al. (1994) and all datasets relying on PRISM, see Table 1 in Lundquist et al. (2015)].

Both interpolation methods have problems—the former due to hard-to-identify errors in high-elevation stations (e.g., Mizukami and Smith 2012) and poorly characterized patterns on the event time scale, and the latter due to spatial patterns of precipitation that differ from climatology on both storm-specific and even annual time scales (Lundquist et al. 2010, 2015). Efforts to avoid ubiquitous errors have used runoff observations and evapotranspiration estimations to “adjust” global precipitation datasets (Adam et al. 2006; Fekete et al. 2002; Milly and Dunne 2002; Xia 2008), with areas of limited gauge coverage and/or severe gauge undercatch, such as the arctic, noted as particularly problematic (Louie et al. 2002; Tian et al. 2007; Ye et al. 2012). While gridded datasets match precipitation observations well at locations that have been incorporated into the interpolation algorithms, most perform significantly less well at locations in between stations, where no training data were available (Currier et al. 2017; Gutmann et al.

2012; Hiemstra et al. 2006). Annual precipitation from many gridded datasets does not even exceed observed runoff in some mountain basins (Henn et al. 2018b); in their snow reanalysis approach Margulis et al. (2015) must adjust incoming solid precipitation to match high-elevation snow observations.

In addition to quantity, hydrologists often infer precipitation phase from an empirical temperature threshold guided by a gridded product of near-surface temperature and sometimes humidity (Harpold et al. 2017; Jennings et al. 2018). The exact parameters in these empirical algorithms are not transferable in space and/or time (Jennings et al. 2018) and are often ill suited to changing synoptic conditions, since conditions aloft also affect precipitation phase at the surface (Wayand et al. 2016). Additionally, the underlying temperature datasets may have significant problems (Feld et al. 2013; Minder et al. 2010; Wayand et al. 2016).

Atmospheric scientist's approach to orographic precipitation.

While most hydrologists view precipitation as an input parameter with some degree of uncertainty, atmospheric scientists view precipitation as the output of dynamic and physical processes (Fig. 2). Precipitation-generation mechanisms over terrain include the lifting of moist air as a result of synoptic-scale patterns (Fig. 2a), thermally driven convection, and orographic lifting when air impinges on topography (Fig. 2b), with multiple review summaries (Barros and Lettenmaier 1994; Houze 2012; Roe 2005; Smith 1979). Orographic precipitation is modified by the moisture flux, the slope of the terrain (Alpert 1986; Neiman et al. 2009, 2013; Roe and Baker 2006; Smith 1979), and degree of blocking (Hughes et al. 2009; Lundquist et al. 2010), as well as cloud microphysical processes (Fig. 2c) (Grubišić et al. 2005; Jankov et al. 2009; Roe and Baker 2006; Yang et al. 2012), which in turn affect the transition between hydrometeor types: snow, ice/graupel, and rain (Fig. 2d); (Minder and Kingsmill 2013; Minder et al. 2011). Upper-level wind speeds and directions control where precipitation reaches the surface, and complex wind patterns near the mountain surface determine the patterns of final snow distribution on the landscape (Greene et al. 1999; Mott and Lehning 2010; Winstral et al. 2002).

All of the above processes, with the exception of finescale snow deposition and redistribution (Liston et al. 1998; Vionnet et al. 2018), are incorporated into modern numerical weather prediction models, which, when using a grid spacing of 12 km or less, are able to resolve the topography that drives major orographic precipitation gradients (Anders et al. 2007; Barros

FIG. 2. (a) Synoptic-scale storms (showing a midlatitude cyclone here) bring moisture to the mountains, and (b) parcels of air may 1) be lifted over the mountains, 2) be blocked by the mountains, or 3) be convectively lifted along the mountains. (c) Microphysical processes determine fall speeds and horizontal transport and influence (d) the phase of precipitation falling at the surface.

and Lettenmaier 1994; Liu et al. 2011). These models provide information for both short-term weather forecasts (Benjamin et al. 2016a; Mass et al. 2002), and downscaled climate projections (Liu et al. 2017; Rasmussen et al. 2019; Salathé et al. 2008).

Mesoscale model precipitation estimates can vary substantially depending on model configuration: for example, boundary layer, convection, and microphysics schemes (Hughes et al. 2019; Jankov et al. 2009; Morales et al. 2018; Yang et al. 2012) and boundary conditions (Hughes et al. 2019). Verification and configuration of these models relies heavily on observations; however, point observations may not be comparable to the model grid scale, and small space-time offsets in the model can appear as large errors (Casati et al. 2008; Cassola et al. 2015), a problem that grows more pronounced with higher-resolution models (Cassola et al. 2015; Lack et al. 2010). Gridded precipitation products are sometimes used to mitigate these mismatches (Ikeda et al. 2010; Liu et al. 2011; Xu et al. 2018), but errors in the statistically interpolated grid can also make results misleading (Henn et al. 2018b; Prein and Gobiet 2017).

MODEL VERSUS OBSERVATIONAL PRECIPITATION SKILL. *Gauge-based assessments.*

Rain gauges undercatch actual rainfall in most environments (Collados-Lara et al. 2018; Liljedahl et al. 2017; Rodda 1968; Rodda and Dixon 2012; Sieck et al. 2007), and snowfall is even harder to measure (Rasmussen et al. 2012). Gridded precipitation products consistently match gauge-based precipitation better than atmospheric models, but given that the gridded products are interpolations between these

very observations, this is not an independent comparison. At independent observation sites, models have generally outperformed gridded estimates for annual precipitation totals by a factor of 2 (Currier et al. 2017; Gutmann et al. 2012), although very few studies have compared with observational stations not used in statistical training, and the best performer may vary between years (Wayand et al. 2013). Compared to snow accumulation measured at snow pillows across California, gridded datasets were unbiased on average, but underpredicted (by as much as 50%) events with a large proportion of postfrontal precipitation (Lundquist et al. 2015). Henn et al. (2018a) demonstrated large spread (typically $\pm 20\%$ in annual means) between gridded datasets across the western United States, and multiple European studies found spread across gridded datasets was as large as, if not larger than, spread between regional climate models, with the greatest spread in areas of low gauge density (Herold et al. 2017; Isotta et al. 2015; Prein and Gobiet 2017). Zhang and Anagnostou (2019) found that mesoscale model output worked better than gauge-based precipitation observations for bias-correcting and downscaling satellite-based precipitation products during convective events in mountains in Colombia, Peru, and Taiwan.

Assessments using a hydrological or land surface model.

Multiple studies have demonstrated that mesoscale model input may be comparable or preferable compared to gauge observations to drive a hydrologic and/or snow model in complex terrain. The majority of studies concluded that hydrologic model performance was similar between the two forcing datasets,

including areas in northwest Montana (Leung et al. 1996), the Pacific Northwest (Currier et al. 2017; Wayand et al. 2016), Colorado (Rasmussen et al. 2011), Northern California (Anderson et al. 2002), and Japan (Yoshitani et al. 2009). In the Pacific Northwest (Wayand et al. 2013; Westrick et al. 2002; Westrick and Mass 2001) and Iceland (Rögnvaldsson et al. 2004, 2007), mesoscale model forced simulations outperformed those using gauge data, which the authors attributed to unrepresentative gauge locations and gauge undercatch, respectively. A few studies found that gauge observations produced better streamflow results, but authors attributed this result to forecast errors because the mesoscale model was run in forecast mode (Anderson et al. 2002; Westrick et al. 2002) or to calibration bias because the hydrological model was calibrated to perform best when using gauge data (Kunstmann and Stadler 2005). A few studies demonstrated skillful mesoscale-model-forced snow simulations compared to snow observations, but did not directly compare with another simulation forced using gridded observations, including studies focused on Colorado (Ikeda et al. 2010; Rasmussen et al. 2011), the California Sierra Nevada (Wrzesien et al. 2015, 2017), and all of North America (Wrzesien et al. 2018). While all of the above results were also sensitive to the configuration and parameters of the selected models, they demonstrate that mesoscale model output is a viable option for mountain precipitation input in hydrologic applications.

To overcome the model dependence of the studies discussed above, some work has focused on a more direct way to extract precipitation from streamflow, while explicitly representing parameter uncertainties. The idea of doing “hydrology backwards” (Kirchner 2009) to infer precipitation from streamflow records has been formalized in a Bayesian framework (Kavetski et al. 2003; Kavetski et al. 2006a,b) and has been applied in a number of settings (Koskela et al. 2012; Kuczera et al. 2006; Renard et al. 2010; Thyer et al. 2009; Vrugt et al. 2008). These efforts attempt to account for uncertainties in other hydrologic fluxes, for example, evapotranspiration and groundwater gains and losses. In these examples, precipitation input uncertainty was accounted for by a set of precipitation multipliers used to relate gauge measured precipitation with basin average precipitation. These multipliers were inferred in conjunction with internal hydrologic model parameters, using Markov chain Monte Carlo (MCMC) iterations. Multiple studies (Henn et al. 2015, 2016, 2018b,c) employed these techniques and demonstrated that precipitation inferred from streamflow and snow

observations shows greater spatial and temporal variability than gridded datasets, including those gridded products that aim to explicitly represent uncertainty (Newman et al. 2015), and that modeled precipitation better matches these estimates than a range of gridded datasets in the California Sierra Nevada (Hughes et al. 2019). Collectively, this work demonstrated that precipitation uncertainty was the dominant source of model error (Kavetski et al. 2006a; Kuczera et al. 2006).

A PATH FORWARD: HOW TO FURTHER IMPROVE.

One of the most alarming conclusions from the cited works is that we cannot trust datasets that are often considered truth. In most cases, problems arise from treating a statistical gridding of limited observations as actual observations. While our direct observations are sometimes flawed, they are the closest we can get to truth. Thus, our strategy should not be to abandon observations in favor of modeling, but rather, to focus on 1) obtaining the best observations possible, including investing in quality controlling those observations so that erroneous measurements are excluded from networks, 2) including all types of related observations (Fig. 3), and 3) working across disciplines to clearly communicate what each observation represents well, its uncertainty, and under which conditions it struggles. These principals are not new; however, the reality of working out what constitutes an error remains an enormous challenge (Dee 2005; Diamond et al. 2013).

Quality measurements in the mountains. We must maintain funding for our core observational networks, and provide increased support for their maintenance and, where and when possible, their expansion. In citizen science efforts, such as CoCoRaHS (Cifelli et al. 2005) or Community Snow Obs (<http://communitysnowobs.org/>), more effort could be made to target people living in remote areas where gauge density is low, or to target areas beyond traditional roads, for example, ski huts, wilderness shelters, or fire lookouts.

However, incorrect observations are worse than no observations, and any effort to expand gauge networks must include adequate maintenance and quality control. River forecast centers regularly check for poor observations, but more effort could be invested into ways to automate such flagging of suspect observations due to their inconsistency with a model (e.g., see Hughes et al. 2012, their Fig. 9) or a range of model outputs, for example, testing of “hydrological coherence” (Laiti et al. 2018).

FIG. 3. Progress will require combining lots of measurements, with understanding their strengths and limitations, to constrain and evaluate models.

Ancillary measurements: Remote sensing, strategic radar, ecology, vapor flux, soil moisture, total storage. Significantly expanding direct observations of precipitation into remote complex terrain is likely not practical or feasible in the short term for many areas. However, models can be evaluated and improved with multiple other measurements that may be more robustly obtained.

Unlike most ground-based radar (Young et al. 1999), satellite views are not blocked by the mountains, and satellite flight frequency and resolution have been increasing in recent years (Entekhabi et al. 1999). The Tropical Rainfall Measurement Mission (TRMM) and the Global Precipitation Measurement (GPM) mission have provided estimates of precipitation over mountains where few in situ measurements exist, providing estimates of liquid precipitation (Houze et al. 2015), characteristics of convective precipitation (Rasmussen et al. 2013), and some options for hydrologic modeling (Xue et al. 2013). However, care must also be taken in the interpretation of satellite data, because these are modeled based on radiance retrieved at multiple wavelengths. To date, satellite algorithms have limited capability for atmospheric river events (Wen et al. 2018) or for estimating mixed or solid precipitation (Cao et al. 2018; Yong et al. 2012), particularly over complex terrain (Ebtehaj and Kummerow 2017; Kummerow et al. 2015). Globally, products that merge gauge data and satellite-based precipitation are often biased by unrepresentative and limited gauge data, particularly in mountains (Derin et al. 2016).

Progress has been made on better observing storm characteristics and vapor fluxes along mountain boundaries. For example, California has added upstream vertically pointing radars of various frequencies, including wind profilers and snow level radars (Ralph et al. 2014). Aircraft-based profile observations are also increasingly available, for example, aircraft meteorological data reports (AMDAR) and Tropospheric Aircraft Meteorological Data Reports (TAMDAR), (Moninger et al. 2003, 2010). These data sources could be assimilated to improve model predictions downwind or could be used to objectively select better performers from a suite of model options.

Although datasets that combine radar with data from other sources, such as the Multi-Radar Multi-Sensor (MRMS) product (Zhang et al. 2016), suffer from the same biases as their underlying datasets in complex terrain (Bytheway et al. 2019), gap-filling radar with a narrow beamwidth (e.g., C band or X band) has been used in the Bay Area of California (Cifelli et al. 2018; Willie et al. 2017), in Utah (Campbell and Steenburgh 2014), and in the Alps (Delrieu et al. 2009; Germann et al. 2006). The Swiss have developed operational methodology for using these observations in complex terrain, taking care to site the radars at high-elevation locations with clear views, to minimize sidelobe effects, and to correct for clutter (Germann et al. 2006). In Taiwan, vertical profile corrections were able to improve radar precipitation estimates where near-ground levels were blocked and only high-altitude signals were available

(Wang et al. 2016). While limited access and funding will still cause problems, other locations could learn from these examples and conduct studies to improve processing algorithms for gap filling radars in locations where they can be feasibly deployed. Another approach to improve the use of radar data is through fusion with weather models. The assimilation of both radar-measured 3D reflectivity and wind velocities with an atmospheric model is an area of active research (Benjamin et al. 2016b; Sun 2005; Wang et al. 2013). However, application in complex terrain requires significant development (Tai et al. 2017).

Another example of the use of nontraditional data sources in precipitation mapping includes the use of ecological information. Maps of ecological communities may not report storm-specific precipitation amounts, but in remote regions, they provide high-resolution spatially detailed patterns of precipitation climatologies. Giambelluca et al. (2013) used both vegetation maps and high-resolution atmospheric model output to guide underlying patterns for precipitation across the mountains of Hawaii, and these climatological maps have been used to create many gridded precipitation products in near-inaccessible and unmonitored locations (Newman et al. 2019).

Soil moisture observations can identify areas where rain occurred recently but either are very localized (point measurements) or have large footprints (~40 km for satellites) and only see the very top layer of the soil, for example, Soil Moisture and Ocean Salinity (SMOS; Srivastava et al. 2015) and Soil Moisture Active Passive (SMAP; Chan et al. 2018). The Gravity Recovery and Climate Experiment (GRACE; and its follow-on mission), can identify total changes in water content at large spatial scales (~200,000 km²), which can benchmark modeled total changes in frozen and subsurface water storage (Chen et al. 2017), but GRACE has a large footprint and cannot identify from which water storage component (snow, ice, or groundwater) a storage change originated from. Alternative soil moisture measurements based on cosmic-ray neutron measurements (Zreda et al. 2008) or GPS interferometry (Larson et al. 2008) both offer intermediate spatial scales (tens of meters), and a large network of GPS installations has existed for decades.

Snow measurements. Of all the ancillary land surface data available, snow observations come closest to directly informing precipitation accuracy, particularly in areas where precipitation is difficult to directly observe. Snow water equivalent (SWE) observations are part of many in situ operational networks and have been used to assess total annual frozen precipitation

as discussed above. Passive microwave observations of SWE from satellites are not robust for snow in the mountains, due to the large spatial footprint (~25 km), and the inability to sense deep (>20 cm) or wet snow (Dietz et al. 2012). Airborne gamma ray estimates of SWE are valid for shallow snow and flat terrain but are inaccurate in deep snow, forests, or complex terrain (Glynn et al. 1988). Thus, most spatially integrated assessments use other snow variables, such as snow depth and snow covered area, as described below.

SNOW DISAPPEARANCE DATE AND THE ENERGY BALANCE. For decades, hydrologists have compared distributed snow simulations with satellite-based estimates of snow-covered area (SCA) or snow cover extent (Shamir and Georgakakos 2006), which provides a time-integrated assessment of combined model performance for both snow accumulation and melt with about 500 m (MODIS) or 30 m (Landsat) spatial resolutions, with new satellites offering the potential for 3 m (Planet) daily observations. A number of studies have utilized the date of snow disappearance to determine how much snow must have fallen at a location in order for snow to disappear on an observed date given modeled melt rates, often termed SWE reconstruction (Bair et al. 2016; Cline et al. 1998; Livneh et al. 2014; Molotch and Margulis 2008; Raleigh and Lundquist 2012). Many of these utilize Bayesian methodology and changing precipitation weights (Margulis et al. 2015, 2016), and some have directly used the reconstructed product to look at snowfall patterns in space and time (Huning and Margulis 2017, 2018). Wrzesien et al. (2017) found that the Margulis et al. (2016) dataset compared favorably with WRF-based snow model simulations over the California Sierra Nevada. These techniques depend on accurately modeling the snow energy balance controlling melt rates. Thus, efforts to improve energy flux observations (e.g., snow albedo, snow surface temperature, radiation) will translate to improved retrospective snowfall estimation.

SNOW DEPTH. Snow depth observations can be obtained across large areas and at high resolutions (1–3-m spatial footprint) by repeat lidar (Painter et al. 2016) or satellite stereo photogrammetry techniques (Shean et al. 2016), but the direct application of these measurements to improving precipitation estimates has only just begun to be explored (Henn et al. 2016, 2018c; Mott et al. 2014). Snow depth is subject to localized effects such as preferential deposition, settling, and wind redistribution but can be aggregated to coarser resolutions for comparison with atmospheric

models (Mott and Lehning 2010; Mott et al. 2011). Most atmospheric and hydrologic models use snow water equivalent as a state variable, which is related to snow depth via bulk density. The density of new snowfall is highly uncertain (Wayand et al. 2017), making density estimates one of the larger sources of error in deriving SWE from lidar measurements (Raleigh and Small 2017). Snow compaction and evolution through time are better understood, resulting in more confidence in density estimates in the spring near peak snow accumulation, making seasonal assessments more accurate than storm-specific assessments. Impacts of snow evolution due to rain on snow (which may increase both water content and density of the snowpack but generally to an unknown extent) are also a source of uncertainty.

Combining disparate data sources through partnerships across disciplines. We propose that two interrelated barriers, time scales and analysis tools, have impeded the rate of crossing historical disciplinary boundaries. Vegetation maps reveal finescale spatial patterns but only long-timescale (e.g., multidecadal average) precipitation patterns. While streamflow gives a solid estimate of total rainfall over a catchment, it is at the cost of smoother spatial and temporal resolutions (Kretschmar et al. 2016). This smoothing in space and time makes streamflow and snow observations useful for evaluating mesoscale model output at an annual time scale, which fits well within a water resources perspective or a climate modeling development framework. Shorter time scales related to floods, particularly those due to convective precipitation events, are more difficult to assess using these techniques. Overall, both point observations and models struggle to accurately represent convective precipitation, and published research to date does not clearly indicate whether model simulations or gridded precipitation products are currently superior in this area. The spatial extent of convective precipitation can sometimes be identified more reliably with streamflow measurements than rain gauges in the mountains (Lundquist et al. 2009), but the precipitation magnitudes are hard to assess due to unknown water storage and evapotranspiration on the landscape over short time scales. Satellite data reveal convective patterns although amounts are biased (Rasmussen et al. 2013). Convection-resolving models (4 km or finer) currently look like a viable path forward, although reproducing specific convective storms remains a challenge (Rasmussen et al. 2019).

Analysis tools also contrast between the two disciplines. Stochastic parameter selection and other

statistical methods are common within hydrology and other land surface sciences, whereas deterministic methods dominate within atmospheric science—due in part to the computational costs involved. Hydrologists frequently run large model ensembles, ranging from one model with many different choices of parameter values, to a suite of models with different model choices in the equations used, often termed structural uncertainty (Clark et al. 2011). Comparisons of multiple long time series with observations are used to select choices that minimize errors. Numerical weather prediction modeling systems also often use ensembles (Hacker et al. 2011; Knierel et al. 2017), although ensembles using perturbed parameters (Greybush et al. 2017; Jankov et al. 2019) are rarer than those generated through perturbed initial conditions. There is invariably a trade-off between the size of the ensemble and the resolution and sophistication of the model (Ferro et al. 2012; Gowan et al. 2018). In general, testing uncertainty in parameter values or in both parameters and structure is more frequent in hydrology, while testing uncertainty in model structure alone is more frequent in atmospheric sciences.

These different approaches impact the necessary duration of measurements and of desired characteristics of mesoscale model output. Statistical methods become more reliable as the amount of training data (e.g., duration of time series) increases. Thus, many flood forecasters use gridded gauge data up until the time of the forecast to provide a long time period for model calibration and spinup, and an ensemble of weather model output for the future forecast (Pagano et al. 2014). Most research hydrologists prefer re-analysis datasets to provide meteorological forcings for their models because these models are run with a constant configuration over a long period of time. These allow hydrologists to calibrate unknown parameters in a hydrologic model (e.g., those related to subsurface storage and conductivity) and accommodate for biases in meteorological input (critical for flood forecasting or water resources operations).

In contrast, the increased value of physically based formulas is much less data volume dependent. Operational numerical weather prediction models change frequently to include the latest improvements in the modeling prediction system, which may be based on one short-term field campaign. However, this near-constant evolution makes them problematic for hydrologists since historic model calibration may introduce compensating biases that lead to the correct answer for the wrong reason (Kirchner 2006).

There are benefits to both stochastic and deterministic approaches, and both communities would benefit

from better sharing and integrating these techniques. For example, fully coupled land–atmosphere models currently lag behind the most promising hydrologic modeling results in mountainous terrain (Clark et al. 2015). This is likely the result of the coupled models being recently developed, often at coarser scales and with fewer processes represented, as well as the relative lack of model parameter estimation implemented in these studies. One path forward requires coupled models developed by actively engaged scientists from multiple disciplines and multiple communities because components that historically were unimportant to one or both disciplines will emerge as important in the coupled system. As atmospheric model output is more routinely adopted as hydrologic model input, as will happen with work currently underway with both the U.S. National Water Model (<http://water.noaa.gov/about/nwm>) and the European Flood Awareness System (www.efas.eu/about-efas.html), greater feedback and assessments will accumulate, which will improve both fields.

Strategically designing field campaigns requires a strong interdisciplinary approach in the earliest stages of planning. While atmospheric events pass within a week, the hydrologic cycle needs to be evaluated over the course of a year at minimum. Recent campaigns such as NOAA's Hydrometeorology Testbed (HMT) in California (Ralph et al. 2005), the Olympic Mountain Experiment (OLYMPEX) in Washington (Houze et al. 2017), and the Integrated Precipitation and Hydrology Experiment (IPHEX) in North Carolina (Barros et al. 2014; Tao et al. 2016) provide examples of how instrument deployments have been balanced across longer and shorter durations to attempt to meet both atmospheric and hydrologic objectives. Further integration across time scales could be achieved by

conducting shorter-duration atmospheric-focused campaigns at field locations that already have a long history of ground-based observations, for example, Long-Term Ecological Research (LTER; Kratz et al. 2003), Critical Zone Observatory (CZO; Lin and Hopmans 2011), or USDA research watersheds (Renard et al. 2008).

Despite efforts to encourage interdisciplinary work (Dirmeyer et al. 2015; National Research Council 1991), strong disciplinary boundaries still exist within universities and many funding agencies. The community should continue to advocate for interdisciplinary graduate education and for projects that combine knowledge and expertise across fields to advance these problems.

CONCLUSIONS. Our computational abilities have advanced more than our observational capabilities. Direct measurements of precipitation in complex terrain continue to be problematic. Even a concerted effort to invest financially in a network of mountain precipitation gauges may not improve our total gridded precipitation estimates significantly due to issues of access, property ownership, representativeness, and maintenance. Multiple types of measurements, including gauges, radars, satellites, streamflow observations, and snow observations (Fig. 3), must be brought to bear to assess our model output of mountain precipitation, and the combined modeling system must strive for excellence in all components relevant to those properties we can reliably measure (e.g., snow and streamflow). In the future, better coordination across disciplines could allow for improved coordination between the communities to improve available datasets available for hydrologic applications (Table 1).

TABLE 1. Summary advice.

Advice for atmospheric scientists

- Do not presume that your model's inability to match a gridded dataset means that it is wrong (Hughes et al. 2019; Liu et al. 2017)
- Investigate multiple sources of data, as well as many different types of storms, to make decisions between different model setups and parameterizations
- Consult hydrologists and snow scientists about the sources of uncertainty in streamflow, soil moisture, and snow measurements (useful, independent data)
- Consult those working in more application-oriented fields to understand what they really need (e.g., getting precipitation on the correct side of an orographic barrier is important)

Advice for land surface scientists

- High-resolution atmospheric model output is likely superior input to your hydrologic model than distributed gauge data
- Atmospheric models can output amounts of mixed-phase precipitation directly, often with better success than empirical methods based on surface observations of air temperature and humidity
- Not all simulations are created equal—befriend some modelers and ask questions about the parameterization schemes, resolution, nesting, etc.
- Work with meteorologists to better understand how atmospheric variables that influence hydrology may be discontinuously distributed in space and time (i.e., stationarity or linearity is often assumed, but may not hold in reality)

Realizing that our current modeling capabilities may now be surpassing our observational abilities in the realm of mountain precipitation speaks well to human ingenuity and scientific improvements over recent decades. We must take this realization as inspiration to now maintain and improve our observational networks and our capabilities in fully utilizing them in order to continue to improve our abilities to understand, model, and forecast our mountain water supplies, transportation hazards, and the elusive deep powder.

ACKNOWLEDGMENTS. We thank Lucas Harris, Chris Milly, Haonan Chen, Rob Cifelli, and two anonymous reviewers for their reviews of the manuscript. We also thank Catherine Raphael for her assistance in graphical design. The National Center for Atmospheric Research is sponsored by the National Science Foundation. J. Lundquist was partially supported by NASA Grant NNX17AL59G.

REFERENCES

- Adam, J. C., E. A. Clark, D. P. Lettenmaier, and E. F. Wood, 2006: Correction of global precipitation products for orographic effects. *J. Climate*, **19**, 15–38, <https://doi.org/10.1175/JCLI3604.1>.
- Alpert, P., 1986: Mesoscale indexing of the distribution of orographic precipitation over high mountains. *J. Climate Appl. Meteor.*, **25**, 532–545, [https://doi.org/10.1175/1520-0450\(1986\)025<0532:MIOTDO>2.0.CO;2](https://doi.org/10.1175/1520-0450(1986)025<0532:MIOTDO>2.0.CO;2).
- Anders, A. M., G. H. Roe, D. R. Durran, and J. R. Minder, 2007: Small-scale spatial gradients in climatological precipitation on the Olympic Peninsula. *J. Hydrometeor.*, **8**, 1068–1081, <https://doi.org/10.1175/JHM610.1>.
- Anderson, M. L., Z.-Q. Chen, M. L. Kavvas, and A. Feldman, 2002: Coupling HEC-HMS with atmospheric models for prediction of watershed runoff. *J. Hydrol. Eng.*, **7**, 312–318, [https://doi.org/10.1061/\(ASCE\)1084-0699\(2002\)7:4\(312\)](https://doi.org/10.1061/(ASCE)1084-0699(2002)7:4(312)).
- Bair, E. H., K. Rittger, R. E. Davis, T. H. Painter, and J. Dozier, 2016: Validating reconstruction of snow water equivalent in California's Sierra Nevada using measurements from the NASA Airborne Snow Observatory. *Water Resour. Res.*, **52**, 8437–8460, <https://doi.org/10.1002/2016WR018704>.
- Barros, A. P., and D. P. Lettenmaier, 1994: Dynamic modeling of orographically induced precipitation. *Rev. Geophys.*, **32**, 265–284, <https://doi.org/10.1029/94RG00625>.
- , and Coauthors, 2014: NASA GPM-Ground Validation Integrated Precipitation and Hydrology Experiment 2014 Science Plan. Duke University Tech. Rep., 64 pp., <https://doi.org/10.7924/G8CC0XMR>.
- Benjamin, S. G., J. M. Brown, and T. G. Smirnova, 2016a: Explicit precipitation-type diagnosis from a model using a mixed-phase bulk cloud–precipitation microphysics parameterization. *Wea. Forecasting*, **31**, 609–619, <https://doi.org/10.1175/WAF-D-15-0136.1>.
- , and Coauthors, 2016b: A North American hourly assimilation and model forecast cycle: The Rapid Refresh. *Mon. Wea. Rev.*, **144**, 1669–1694, <https://doi.org/10.1175/MWR-D-15-0242.1>.
- Bytheway, J. L., M. Hughes, K. Mahoney, and R. Cifelli, 2019: A multiscale evaluation of multisensor quantitative precipitation estimates in the Russian River Basin. *J. Hydrometeor.*, **20**, 447–466, <https://doi.org/10.1175/JHM-D-18-0142.1>.
- Campbell, L. S., and W. J. Steenburgh, 2014: Finescale orographic precipitation variability and gap-filling radar potential in Little Cottonwood Canyon, Utah. *Wea. Forecasting*, **29**, 912–935, <https://doi.org/10.1175/WAF-D-13-00129.1>.
- Cao, Q., T. H. Painter, W. R. Currier, J. D. Lundquist, and D. P. Lettenmaier, 2018: Estimation of precipitation over the OLYMPEX domain during winter 2015/16. *J. Hydrometeor.*, **19**, 143–160, <https://doi.org/10.1175/JHM-D-17-0076.1>.
- Casati, B., and Coauthors, 2008: Forecast verification: current status and future directions. *Meteor. Appl.*, **15**, 3–18, <https://doi.org/10.1002/met.52>.
- Cassola, F., F. Ferrari, and A. Mazzino, 2015: Numerical simulations of Mediterranean heavy precipitation events with the WRF model: A verification exercise using different approaches. *Atmos. Res.*, **164–165**, 210–225, <https://doi.org/10.1016/j.atmosres.2015.05.010>.
- Chan, S., and Coauthors, 2018: Development and assessment of the SMAP enhanced passive soil moisture product. *Remote Sens. Environ.*, **204**, 931–941, <https://doi.org/10.1016/j.rse.2017.08.025>.
- Chen, X., D. Long, Y. Hong, C. Zeng, and D. Yan, 2017: Improved modeling of snow and glacier melting by a progressive two-stage calibration strategy with GRACE and multisource data: How snow and glacier meltwater contributes to the runoff of the Upper Brahmaputra River basin? *Water Resour. Res.*, **53**, 2431–2466, <https://doi.org/10.1002/2016WR019656>.
- Cifelli, R., N. Doesken, P. Kennedy, L. D. Carey, S. A. Rutledge, C. Gimmestad, and T. Depue, 2005: The community collaborative rain, hail, and snow network: Informal education for scientists and citizens. *Bull. Amer. Meteor. Soc.*, **86**, 1069–1078, <https://doi.org/10.1175/BAMS-86-8-1069>.

- , V. Chandrasekar, H. Chen, and L. E. Johnson, 2018: High resolution radar quantitative precipitation estimation in the San Francisco Bay Area: Rainfall monitoring for the urban environment. *J. Meteor. Soc. Japan Ser. II*, **96A**, 141–155, <https://doi.org/10.2151/jmsj.2018-016>.
- Clark, M. P., and A. G. Slater, 2006: Probabilistic quantitative precipitation estimation in complex terrain. *J. Hydrometeor.*, **7**, 3–22, <https://doi.org/10.1175/JHM474.1>.
- , D. Kavetski, and F. Fenicia, 2011: Pursuing the method of multiple working hypotheses for hydrological modeling. *Water Resour. Res.*, **47**, W09301, <https://doi.org/10.1029/2010WR009827>.
- , and Coauthors, 2015: Improving the representation of hydrologic processes in Earth system models. *Water Resour. Res.*, **51**, 5929–5956, <https://doi.org/10.1002/2015WR017096>.
- Cline, D. W., R. C. Bales, and J. Dozier, 1998: Estimating the spatial distribution of snow in mountain basins using remote sensing and energy balance modeling. *Water Resour. Res.*, **34**, 1275–1285, <https://doi.org/10.1029/97WR03755>.
- Collados-Lara, A., E. Pardo-Iguzquiza, D. Pulido-Velazquez, and J. Jimenez-Sanchez, 2018: Precipitation fields in an alpine Mediterranean catchment: Inversion of precipitation gradient with elevation or undercatch of snowfall? *Int. J. Climatol.*, **38**, 3565–3578, <https://doi.org/10.1002/joc.5517>.
- Currier, W. R., T. Thorson, and J. D. Lundquist, 2017: Independent evaluation of frozen precipitation from WRF and PRISM in the Olympic Mountains. *J. Hydrometeor.*, **18**, 2681–2703, <https://doi.org/10.1175/JHM-D-17-0026.1>.
- Daly, C., R. P. Neilson, and D. L. Phillips, 1994: A statistical-topographic model for mapping climatological precipitation over mountainous terrain. *J. Appl. Meteor.*, **33**, 140–158, [https://doi.org/10.1175/1520-0450\(1994\)033<0140:ASTMFM>2.0.CO;2](https://doi.org/10.1175/1520-0450(1994)033<0140:ASTMFM>2.0.CO;2).
- , and Coauthors, 2008: Physiographically sensitive mapping of climatological temperature and precipitation across the conterminous United States. *Int. J. Climatol.*, **28**, 2031–2064, <https://doi.org/10.1002/joc.1688>.
- Dee, D. P., 2005: Bias and data assimilation. *Quart. J. Roy. Meteor. Soc.*, **131**, 3323–3343, <https://doi.org/10.1256/qj.05.137>.
- Delrieu, G., B. Boudevillain, J. Nicol, B. Chapon, P.-E. Kirstetter, H. Andrieu, and D. Faure, 2009: Bolène-2002 Experiment: Radar quantitative precipitation estimation in the Cévennes–Vivarais region, France. *J. Appl. Meteor. Climatol.*, **48**, 1422–1447, <https://doi.org/10.1175/2008JAMC1987.1>.
- Derin, Y., and Coauthors, 2016: Multiregional satellite precipitation products evaluation over complex terrain. *J. Hydrometeor.*, **17**, 1817–1836, <https://doi.org/10.1175/JHM-D-15-0197.1>.
- Dettinger, M., K. Redmond, and D. Cayan, 2004: Winter orographic precipitation ratios in the Sierra Nevada—Large-scale atmospheric circulations and hydrologic consequences. *J. Hydrometeor.*, **5**, 1102–1116, <https://doi.org/10.1175/JHM-390.1>.
- Diamond, H. J., and Coauthors, 2013: U.S. Climate Reference Network after one decade of operations: Status and assessment. *Bull. Amer. Meteor. Soc.*, **94**, 485–498, <https://doi.org/10.1175/BAMS-D-12-00170.1>.
- Dietz, A. J., C. Kuenzer, U. Gessner, and S. Dech, 2012: Remote sensing of snow—A review of available methods. *Int. J. Remote Sens.*, **33**, 4094–4134, <https://doi.org/10.1080/01431161.2011.640964>.
- Dirmeyer, P. A., and Coauthors, 2015: Advancing hydrometeorological-hydroclimatic-ecohydrological process understanding and predictions: Summary and findings from a community workshop at the Colorado School of Mines. Tech. Rep., 14 pp., www.nsf.gov/geo/ear/programs/nsf-hydro-atmos-workshop-whitepaper-dec2014.pdf.
- Ebtehaj, A., and C. Kummerow, 2017: Microwave retrievals of terrestrial precipitation over snow-covered surfaces: A lesson from the GPM satellite. *Geophys. Res. Lett.*, **44**, 6154–6162, <https://doi.org/10.1002/2017GL073451>.
- Entekhabi, D., and Coauthors, 1999: An agenda for land surface hydrology research and a call for the second international hydrological decade. *Bull. Amer. Meteor. Soc.*, **80**, 2043–2058, [https://doi.org/10.1175/1520-0477\(1999\)080<2043:AAFLSH>2.0.CO;2](https://doi.org/10.1175/1520-0477(1999)080<2043:AAFLSH>2.0.CO;2).
- Fekete, B. M., C. J. Vörösmarty, and W. Grabs, 2002: High-resolution fields of global runoff combining observed river discharge and simulated water balances. *Global Biogeochem. Cycles*, **16**, 1042, <https://doi.org/10.1029/1999GB001254>.
- Feld, S. I., N. C. Cristea, and J. D. Lundquist, 2013: Representing atmospheric moisture content along mountain slopes: Examination using distributed sensors in the Sierra Nevada, California. *Water Resour. Res.*, **49**, 4424–4441, <https://doi.org/10.1002/wrcr.20318>.
- Ferro, C. A., T. E. Jupp, F. H. Lambert, C. Huntingford, and P. M. Cox, 2012: Model complexity versus ensemble size: allocating resources for climate prediction. *Philos. Trans. Roy. Soc. London*, **370**, 1087–1099, <https://doi.org/10.1098/rsta.2011.0307>.
- Galewsky, J., and A. Sobel, 2005: Moist dynamics and orographic precipitation in northern and central California during the New Year's flood of 1997. *Mon. Wea. Rev.*, **133**, 1594–1612, <https://doi.org/10.1175/MWR2943.1>.

- Germann, U., G. Galli, M. Boscacci, and M. Bolliger, 2006: Radar precipitation measurement in a mountainous region. *Quart. J. Roy. Meteor. Soc.*, **132**, 1669–1692, <https://doi.org/10.1256/qj.05.190>.
- Giambelluca, T. W., Q. Chen, A. G. Frazier, J. P. Price, Y. Chen, P. Chu, J. K. Eischeid, and D.M. Delparte, 2013: Online rainfall atlas of Hawai‘i. *Bull. Amer. Meteor. Soc.*, **94**, 313–316, <https://doi.org/10.1175/BAMS-D-11-00228.1>.
- Glynn, J., T. Carroll, P. Holman, and R. Grasty, 1988: An airborne gamma ray snow survey of a forest covered area with a deep snowpack. *Remote Sens. Environ.*, **26**, 149–160, [https://doi.org/10.1016/0034-4257\(88\)90093-4](https://doi.org/10.1016/0034-4257(88)90093-4).
- Gowan, T. M., W. J. Steenburgh, and C. S. Schwartz, 2018: Validation of mountain precipitation forecasts from the convection-permitting NCAR ensemble and operational forecast systems over the western United States. *Wea. Forecasting*, **33**, 739–765, <https://doi.org/10.1175/WAF-D-17-0144.1>.
- Greene, E. M., G. E. Liston, and R. A. Pielke, 1999: Simulation of above treeline snowdrift formation using a numerical snow-transport model. *Cold Reg. Sci. Technol.*, **30**, 135–144, [https://doi.org/10.1016/S0165-232X\(99\)00008-7](https://doi.org/10.1016/S0165-232X(99)00008-7).
- Greybush, S. J., S. Saslo, and R. Grumm, 2017: Assessing the ensemble predictability of precipitation forecasts for the January 2015 and 2016 East Coast winter storms. *Wea. Forecasting*, **32**, 1057–1078, <https://doi.org/10.1175/WAF-D-16-0153.1>.
- Grubišić, V., R. Vellore, and A. Huggins, 2005: Quantitative precipitation forecasting of wintertime storms in the Sierra Nevada: Sensitivity to the microphysical parameterization and horizontal resolution. *Mon. Wea. Rev.*, **133**, 2834–2859, <https://doi.org/10.1175/MWR3004.1>.
- Gutmann, E. D., R. M. Rasmussen, C. Liu, K. Ikeda, D. J. Gochis, M. P. Clark, J. Dudhia, and G. Thompson, 2012: A comparison of statistical and dynamical downscaling of winter precipitation over complex terrain. *J. Climate*, **25**, 262–281, <https://doi.org/10.1175/2011JCLI4109.1>.
- Hacker, J. P., and Coauthors, 2011: The US Air Force Weather Agency’s mesoscale ensemble: scientific description and performance results. *Tellus*, **63A**, 625–641, <https://doi.org/10.1111/j.1600-0870.2010.00497.x>.
- Harpold, A., S. Rajagopal, J. Crews, T. Winchell, and R. Schumer, 2017: Relative humidity has uneven effects on shifts from snow to rain over the western U.S. *Geophys. Res. Lett.*, **44**, 9742–9750, <https://doi.org/10.1002/2017GL075046>.
- Hay, L. E., and M. Clark, 2003: Use of statistically and dynamically downscaled atmospheric model output for hydrologic simulations in three mountainous basins in the western United States. *J. Hydrol.*, **282**, 56–75, [https://doi.org/10.1016/S0022-1694\(03\)00252-X](https://doi.org/10.1016/S0022-1694(03)00252-X).
- Heggli, M. F., and R. M. Rauber, 1988: The characteristics and evolution of supercooled water in wintertime storms over the Sierra Nevada: A summary of microwave radiometric measurements taken during the Sierra Cooperative Pilot Project. *J. Appl. Meteor.*, **27**, 989–1015, [https://doi.org/10.1175/1520-0450\(1988\)027<0989:TCAEOS>2.0.CO;2](https://doi.org/10.1175/1520-0450(1988)027<0989:TCAEOS>2.0.CO;2).
- Henn, B., M. P. Clark, D. Kavetski, and J. D. Lundquist, 2015: Estimating mountain basin-mean precipitation from streamflow using Bayesian inference. *Water Resour. Res.*, **51**, 8012–8033, <https://doi.org/10.1002/2014WR016736>.
- , —, —, B. McGurk, T. H. Painter, and J. D. Lundquist, 2016: Combining snow, streamflow, and precipitation gauge observations to infer basin-mean precipitation. *Water Resour. Res.*, **52**, 8700–8723, <https://doi.org/10.1002/2015WR018564>.
- , A. J. Newman, B. Livneh, C. Daly, and J. D. Lundquist, 2018a: An assessment of differences in gridded precipitation datasets in complex terrain. *J. Hydrol.*, **556**, 1205–1219, <https://doi.org/10.1016/j.jhydrol.2017.03.008>.
- , M. P. Clark, D. Kavetski, A. J. Newman, M. Hughes, B. McGurk, and J. D. Lundquist, 2018b: Spatio-temporal patterns of precipitation inferred from streamflow observations across the Sierra Nevada mountain range. *J. Hydrol.*, **556**, 993–1012, <https://doi.org/10.1016/j.jhydrol.2016.08.009>.
- , and Coauthors, 2018c: High-elevation evapotranspiration estimates during drought: using streamflow and NASA airborne snow observatory SWE observations to close the Upper Tuolumne River basin water balance. *Water Resour. Res.*, **54**, 746–766, <https://doi.org/10.1002/2017WR020473>.
- Herold, N., A. Behrangi, and L. V. Alexander, 2017: Large uncertainties in observed daily precipitation extremes over land. *J. Geophys. Res. Atmos.*, **122**, 668–681, <https://doi.org/10.1002/2016JD025842>.
- Hiemstra, C. A., G. E. Liston, R. A. Pielke Sr., D. L. Birkenheuer, and S. C. Albers, 2006: Comparing Local Analysis and Prediction System (LAPS) assimilations with independent observations. *Wea. Forecasting*, **21**, 1024–1040, <https://doi.org/10.1175/WAF961.1>.
- Houze, R. A., Jr., 2012: Orographic effects on precipitating clouds. *Rev. Geophys.*, **50**, RG1001, <https://doi.org/10.1029/2011RG000365>.
- , K. L. Rasmussen, M. D. Zuluaga, and S. R. Brodzik, 2015: The variable nature of convection in the tropics and subtropics: A legacy of 16 years of

- the Tropical Rainfall Measuring Mission satellite. *Rev. Geophys.*, **53**, 994–1021, <https://doi.org/10.1002/2015RG000488>.
- , and Coauthors, 2017: The Olympic Mountains Experiment (OLYMPEX). *Bull. Amer. Meteor. Soc.*, **98**, 2167–2188, <https://doi.org/10.1175/BAMS-D-16-0182.1>.
- Hughes, M., A. Hall, and R. G. Fovell, 2009: Blocking in areas of complex topography, and its influence on rainfall distribution. *J. Atmos. Sci.*, **66**, 508–518, <https://doi.org/10.1175/2008JAS2689.1>.
- , P. J. Neiman, E. Sukovich, and M. Ralph, 2012: Representation of the Sierra Barrier Jet in 11 years of a high-resolution dynamical reanalysis downscaling compared with long-term wind profiler observations. *J. Geophys. Res.*, **117**, D18116, <https://doi.org/10.1029/2012JD017869>.
- , J. D. Lundquist, and B. Henn, 2019: Dynamical downscaling improves upon gridded precipitation products in the Sierra Nevada, California. *Climate Dyn.*, <https://doi.org/10.1007/s00382-017-3631-z>, in press.
- Huning, L. S., and S. A. Margulis, 2017: Climatology of seasonal snowfall accumulation across the Sierra Nevada (USA): Accumulation rates, distributions, and variability. *Water Resour. Res.*, **53**, 6033–6049, <https://doi.org/10.1002/2017WR020915>.
- , and —, 2018: Investigating the variability of high-elevation seasonal orographic snowfall enhancement and its drivers across Sierra Nevada, California. *J. Hydrometeor.*, **19**, 47–67, <https://doi.org/10.1175/JHM-D-16-0254.1>.
- Ikeda, K., and Coauthors, 2010: Simulation of seasonal snowfall over Colorado. *Atmos. Res.*, **97**, 462–477, <https://doi.org/10.1016/j.atmosres.2010.04.010>.
- Isotta, F. A., R. Vogel, and C. Frei, 2015: Evaluation of European regional reanalyses and downscalings for precipitation in the Alpine region. *Meteor. Z.*, **24**, 15–37, <https://doi.org/10.1127/metz/2014/0584>.
- Jankov, I., J.-W. Bao, P. J. Neiman, P. J. Schultz, H. Yuan, and A. B. White, 2009: Evaluation and comparison of microphysical algorithms in ARW-WRF model simulations of atmospheric river events affecting the California coast. *J. Hydrometeor.*, **10**, 847–870, <https://doi.org/10.1175/2009JHM1059.1>.
- , J. Beck, J. Wolff, M. Harrold, J. B. Olson, T. Smirnova, C. Alexander, and J. Berner, 2019: Stochastically perturbed parameterizations in an HRRR-based ensemble. *Mon. Wea. Rev.*, **147**, 153–173, <https://doi.org/10.1175/MWR-D-18-0092.1>.
- Jennings, K. S., T. S. Winchell, B. Livneh, and N. P. Molotch, 2018: Spatial variation of the rain–snow temperature threshold across the Northern Hemisphere. *Nat. Commun.*, **9**, 1148, <https://doi.org/10.1038/s41467-018-03629-7>.
- Jeton, A. E., M. D. Dettinger, and J. L. Smith, 1996: Potential effects of climate change on streamflow, eastern and western slopes of the Sierra Nevada, California and Nevada. USGS Water-Resources Investigations Rep. 95-4260, 44 pp., <https://doi.org/10.3133/wri954260>.
- Kavetski, D., S. W. Franks, and G. Kuczera, 2003: Confronting input uncertainty in environmental modeling. *Calibration of Watershed Models*, Q. Duan et al., Eds., Water Science and Application, Vol. 6, Amer. Geophys. Union, 49–68.
- , G. Kuczera, and S. W. Franks, 2006a: Bayesian analysis of input uncertainty in hydrological modeling: 1. Theory. *Water Resour. Res.*, **42**, W03407, <https://doi.org/10.1029/2005WR004368>.
- , —, and —, 2006b: Bayesian analysis of input uncertainty in hydrological modeling: 2. Application. *Water Resour. Res.*, **42**, W03408, <https://doi.org/10.1029/2005WR004376>.
- Kirchner, J. W., 2006: Getting the right answers for the right reasons: Linking measurements, analyses, and models to advance the science of hydrology. *Water Resour. Res.*, **42**, W03S04, <https://doi.org/10.1029/2005WR004362>.
- , 2009: Catchments as simple dynamical systems: Catchment characterization, rainfall-runoff modeling, and doing hydrology backward. *Water Resour. Res.*, **45**, W02429, <https://doi.org/10.1029/2008WR006912>.
- Knievel, J. C., and Coauthors, 2017: Mesoscale ensemble weather prediction at U.S. Army Dugway Proving Ground, Utah. *Wea. Forecasting*, **32**, 2195–2216, <https://doi.org/10.1175/WAF-D-17-0049.1>.
- Koskela, J. J., B. Croke, H. Koivusalo, A. Jakeman, and T. Kokkonen, 2012: Bayesian inference of uncertainties in precipitation-streamflow modeling in a snow affected catchment. *Water Resour. Res.*, **48**, W11513, <https://doi.org/10.1029/2011WR011773>.
- Kratz, T. K., L. A. Deegan, M. E. Harmon, and W. K. Lauenroth, 2003: Ecological variability in space and time: Insights gained from the US LTER program. *BioScience*, **53**, 57–67, [https://doi.org/10.1641/0006-3568\(2003\)053\[0057:EVISAT\]2.0.CO;2](https://doi.org/10.1641/0006-3568(2003)053[0057:EVISAT]2.0.CO;2).
- Kretzschmar, A., W. Tych, N. Chappell, and K. Beven, 2016: What really happens at the end of the rainbow? Paying the price for reducing uncertainty (using reverse hydrology models). *Procedia Eng.*, **154**, 1333–1340, <https://doi.org/10.1016/j.proeng.2016.07.485>.
- Kuczera, G., D. Kavetski, S. Franks, and M. Thyer, 2006: Towards a Bayesian total error analysis of conceptual rainfall-runoff models: Characterising

- model error using storm-dependent parameters. *J. Hydrol.*, **331**, 161–177, <https://doi.org/10.1016/j.jhydrol.2006.05.010>.
- Kummerow, C. D., D. L. Randel, M. Kulie, N.-Y. Wang, R. Ferraro, S. J. Munchak, and V. Petkovic, 2015: The evolution of the Goddard profiling algorithm to a fully parametric scheme. *J. Atmos. Oceanic Technol.*, **32**, 2265–2280, <https://doi.org/10.1175/JTECH-D-15-0039.1>.
- Kunstmann, H., and C. Stadler, 2005: High resolution distributed atmospheric-hydrological modelling for Alpine catchments. *J. Hydrol.*, **314**, 105–124, <https://doi.org/10.1016/j.jhydrol.2005.03.033>.
- Lack, S., G. Limpert, and N. Fox, 2010: An object-oriented multiscale verification scheme. *Wea. Forecasting*, **25**, 79–92, <https://doi.org/10.1175/2009WAF2222245.1>.
- Laiti, L., and Coauthors, 2018: Testing the hydrological coherence of high-resolution gridded precipitation and temperature data sets. *Water Resour. Res.*, **54**, 1999–2016, <https://doi.org/10.1002/2017WR021633>.
- Larson, K. M., E. E. Small, E. D. Gutmann, A. L. Bilich, J. J. Braun, and V. U. Zavorotny, 2008: Use of GPS receivers as a soil moisture network for water cycle studies. *Geophys. Res. Lett.*, **35**, L24405, <https://doi.org/10.1029/2008GL036013>.
- Leung, L. R., M. S. Wigmosta, S. J. Ghan, D. J. Epstein, and L. W. Vail, 1996: Application of a subgrid orographic precipitation/surface hydrology scheme to a mountain watershed. *J. Geophys. Res.*, **101**, 12 803–12 817, <https://doi.org/10.1029/96JD00441>.
- Liljedahl, A. K., L. D. Hinzman, D. L. Kane, W. C. Oechel, C. E. Tweedie, and D. Zona, 2017: Tundra water budget and implications of precipitation underestimation. *Water Resour. Res.*, **53**, 6472–6486, <https://doi.org/10.1002/2016WR020001>.
- Lin, H., and J. W. Hopmans, 2011: Interdisciplinary sciences in a global network of critical zone observatories. *Vadose Zone J.*, **10**, 781–785, <https://doi.org/10.2136/vzj2011.0084>.
- Liston, G. E., G. Liston, and M. Sturm, 1998: A snow-transport model for complex terrain. *J. Glaciol.*, **44**, 498–516, <https://doi.org/10.1017/S002214300002021>.
- Liu, C., K. Ikeda, G. Thompson, R. Rasmussen, and J. Dudhia, 2011: High-resolution simulations of wintertime precipitation in the Colorado headwaters region: Sensitivity to physics parameterizations. *Mon. Wea. Rev.*, **139**, 3533–3553, <https://doi.org/10.1175/MWR-D-11-00009.1>.
- , and Coauthors, 2017: Continental-scale convection-permitting modeling of the current and future climate of North America. *Climate Dyn.*, **49**, 71–95, <https://doi.org/10.1007/s00382-016-3327-9>.
- Livneh, B., J. S. Deems, D. Schneider, J. J. Barsugli, and N. P. Molotch, 2014: Filling in the gaps: Inferring spatially distributed precipitation from gauge observations over complex terrain. *Water Resour. Res.*, **50**, 8589–8610, <https://doi.org/10.1002/2014WR015442>.
- Louie, P. Y. T., W. D. Hogg, M. D. MacKay, X. Zhang, and R. F. Hopkinson, 2002: The water balance climatology of the Mackenzie basin with reference to the 1994/95 water year. *Atmos.–Ocean*, **40**, 159–180, <https://doi.org/10.3137/ao.400206>.
- Lundquist, J., B. Huggett, H. Roop, and N. Low, 2009: Use of spatially distributed stream stage recorders to augment rain gages by identifying locations of thunderstorm precipitation and distinguishing rain from snow. *Water Resour. Res.*, **45**, W00D25, <https://doi.org/10.1029/2008WR006995>.
- , J. R. Minder, P. J. Neiman, and E. Sukovich, 2010: Relationships between barrier jet heights, orographic precipitation gradients, and streamflow in the northern Sierra Nevada. *J. Hydrometeorol.*, **11**, 1141–1156, <https://doi.org/10.1175/2010JHM1264.1>.
- , M. Hughes, B. Henn, E. D. Gutmann, B. Livneh, J. Dozier, and P. Neiman, 2015: High-elevation precipitation patterns: Using snow measurements to assess daily gridded datasets across the Sierra Nevada, California. *J. Hydrometeorol.*, **16**, 1773–1792, <https://doi.org/10.1175/JHM-D-15-0019.1>.
- Margulis, S. A., M. Girotto, G. Cortés, and M. Durand, 2015: A particle batch smoother approach to snow water equivalent estimation. *J. Hydrometeorol.*, **16**, 1752–1772, <https://doi.org/10.1175/JHM-D-14-0177.1>.
- , G. Cortés, M. Girotto, and M. Durand, 2016: A Landsat-era Sierra Nevada snow reanalysis (1985–2015). *J. Hydrometeorol.*, **17**, 1203–1221, <https://doi.org/10.1175/JHM-D-15-0177.1>.
- Marwitz, J. D., 1983: The kinematics of orographic airflow during Sierra storms. *J. Atmos. Sci.*, **40**, 1218–1227, [https://doi.org/10.1175/1520-0469\(1983\)040<1218:TKOAOAD>2.0.CO;2](https://doi.org/10.1175/1520-0469(1983)040<1218:TKOAOAD>2.0.CO;2).
- Mass, C., D. Ovens, K. Westrick, and B. Colle, 2002: Does increasing horizontal resolution produce more skillful forecasts? The results of two years of real-time numerical weather prediction over the Pacific Northwest. *Bull. Amer. Meteor. Soc.*, **83**, 407–430, [https://doi.org/10.1175/1520-0477\(2002\)083<0407:DIHRPM>2.3.CO;2](https://doi.org/10.1175/1520-0477(2002)083<0407:DIHRPM>2.3.CO;2).
- Milly, P., and K. Dunne, 2002: Macroscale water fluxes 1. Quantifying errors in the estimation of basin mean precipitation. *Water Resour. Res.*, **38**, 1205, <https://doi.org/10.1029/2001WR000759>.
- Minder, J. R., and D. E. Kingsmill, 2013: Mesoscale variations of the atmospheric snow line over the northern Sierra Nevada: Multiyear statistics, case

- study, and mechanisms. *J. Atmos. Sci.*, **70**, 916–938, <https://doi.org/10.1175/JAS-D-12-0194.1>.
- , P. W. Mote, and J. D. Lundquist, 2010: Surface temperature lapse rates over complex terrain: Lessons from the Cascade Mountains. *J. Geophys. Res.*, **115**, D14122, <https://doi.org/10.1029/2009JD013493>.
- , D. R. Durran, and G. H. Roe, 2011: Mesoscale controls on the mountainside snow line. *J. Atmos. Sci.*, **68**, 2107–2127, <https://doi.org/10.1175/JAS-D-10-05006.1>.
- Mizukami, N., and M. B. Smith, 2012: Analysis of inconsistencies in multi-year gridded quantitative precipitation estimate over complex terrain and its impact on hydrologic modeling. *J. Hydrol.*, **428–429**, 129–141, <https://doi.org/10.1016/j.jhydrol.2012.01.030>.
- Molotch, N. P., and S. A. Margulis, 2008: Estimating the distribution of snow water equivalent using remotely sensed snow cover data and a spatially distributed snowmelt model: A multi-resolution, multi-sensor comparison. *Adv. Water Resour.*, **31**, 1503–1514, <https://doi.org/10.1016/j.advwatres.2008.07.017>.
- Moninger, W. R., R. D. Mamrosh, and P. M. Pauley, 2003: Automated meteorological reports from commercial aircraft. *Bull. Amer. Meteor. Soc.*, **84**, 203–216, <https://doi.org/10.1175/BAMS-84-2-203>.
- , S. G. Benjamin, B. D. Jamison, T. W. Schlatter, T. L. Smith, and E. J. Szoke, 2010: Evaluation of regional aircraft observations using TAMDAR. *Wea. Forecasting*, **25**, 627–645, <https://doi.org/10.1175/2009WAF2222321.1>.
- Morales, A., H. Morrison, and D. Posselt, 2018: Orographic precipitation response to microphysical parameter perturbations for idealized moist nearly neutral flow. *J. Atmos. Sci.*, **75**, 1933–1953, <https://doi.org/10.1175/JAS-D-17-0389.1>.
- Mott, R., and M. Lehning, 2010: Meteorological modeling of very high-resolution wind fields and snow deposition for mountains. *J. Hydrometeor.*, **11**, 934–949, <https://doi.org/10.1175/2010JHM1216.1>.
- , M. Schirmer, and M. Lehning, 2011: Scaling properties of wind and snow depth distribution in an Alpine catchment. *J. Geophys. Res.*, **116**, D06106, <https://doi.org/10.1029/2010JD014886>.
- , D. Scipion, M. Schneebeli, N. Dawes, A. Berne, and M. Lehning, 2014: Orographic effects on snow deposition patterns in mountainous terrain. *J. Geophys. Res. Atmos.*, **119**, 1363–1385, <https://doi.org/10.1002/2013JD019880>.
- National Research Council, 1991: *Opportunities in the Hydrologic Sciences*. National Academies Press, 371 pp., <https://doi.org/10.17226/1543>.
- Neiman, P. J., A. B. White, F. M. Ralph, D. J. Gottas, and S. I. Gutman, 2009: A water vapour flux tool for precipitation forecasting. *Proc. Inst. Civ. Eng.: Water Manage.*, **162**, 83–94, <https://doi.org/10.1680/wama.2009.162.2.83>.
- , M. Hughes, B. J. Moore, F. M. Ralph, and E. M. Sukovich, 2013: Sierra barrier jets, atmospheric rivers, and precipitation characteristics in northern California: A composite perspective based on a network of wind profilers. *Mon. Wea. Rev.*, **141**, 4211–4233, <https://doi.org/10.1175/MWR-D-13-00112.1>.
- Nelson, B. R., O. P. Prat, D.-J. Seo, and E. Habib, 2016: Assessment and implications of NCEP Stage IV quantitative precipitation estimates for product inter-comparisons. *Wea. Forecasting*, **31**, 371–394, <https://doi.org/10.1175/WAF-D-14-00112.1>.
- Newman, A. J., and Coauthors, 2015: Gridded ensemble precipitation and temperature estimates for the contiguous United States. *J. Hydrometeor.*, **16**, 2481–2500, <https://doi.org/10.1175/JHM-D-15-0026.1>.
- , M. P. Clark, R. J. Longman, and T. W. Giambelluca, 2019: Methodological intercomparisons of station-based gridded meteorological products: Utility, limitations, and paths forward. *J. Hydrometeor.*, **20**, 531–547, <https://doi.org/10.1175/JHM-D-18-0114.1>.
- Ohara, N., M. Kavvas, S. Kure, Z. Chen, S. Jang, and E. Tan, 2011: Physically based estimation of maximum precipitation over American River watershed, California. *J. Hydrol. Eng.*, **16**, 351–361, [https://doi.org/10.1061/\(ASCE\)HE.1943-5584.0000324](https://doi.org/10.1061/(ASCE)HE.1943-5584.0000324).
- Pagano, T. C., and Coauthors, 2014: Challenges of operational river forecasting. *J. Hydrometeor.*, **15**, 1692–1707, <https://doi.org/10.1175/JHM-D-13-0188.1>.
- Painter, T. H., and Coauthors, 2016: The Airborne Snow Observatory: Fusion of scanning lidar, imaging spectrometer, and physically-based modeling for mapping snow water equivalent and snow albedo. *Remote Sens. Environ.*, **184**, 139–152, <https://doi.org/10.1016/j.rse.2016.06.018>.
- Pandey, G. R., D. R. Cayan, and K. P. Georgakakos, 1999: Precipitation structure in the Sierra Nevada of California during winter. *J. Geophys. Res.*, **104**, 12019–12030, <https://doi.org/10.1029/1999JD900103>.
- Panziera, L., M. Gabella, U. Germann, and O. Martius, 2018: A 12-year radar-based climatology of daily and sub-daily extreme precipitation over the Swiss Alps. *Int. J. Climatol.*, **38**, 3749–3769, <https://doi.org/10.1002/joc.5528>.
- Parish, T. R., 1982: Barrier winds along the Sierra Nevada mountains. *J. Appl. Meteor.*, **21**, 925–930, [https://doi.org/10.1175/1520-0450\(1982\)021<0925:BWATSN>2.0.CO;2](https://doi.org/10.1175/1520-0450(1982)021<0925:BWATSN>2.0.CO;2).
- Prein, A. F., and A. Gobiet, 2017: Impacts of uncertainties in European gridded precipitation observations

- on regional climate analysis. *Int. J. Climatol.*, **37**, 305–327, <https://doi.org/10.1002/joc.4706>.
- Raleigh, M. S., and J. D. Lundquist, 2012: Comparing and combining SWE estimates from the SNOW-17 model using PRISM and SWE reconstruction. *Water Resour. Res.*, **48**, W01506, <https://doi.org/10.1029/2011WR010542>.
- Raleigh, M. S., and E. E. Small, 2017: Snowpack density modeling is the primary source of uncertainty when mapping basin-wide SWE with lidar. *Geophys. Res. Lett.*, **44**, 3700–3709, <https://doi.org/10.1002/2016GL071999>.
- Ralph, F. M., and Coauthors, 2005: Improving short-term (0–48 h) cool-season quantitative precipitation forecasting: Recommendations from a USWRP workshop. *Bull. Amer. Meteor. Soc.*, **86**, 1619–1632, <https://doi.org/10.1175/BAMS-86-11-1615>.
- , and Coauthors, 2014: A vision for future observations for western US extreme precipitation and flooding. *J. Contemp. Water Res. Educ.*, **153**, 16–32, <https://doi.org/10.1111/j.1936-704X.2014.03176.x>.
- Rasmussen, K. L., S. L. Choi, M. D. Zuluaga, and R. A. Houze Jr., 2013: TRMM precipitation bias in extreme storms in South America. *Geophys. Res. Lett.*, **40**, 3457–3461, <https://doi.org/10.1002/grl.50651>.
- , A. F. Prein, R. M. Rasmussen, K. Ikeda, and C. Liu, 2019: Changes in the convective population and thermodynamic environments in convection-permitting regional climate simulations over the United States. *Climate Dyn.*, <https://doi.org/10.1007/s00382-017-4000-7>, in press.
- Rasmussen, R., and Coauthors, 2011: High-resolution coupled climate runoff simulations of seasonal snowfall over Colorado: A process study of current and warmer climate. *J. Climate*, **24**, 3015–3048, <https://doi.org/10.1175/2010JCLI3985.1>.
- , and Coauthors, 2012: How well are we measuring snow: The NOAA/FAA/NCAR winter precipitation test bed. *Bull. Amer. Meteor. Soc.*, **93**, 811–829, <https://doi.org/10.1175/BAMS-D-11-00052.1>.
- Reeves, H. D., Y.-L. Lin, and R. Rotunno, 2008: Dynamic forcing and mesoscale variability of heavy precipitation events over the Sierra Nevada Mountains. *Mon. Wea. Rev.*, **136**, 62–77, <https://doi.org/10.1175/2007MWR2164.1>.
- Renard, B., D. Kavetski, G. Kuczera, M. Thyer, and S. W. Franks, 2010: Understanding predictive uncertainty in hydrologic modeling: The challenge of identifying input and structural errors. *Water Resour. Res.*, **46**, W05521, <https://doi.org/10.1029/2009WR008328>.
- Renard, K. G., M. Nichols, D. Woolhiser, and H. Osborn, 2008: A brief background on the U.S. Department of Agriculture Agricultural Research Service Walnut Gulch Experimental Watershed. *Water Resour. Res.*, **44**, W05S02, <https://doi.org/10.1029/2006WR005691>.
- Rodda, J., 1968: The rainfall measurement problem. *IAHS Publ.*, **78**, 215–231.
- , and H. Dixon, 2012: Rainfall measurement revisited. *Weather*, **67**, 131–136, <https://doi.org/10.1002/wea.875>.
- Roe, G. H., 2005: Orographic precipitation. *Annu. Rev. Earth Planet. Sci.*, **33**, 645–671, <https://doi.org/10.1146/annurev.earth.33.092203.122541>.
- , and M. B. Baker, 2006: Microphysical and geometrical controls on the pattern of orographic precipitation. *J. Atmos. Sci.*, **63**, 861–880, <https://doi.org/10.1175/JAS3619.1>.
- Rögnvaldsson, Ó., P. Crochet, and H. Ólafsson, 2004: Mapping of precipitation in Iceland using numerical simulations and statistical modeling. *Meteor. Z.*, **13**, 209–219, <https://doi.org/10.1127/0941-2948/2004/0013-0209>.
- , J. F. Jónsdóttir, and H. Ólafsson, 2007: Numerical simulations of precipitation in the complex terrain of Iceland—Comparison with glaciological and hydrological data. *Meteor. Z.*, **16**, 71–85, <https://doi.org/10.1127/0941-2948/2007/0174>.
- Salathé, E. P., R. Steed, C. F. Mass, and P. H. Zahn, 2008: A high-resolution climate model for the U.S. Pacific Northwest mesoscale feedbacks and local responses to climate change. *J. Climate*, **21**, 5708–5726, <https://doi.org/10.1175/2008JCLI2090.1>.
- Shamir, E., and K. P. Georgakakos, 2006: Distributed snow accumulation and ablation modeling in the American River basin. *Adv. Water Resour.*, **29**, 558–570, <https://doi.org/10.1016/j.advwatres.2005.06.010>.
- Shean, D. E., O. Alexandrov, Z. M. Moratto, B. E. Smith, I. R. Joughin, C. Porter, and P. Morin, 2016: An automated, open-source pipeline for mass production of digital elevation models (DEMs) from very-high-resolution commercial stereo satellite imagery. *ISPRS J. Photogramm. Remote Sens.*, **116**, 101–117, <https://doi.org/10.1016/j.isprsjprs.2016.03.012>.
- Sieck, L. C., S. J. Burges, and M. Steiner, 2007: Challenges in obtaining reliable measurements of point rainfall. *Water Resour. Res.*, **43**, W01420, <https://doi.org/10.1029/2005WR004519>.
- Smith, J. A., M. L. Baeck, G. Villarini, and W. F. Krajewski, 2010: The hydrology and hydrometeorology of flooding in the Delaware River Basin. *J. Hydrometeorol.*, **11**, 841–859, <https://doi.org/10.1175/2010JHM1236.1>.
- Smith, R. B., 1979: The influence of mountains on the atmosphere. *Advances in Geophysics*, Vol. 21, Academic Press, 87–230, [https://doi.org/10.1016/S0065-2687\(08\)60262-9](https://doi.org/10.1016/S0065-2687(08)60262-9).
- Srivastava, P. K., D. Han, M. A. Rico-Ramirez, P. O'Neill, T. Islam, M. Gupta, and Q. Dai, 2015: Performance

- evaluation of WRF-Noah Land surface model estimated soil moisture for hydrological application: Synergistic evaluation using SMOS retrieved soil moisture. *J. Hydrol.*, **529**, 200–212, <https://doi.org/10.1016/j.jhydrol.2015.07.041>.
- Sun, J., 2005: Convective-scale assimilation of radar data: Progress and challenges. *Quart. J. Roy. Meteor. Soc.*, **131**, 3439–3463, <https://doi.org/10.1256/qj.05.149>.
- Tai, S.-L., Y.-C. Liou, J. Sun, and S.-F. Chang, 2017: The development of a terrain-resolving scheme for the forward model and its adjoint in the Four-Dimensional Variational Doppler Radar Analysis System (VDRAS). *Mon. Wea. Rev.*, **145**, 289–306, <https://doi.org/10.1175/MWR-D-16-0092.1>.
- Tao, J., D. Wu, J. Gourley, S. Q. Zhang, W. Crow, C. Peters-Lidard, and A. P. Barros, 2016: Operational hydrological forecasting during the IPHEX-IOP campaign—Meet the challenge. *J. Hydrol.*, **541**, 434–456, <https://doi.org/10.1016/j.jhydrol.2016.02.019>.
- Thyer, M., B. Renard, D. Kavetski, G. Kuczera, S. W. Franks, and S. Srikanthan, 2009: Critical evaluation of parameter consistency and predictive uncertainty in hydrological modeling: A case study using Bayesian total error analysis. *Water Resour. Res.*, **45**, W00B14, <https://doi.org/10.1029/2008WR006825>.
- Tian, Y., C. D. Peters-Lidard, B. J. Choudhury, and M. Garcia, 2007: Multitemporal analysis of TRMM-based satellite precipitation products for land data assimilation applications. *J. Hydrometeorol.*, **8**, 1165–1183, <https://doi.org/10.1175/2007JHM859.1>.
- Trapero, L., J. Bech, T. Rigo, N. Pineda, and D. Forcadell, 2009: Uncertainty of precipitation estimates in convective events by the Meteorological Service of Catalonia radar network. *Atmos. Res.*, **93**, 408–418, <https://doi.org/10.1016/j.atmosres.2009.01.021>.
- Vionnet, V., G. Guyomarc'h, M. Lafaysse, F. Naaim-Bouvet, G. Giraud, and Y. Deliot, 2018: Operational implementation and evaluation of a blowing snow scheme for avalanche hazard forecasting. *Cold Reg. Sci. Technol.*, **147**, 1–10, <https://doi.org/10.1016/j.coldregions.2017.12.006>.
- Vrugt, J. A., C. G. Diks, and M. P. Clark, 2008: Ensemble Bayesian model averaging using Markov chain Monte Carlo sampling. *Environ. Fluid Mech.*, **8**, 579–595, <https://doi.org/10.1007/s10652-008-9106-3>.
- Wang, H., J. Sun, X. Zhang, X.-Y. Huang, and T. Auligné, 2013: Radar data assimilation with WRF 4D-Var. Part I: System development and preliminary testing. *Mon. Wea. Rev.*, **141**, 2224–2244, <https://doi.org/10.1175/MWR-D-12-00168.1>.
- Wang, Y., J. Zhang, P.-L. Chang, C. Langston, B. Kaney, and L. Tang, 2016: Operational C-band dual-polarization radar QPE for the subtropical complex terrain of Taiwan. *Adv. Meteor.*, **2016**, 15, <https://doi.org/10.1155/2016/4294271>.
- Wayand, N. E., A. F. Hamlet, M. Hughes, S. I. Feld, and J. D. Lundquist, 2013: Intercomparison of meteorological forcing data from empirical and meso-scale model sources in the North Fork American River basin in northern Sierra Nevada, California. *J. Hydrometeorol.*, **14**, 677–699, <https://doi.org/10.1175/JHM-D-12-0102.1>.
- , J. Stemberis, J. P. Zagrodnik, C. F. Mass, and J. D. Lundquist, 2016: Improving simulations of precipitation phase and snowpack at a site subject to cold air intrusions: Snoqualmie Pass, WA. *J. Geophys. Res. Atmos.*, **121**, 9929–9942, <https://doi.org/10.1002/2016JD025387>.
- , M. P. Clark, and J. D. Lundquist, 2017: Diagnosing snow accumulation errors in a rain-snow transitional environment with snow board observations. *Hydrol. Processes*, **31**, 349–363, <https://doi.org/10.1002/hyp.11002>.
- Wen, Y., A. Behrangi, H. Chen, and B. Lambriksen, 2018: How well were the early 2017 California Atmospheric River precipitation events captured by satellite products and ground-based radars? *Quart. J. Roy. Meteor. Soc.*, **144**, 344–359, <https://doi.org/10.1002/qj.3253>.
- Westrick, K. J., and C. F. Mass, 2001: An evaluation of a high-resolution hydrometeorological modeling system for prediction of a cool-season flood event in a coastal mountainous watershed. *J. Hydrometeorol.*, **2**, 161–180, [https://doi.org/10.1175/1525-7541\(2001\)002<0161:AEOAHR>2.0.CO;2](https://doi.org/10.1175/1525-7541(2001)002<0161:AEOAHR>2.0.CO;2).
- , —, and B. A. Colle, 1999: The limitations of the WSR-88D radar network for quantitative precipitation measurement over the coastal western United States. *Bull. Amer. Meteor. Soc.*, **80**, 2289–2298, [https://doi.org/10.1175/1520-0477\(1999\)080<2289:TL OTWR>2.0.CO;2](https://doi.org/10.1175/1520-0477(1999)080<2289:TL OTWR>2.0.CO;2).
- , P. Storck, and C. Mass, 2002: Description and evaluation of a hydrometeorological forecast system for mountainous watersheds. *Wea. Forecasting*, **17**, 250–262, [https://doi.org/10.1175/1520-0434\(2002\)017<0250:DAEOAH>2.0.CO;2](https://doi.org/10.1175/1520-0434(2002)017<0250:DAEOAH>2.0.CO;2).
- Willie, D., and Coauthors, 2017: Evaluation of multisensor quantitative precipitation estimation in Russian River Basin. *J. Hydrol. Eng.*, **22**, E5016002, [https://doi.org/10.1061/\(ASCE\)HE.1943-5584.0001422](https://doi.org/10.1061/(ASCE)HE.1943-5584.0001422).
- Winstral, A., K. Elder, and R. E. Davis, 2002: Spatial snow modeling of wind-redistributed snow using terrain-based parameters. *J. Hydrometeorol.*, **3**, 524–538, [https://doi.org/10.1175/1525-7541\(2002\)003<0524:SS MOWR>2.0.CO;2](https://doi.org/10.1175/1525-7541(2002)003<0524:SS MOWR>2.0.CO;2).

- Wrzesien, M. L., T. M. Pavelsky, S. B. Kapnick, M. T. Durand, and T. H. Painter, 2015: Evaluation of snow cover fraction for regional climate simulations in the Sierra Nevada. *Int. J. Climatol.*, **35**, 2472–2484, <https://doi.org/10.1002/joc.4136>.
- , M. T. Durand, T. M. Pavelsky, I. M. Howat, S. A. Margulis, and L. S. Huning, 2017: Comparison of methods to estimate snow water equivalent at the mountain range scale: A case study of the California Sierra Nevada. *J. Hydrometeorol.*, **18**, 1101–1119, <https://doi.org/10.1175/JHM-D-16-0246.1>.
- , —, —, S. B. Kapnick, Y. Zhang, J. Guo, and C. K. Shum, 2018: A new estimate of North American mountain snow accumulation from regional climate model simulations. *Geophys. Res. Lett.*, **45**, 1423–1432, <https://doi.org/10.1002/2017GL076664>.
- Xia, Y., 2008: Adjustment of global precipitation data for orographic effects using observed annual streamflow and the LaD model. *J. Geophys. Res.*, **113**, D04106, <https://doi.org/10.1029/2007JD008545>.
- Xu, Z., A. M. Rhoades, H. Johansen, P. A. Ullrich, and W. D. Collins, 2018: An intercomparison of GCM and RCM dynamical downscaling for characterizing the hydroclimatology of California and Nevada. *J. Hydrometeorol.*, **19**, 1485–1506, <https://doi.org/10.1175/JHM-D-17-0181.1>.
- Xue, X., and Coauthors, 2013: Statistical and hydrological evaluation of TRMM-based Multi-satellite Precipitation Analysis over the Wangchu Basin of Bhutan: Are the latest satellite precipitation products 3B42V7 ready for use in ungauged basins? *J. Hydrol.*, **499**, 91–99, <https://doi.org/10.1016/j.jhydrol.2013.06.042>.
- Yang, J., K. F. Jones, W. Yu, and R. Morris, 2012: Simulation of in-cloud icing events on Mount Washington with the GEM-LAM. *J. Geophys. Res.*, **117**, D17204, <https://doi.org/10.1029/2012JD017520>.
- Ye, B., D. Yang, and L. Ma, 2012: Effect of precipitation bias correction on water budget calculation in Upper Yellow River, China. *Environ. Res. Lett.*, **7**, 025201, <https://doi.org/10.1088/1748-9326/7/2/025201>.
- Yong, B., and Coauthors, 2012: Assessment of evolving TRMM-based multisatellite real-time precipitation estimation methods and their impacts on hydrologic prediction in a high latitude basin. *J. Geophys. Res.*, **117**, D09108, <https://doi.org/10.1029/2011JD017069>.
- Yoshitani, J., Z. Chen, M. Kavvas, and K. Fukami, 2009: Atmospheric model-based streamflow forecasting at small, mountainous watersheds by a distributed hydrologic model: Application to a watershed in Japan. *J. Hydrol. Eng.*, **14**, 1107–1118, [https://doi.org/10.1061/\(ASCE\)HE.1943-5584.0000111](https://doi.org/10.1061/(ASCE)HE.1943-5584.0000111).
- Young, C. B., B. R. Nelson, A. A. Bradley, J. A. Smith, C. D. Peters-Lidard, A. Kruger, and M. L. Baeck, 1999: An evaluation of NEXRAD precipitation estimates in complex terrain. *J. Geophys. Res.*, **104**, 19691–19703, <https://doi.org/10.1029/1999JD900123>.
- Zhang, J., Y. Qi, D. Kingsmill, and K. Howard, 2012: Radar-based quantitative precipitation estimation for the cool season in complex terrain: Case studies from the NOAA Hydrometeorology Testbed. *J. Hydrometeorol.*, **13**, 1836–1854, <https://doi.org/10.1175/JHM-D-11-0145.1>.
- , and Coauthors, 2016: Multi-Radar Multi-Sensor (MRMS) quantitative precipitation estimation: Initial operating capabilities. *Bull. Amer. Meteor. Soc.*, **97**, 621–638, <https://doi.org/10.1175/BAMS-D-14-00174.1>.
- Zhang, X., and E. N. Anagnostou, 2019: Evaluation of numerical weather model-based satellite precipitation adjustment in tropical mountainous regions. *J. Hydrometeorol.*, **20**, 431–445, <https://doi.org/10.1175/JHM-D-18-0008.1>.
- Zreda, M., D. Desilets, T. Ferré, and R. L. Scott, 2008: Measuring soil moisture content non-invasively at intermediate spatial scale using cosmic-ray neutrons. *Geophys. Res. Lett.*, **35**, L21402, <https://doi.org/10.1029/2008GL035655>.