EXPLAINING EXTREME EVENTS OF 2018

From a Climate Perspective

Special Supplement to the
Bulletin of the American Meteorological Society
Vol. 101, No. 1, January 2020

EXPLAINING EXTREME EVENTS OF 2018 FROM A CLIMATE PERSPECTIVE

Editors

Stephanie C. Herring, Nikolaos Christidis, Andrew Hoell,
Martin P. Hoerling, and Peter A. Stott

BAMS Special Editors for Climate

Andrew King, Thomas Knutson,
John Nielsen-Gammon, and Friederike Otto

Special Supplement to the

Bulletin of the American Meteorological Society

Vol. 101, No. 1, January 2020

AMERICAN METEOROLOGICAL SOCIETY

CORRESPONDING EDITOR:

Stephanie C. Herring, PhD
NOAA National Centers for Environmental Information
325 Broadway, E/CC23, Rm 1B-131
Boulder, CO, 80305-3328
E-mail: stephanie.herring@noaa.gov

COVER CREDIT: iStock.com/Alena Kravchenko—River Thames receded during a heatwave in summer 2018 in London, United Kingdom.

HOW TO CITE THIS DOCUMENT

Citing the complete report:

Herring, S. C., N. Christidis, A. Hoell, M. P. Hoerling, and P. A. Stott, Eds., 2020: Explaining Extreme Events of 2018 from a Climate Perspective. *Bull. Amer. Meteor. Soc.*, **101** (1), S1–S128, doi:10.1175/BAMS-ExplainingExtremeEvents2018.1.

Citing a section (example):

Mahoney, K., 2020: Extreme Hail Storms and Climate Change: Foretelling the Future in Tiny, Turbulent Crystal Balls? [in “Explaining Extremes of 2018 from a Climate Perspective”]. *Bull. Amer. Meteor. Soc.*, **101** (1), S17–S22, doi:10.1175/BAMS-D-19-0233.1.

TABLE OF CONTENTS

1. The Extreme 2018 Northern California Fire Season	I
2. Anthropogenic Impacts on the Exceptional Precipitation of 2018 in the Mid-Atlantic United States	5
3. Quantifying Human-Induced Temperature Impacts on the 2018 United States Four Corners Hydrologic and Agro-Pastoral Drought	11
4. Extreme Hail Storms and Climate Change: Foretelling the Future in Tiny, Turbulent Crystal Balls?	17
5. The Extremely Cold Start of the Spring of 2018 in the United Kingdom	23
6. The Exceptional Iberian Heatwave of Summer 2018	29
7. Analyses of the Northern European Summer Heatwave of 2018	35
8. Anthropogenic Influence on the 2018 Summer Warm Spell in Europe: The Impact of Different Spatio-Temporal Scales	41
9. On High Precipitation in Mozambique, Zimbabwe and Zambia in February 2018	47
10. The Record Low Bering Sea Ice Extent in 2018: Context, Impacts, and an Assessment of the Role of Anthropogenic Climate Change	53
11. The Late Spring Drought of 2018 in South China	59
12. Anthropogenic Influence on 2018 Summer Persistent Heavy Rainfall in Central Western China	65
13. Conditional Attribution of the 2018 Summer Extreme Heat over Northeast China: Roles of Urbanization, Global Warming, and Warming-Induced Circulation Changes	71
14. Effects of Anthropogenic Forcing and Natural Variability on the 2018 Heatwave in Northeast Asia	77
15. Anthropogenic Influences on the Persistent Night-Time Heat Wave in Summer 2018 over Northeast China	83
16. Anthropogenic Contributions to the 2018 Extreme Flooding over the Upper Yellow River Basin in China	89
17. Attribution of the Record-Breaking Consecutive Dry Days in Winter 2017/18 in Beijing	95
18. Quantifying Human Impact on the 2018 Summer Longest Heat Wave in South Korea	103
19. The Heavy Rain Event of July 2018 in Japan Enhanced by Historical Warming	109
20. Deconstructing Factors Contributing to the 2018 Fire Weather in Queensland, Australia	115
21. A 1-Day Extreme Rainfall Event in Tasmania: Process Evaluation and Long Tail Attribution	123
22. Attribution of the Warmest Spring of 2018 in Northeastern Asia Using Simulations of a Coupled and an Atmospheric Model	129
23. Attribution of the 2018 October–December Drought over South Southern Africa	135

ATTRIBUTION OF THE 2018 OCTOBER–DECEMBER DROUGHT OVER SOUTH SOUTHERN AFRICA

SHINGIRAI NANGOMBE, TIANJUN ZHOU, LIXIA ZHANG, AND WENXIA ZHANG

The HadGEM3 (CAM5) model suggests that anthropogenic emissions increased the chances of the October–December 2018 drought over the southernmost area of southern Africa by 1.4 (0.97–1.96) times [4.3 (3.43–5.46) times], with higher influence on reference evapotranspiration than on precipitation.

Across Africa, droughts are a major type of natural disaster and a household insecurity trigger. The south southern Africa region (SSA; encompassed by the 22°–35°S, 18°–32°E rectangle) experienced extremely poor rainfall during the first half [i.e., October–December 2018 (OND)] of the October 2018 to March 2019 rainfall season (Fig. 1a). The OND subseason marks the important first half of southern Africa's rainfall season (Landman and Mason 1999). Late rainfall onset dates normally lead to below normal seasonal rainfall in southern Africa (Dunning et al. 2016). Dry conditions during OND 2018 led to a 2-month delay in sowing in parts of southern Africa (SADC CSC 2018), which translated

to reduced regional crop yield, where livelihoods are principally agriculture-based. In South Africa's large cereal-producing provinces, 40%–50% reduced rains delayed plantings and curbed sown areas (FAO 2019; FEWSNET 2019). Late rain onset acted as a disincentive to farmers to take risks of sowing outside of the optimal planting window (FAO 2019). This contributed to Zambia and South Africa (the main regional grain suppliers) recording below-average harvests, thereby reducing exportable surplus from 7.5 to 1.4 million tons (SADC RVAA 2019), hence affecting regional food security.

The poor rains were also accompanied by water loss through above average reference evapotranspiration (Fig. 1b). This exacerbated the drought impact, considering the region's large population. Water loss worsened the water supply situation since dam and river water levels were already low due to the three previous consecutive droughts (Richman and Leslie 2018).

This study investigates how extreme the OND 2018 SSA drought was, and explores the related anthropogenic influences.

DATA AND METHODS. Gridded observational temperature, from the Global Historical Climatology Network Gridded v2 and Climate Anomaly Monitoring System dataset (GHCN-CAMS; Fan and van den Dool 2008) is used here available at 0.5° horizontal grid resolution. A precipitation average of the Global Precipitation Climatology Centre v2018 (GPCC-FD v2018; Schneider et al. 2015) dataset available at 1° horizontal grid resolution and the Climate Research Unit v4.0.3 (CRU-TS 4.03; Harris et al. 2014) dataset available since 1901 at 0.5° horizontal resolution is also utilized.

Two C20C+ Detection and Attribution project (Folland et al. 2014; Stone et al. 2019) global mod-

AFFILIATIONS: NANGOMBE—LASG, Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing, China; Meteorological Services Department, Harare, Zimbabwe; University of Chinese Academy of Sciences, Beijing, China; ZHOU—LASG, Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing, China; CAS Centre for Excellence in Tibetan Plateau Earth Sciences, Chinese Academy of Sciences, Beijing, China; University of Chinese Academy of Sciences, Beijing, China; ZHANG—LASG, Institute of Atmospheric Physics, and CAS Centre for Excellence in Tibetan Plateau Earth Sciences, Chinese Academy of Sciences, Beijing, China; ZHANG—LASG, Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing, China

CORRESPONDING AUTHOR: Shingirai Nangombe, ssnangombe@lasg.iap.ac.cn

DOI:10.1175/BAMS-D-19-0179.1

A supplement to this article is available online (10.1175/BAMS-D-19-0179.2)

© 2020 American Meteorological Society
For information regarding reuse of this content and general copyright information, consult the [AMS Copyright Policy](#).

FIG. 1. The OND 2018 anomalies of (a) precipitation derived from the GPCC and CRU averaged datasets, (b) reference evapotranspiration calculated from GHCN-CAMS, and (c) difference between precipitation and reference evapotranspiration (P -refET) denoting the drought index. (d) Map of geopotential height (shading; hPa) and moisture transport (vector; $\text{kg kg}^{-1} \text{m s}^{-1}$) anomalies at 850 hPa during OND 2018 derived from the JRA-55 reanalysis (Kobayashi et al. 2015). (e) SSA area-averaged OND P (blue) and P -refET (red) with overlaid refET (sky-blue and brown bars). (f) Map of OND 2018 outgoing longwave radiation flux anomalies (W m^{-2}). SSA is shown by a black box in (a)–(d) and (f). All the anomalies are calculated from a 1960–2013 climatology. The units are in mm day^{-1} unless stated otherwise.

els are employed here in evaluating anthropogenic influence on this drought occurrence: HadGEM3 (Ciavarella et al. 2018) and CAM5 (Risser et al. 2018). These comprise two suites of simulations representing the actual climate (All-Hist) and a hypothetical natural climate excluding anthropogenic influence (Nat-Hist). In HadGEM3, both All-Hist and Nat-Hist have 15 realizations ranging from 1960 to 2013; in CAM5, there are 50 realizations ranging from 1960 to 2017.

To factor in high temperatures, a drought index that incorporates reference evapotranspiration is used: specifically, OND averaged P -refET, where P and refET denote precipitation and reference evapotranspiration, respectively. The Hamon method is used to estimate refET (Haith and Shoemaker 1987) from temperature data using the *refevt_hamon* function (Brown et al. 2019). The online supplemental

information and Eq. (7) in Haith and Shoemaker (1987) provide details of refET calculation.

The Gaussian distribution (Wellmer 1998) was fitted to the observed OND averaged data, and return period of the 2018 event derived, separately for P , refET, and P -refET. Using 810 and 2900 model years from HadGEM3 and CAM5 All-Hist simulations respectively, OND averaged P , refET, and P -refET indices were extracted. Then Gaussian distributions are fitted to the All-Hist-based indices and used to derive the drought events that have similar return periods as the 2018 observed one. Last, as in Stott et al. (2004) and Ma et al. (2017), the probability ratio ($\text{PR} = P_1 / P_0$) and the fraction of attributable risk ($\text{FAR} = 1 - P_0 / P_1$) are estimated, where P_1 (P_0) is the likelihood of an event occurring in the All-Hist (Nat-Hist) simulations. Here, a null hypothesis of existing human influence against an alternative hypothesis of no human influence on the drought occurrence is used.

However, in some cases like precipitation where high model uncertainty exists, the null hypothesis may be disproved. A total of 1,000 bootstrapping samples are used to estimate the 5th–95th confidence intervals for PR and FAR.

RESULTS. The observed OND precipitation area-averaged over SSA in 2018 was 0.68 mm day^{-1} (46%) below normal (1960–2013 baseline), while refET was 0.23 mm day^{-1} above normal (Fig. 1e). These precipitation and refET anomalies were ranked as the second lowest and third highest since 1950, respectively. Consequently, P -refET was the second record-breaking event with $-0.91 \text{ mm day}^{-1}$ (−60%) anomaly (Figs. 1c,e) and a 72-yr return period. Precipitation (refET) had a 40-yr (32-yr) return period (not shown).

SSA in OND is normally characterized by northeasterly moisture transport from the tropical Indian Ocean (Fig. S1a), leading to precipitation. However, in 2018, there was weak southerly moisture flux into SSA (Fig. 1d). A cyclonic anomaly in the Indian and Atlantic Oceans dominated the 850-mb moisture flux, hence diverting moisture flow away from SSA (Fig. 1d). Moreover, late 2018 was associated with El Niño conditions (Fig. S1b). Typically, during El Niño in austral summer, southern Africa receives drought (Hoell and Cheng 2018; Manatsa et al. 2017) while temperatures are usually above average (Nangombe et al. 2018). Likewise, Figs. S1c and S1d show a positive (negative) relationship between the OND ENSO index and OND SSA refET (*P*-refET) index.

Meanwhile, using outgoing longwave radiation (OLR) as a proxy for tropical temperate trough (TTT) events, anomalously high OLR over SSA in OND 2018 indicates a decrease in convection linked to TTTs (Fig. 1f). TTTs are the main rainfall bearers to SSA (Cook et al. 2004). Therefore, sea surface temperatures in the Pacific and Indian Oceans accompanied by anomalous low TTTs over SSA contributed to below normal 2018 OND precipitation in SSA.

On evaluating the model performance, correlation coefficients of *P*-refET, *P*, and refET area-averaged over SSA between the HadGEM3 (CAM5) All-Hist ensemble-mean and observation are 0.36, 0.31, and 0.38 (0.34, 0.21, and -0.1), respectively, with about 1 to 2 mm day⁻¹ overestimation of the models. All the correlations based on the HadGEM3 (CAM5) model are significant (insignificant) at the 0.05 level. Nonetheless, results from both models under All-Hist and Nat-Hist cover the observed range (Figs. S2a,b). Additionally, there are statistical similarities between the density distributions of the All-Hist simulations and observation/reanalysis (Figs. S2c,d), hence our confidence in using these models.

In exploring the anthropogenic influence, CAM5 depicts a distinct leftward shift of the probability functions of the OND *P*-refET in All-Hist compared to Nat-Hist, while HadGEM3 suggests a slight shift (Figs. 2a,b). This signifies a likely anthropogenic influence on *P*-refET in the SSA. The shift is even more pronounced for refET, albeit in the rightward direction (Figs. 2e,f). Conversely, for precipitation, there is no obvious shift in All-Hist compared to Nat-Hist simulations of HadGEM3, while CAM5 depicts a moderate shift (Figs. S2e,f). In HadGEM3 (CAM5), -2.02 mm day⁻¹ (-1.79 mm day⁻¹) of *P*-refET is selected as the threshold for attribution of SSA drought since it has a return period similar to the observed 2018 *P*-refET. For refET (*P*), the thresholds used are

3.32 (1.33) mm day⁻¹ and 3.59 (1.84) mm day⁻¹, respectively. The HadGEM3 (CAM5) model estimates PR and FAR to be 1.36 (0.97–1.96) [4.34 (3.43–5.46)] and 26% (-3% to 49%) [77% (71%–82%)], respectively, for *P*-refET. The attributable influence from anthropogenic forcings is weaker in HadGEM3 than in CAM5, as the lower bound of the confidence interval of PR from HadGEM3 is slightly less than 1. Nonetheless, HadGEM3 simulations suggest PR and FAR values of 157.16 (86.90–310.74) and 99% (99%–100%) for refET while CAM5 suggests 105.75 (77.04–146.45) and 99% (99%–99%), respectively. We note that the PR and FAR estimates calculated from the CAM5 simulations are substantially affected by the nonstationarity of the ALL-Hist simulations (in terms of greenhouse gas concentrations) compared to estimates from the HadGEM3 simulations (not shown).

Furthermore, using simulations from composite El Niño years in estimating *P*-refET, HadGEM3 and CAM5 estimate PR values of 0.75 (0.24–2.77) and 2.19 (0.68–8.03) respectively. Thus, comparing with the other PR values, this portrays a reduction in the anthropogenic influence on drought of similar magnitude to that of OND 2018 in SSA during El Niño years.

Examining precipitation associated with the 2018 drought, CAM5 estimates the related PR (FAR) to be 1.95 (1.63–2.33) [49% (39%–57%)], whereas HadGEM3 suggests 0.76 (0.59–0.98) and -32% (-2% to -69%) PR (FAR) values, respectively, hence implying a reduced probability of occurrence due to anthropogenic forcing. Thus, comparing the two models, the anthropogenic forcing impact on precipitation-based drought in SSA is uncertain. Also, Yuan et al. (2018) highlighted that the major sources of uncertainty in detecting/attributing flash droughts over southern Africa are soil moisture and precipitation variation. Plus, the interannual variability of precipitation is much higher in African regions, hence making it difficult to distinguish anthropogenic signal from noise (Otto et al. 2013). Nonetheless, although low precipitation seasons such as OND 2018 are not necessarily more likely due to anthropogenic emissions, they may have severe impacts in an anthropogenically warmed world due to increased temperatures and hence increased refET.

CONCLUDING REMARKS. SSA experienced an extreme drought event in OND 2018 accompanied by extremely low rainfall and high reference evapotranspiration. This translated to substantial delay in the planting season hence affected the regional 2018/19 crop yield where livelihoods are largely agriculture-based.

FIG. 2. HadGEM3's probability density functions (PDFs) of October–December (a) *P*-refET from all years, (c) *P*-refET El Niño years, (e) refET all years, and (g) refET El Niño years, area-averaged over the SSA region. (b),(d),(f),(h) As in (a),(c),(e),(g), but for the CAM5 model. All-Hist simulations and Nat-Hist simulations are represented by red and blue PDFs, respectively. The PDFs of HadGEM3 are based on 15-member realizations of both All-Hist and Nat-Hist ranging from 1960 to 2013 while CAM5 PDFs are based on 50-member realizations of both All-Hist and Nat-Hist ranging from 1959 to 2016. The dashed lines are the thresholds similar to the observed 2018 values. PDFs are calculated by the Gaussian kernel density estimator.

Here, the HadGEM3 model suggests that anthropogenic influence may have increased the probability of SSA droughts, such as the OND 2018 drought (*P*-refET), by 1.4 times [0.97, 1.96], and the CAM5 model indicates 4.3 times [3.43, 5.46]. The high anthropogenic influence on refET mostly contributed to this drought occurrence. El Niño conditions may also have had an influence. Therefore, the increase in temperature accompanied by rainfall decrease projected over southern Africa might result in increase in drought occurrence of at least OND 2018 magnitudes in the future (Kusangaya et al. 2014; Niang et al. 2014; Maure et al. 2018; Nikulin et al. 2018; Nangombe et al. 2018).

ACKNOWLEDGMENTS. This research was supported by the National Natural Science Foundation of China Grants 41661144009 and 41675076. The first author is a recipient of a research studentship provided by the CAS-TWAS President fellowship. Special thanks go to the C20C+ Detection and Attribution project for providing global models we used.

REFERENCES

- Allen, R. G., L. S. Pereira, D. Raes, and M. Smith, 1998: Crop evapotranspiration—Guidelines for computing crop water requirements. FAO Irrigation and Drainage Paper 56, 300 pp., www.fao.org/docrep/X0490E/X0490E00.htm.
- Brown, D., R. Brownrigg, M. Haley, and W. Huang, 2019: The NCAR NCAR Command Language (NCL) (version 6.6.2). UCAR/NCAR Computational and Information Systems Laboratory. Command Language. <https://doi.org/10.5065/D6WD3XH>
- Ciavarella, A., and Coauthors, 2018: Upgrade of the HadGEM3-A based attribution system to high resolution and a new validation framework for probabilistic event attribution. *Wea. Climate Extremes*, **20**, 9–32, <https://doi.org/10.1016/j.wace.2018.03.003>.
- Compo, G. P., and Coauthors, 2011: The Twentieth Century Reanalysis Project. *Quart. J. Roy. Meteor. Soc.*, **137**, 1–28, <https://doi.org/10.1002/qj.776>.
- Cook, C., C. J. C. Reason, and B. C. Hewitson, 2004: Wet and dry spells within particularly wet and dry summers in the South African summer rainfall region. *Climate Res.*, **26**, 17–31, <https://doi.org/10.3354/cr026017>.
- Fan, Y., and H. van den Dool, 2008: A global monthly land surface air temperature analysis for 1948–present. *J. Geophys. Res.*, **113**, D01103, <https://doi.org/10.1029/2007JD008470>.
- Dunning, C. M., E. C. L. Black, and R. P. Allan, 2016: The onset and cessation of seasonal rainfall over Africa. *J. Geophys. Res.*, **121**, 11 405–11 424, <https://doi.org/10.1002/2016JD025428>
- FAO, 2019: Southern Africa: Dry weather conditions reduce agricultural production prospects in 2019, Food and Agriculture Organization of the United Nations, 4 pp., https://reliefweb.int/sites/reliefweb.int/files/resources/ca3071en_0.pdf.
- FEWSNET, 2019: Delayed start of season and significant rainfall deficits negatively impacting cropping conditions. Famine Early Warning System Network, 2 pp., <https://fewsn.net/southern-africa/alert/february-1-2019>.
- Folland, C., D. Stone, C. Frederiksen, D. Karoly, and J. Kinter, 2014: The International CLIVAR Climate of the 20th Century Plus (C20C+) Project: Report of the Sixth Workshop. *CLIVAR Exchanges*, CLIVAR, Southampton, United Kingdom, 57–59, https://crd.lbl.gov/assets/pubs_presos/FollandCStoneDetalii2014.pdf.
- Haith, D. A., and L. L. Shoemaker, 1987: Generalized watershed loading functions for stream flow nutrients. *J. Amer. Water Resour. Assoc.*, **23**, 471–478, <https://doi.org/10.1111/j.1752-1688.1987.tb00825.x>.
- Harris, I., P. D. Jones, T. J. Osborn, and D. H. Lister, 2014: Updated high-resolution grids of monthly climatic observations—The CRU TS3.10 dataset. *Int. J. Climatol.*, **34**, 623–642, <https://doi.org/10.1002/joc.3711>.
- Hoell, A., and L. Cheng, 2018: Austral summer southern Africa precipitation extremes forced by the El Niño–Southern Oscillation and the subtropical Indian Ocean dipole. *Climate Dyn.*, **50**, 3219–3236, <https://doi.org/10.1007/s00382-017-3801-z>.
- Kalnay, E., and Coauthors, 1996: The NCEP/NCAR 40-Year Reanalysis Project. *Bull. Amer. Meteor. Soc.*, **77**, 437–471, [https://doi.org/10.1175/1520-0477\(1996\)077<0437:TNYRP>2.0.CO;2](https://doi.org/10.1175/1520-0477(1996)077<0437:TNYRP>2.0.CO;2).
- Kobayashi, S., and Coauthors, 2015: The JRA-55 reanalysis: General specifications and basic characteristics. *J. Meteor. Soc. Japan*, **93**, 5–48, <https://doi.org/10.2151/jmsj.2015-001>.
- Kusangaya, S., M. L. Warburton, E. A. van Garderen, and G. P. Jewitt, 2014: Impacts of climate change on water resources in southern Africa: A review. *Phys. Chem. Earth*, **67–69**, 47–54, <https://doi.org/10.1016/j.pce.2013.09.014>.
- Landman, W. A., and S. J. Mason, 1999: Operational long-lead prediction of South African rainfall using canonical correlation analysis. *Int. J. Climatol.*, **19**, 1073–1090, [https://doi.org/10.1002/\(SICI\)1097-0088\(199908\)19:10<1073::AID-JOC415>3.0.CO;2-J](https://doi.org/10.1002/(SICI)1097-0088(199908)19:10<1073::AID-JOC415>3.0.CO;2-J)
- Ma, S., T. Zhou, O. Angéil, and H. Shiogama, 2017: Increased chances of drought in southeastern periphery of the Tibetan Plateau induced by anthropogenic

- warming. *J. Climate*, **30**, 6543–6560, <https://doi.org/10.1175/JCLI-D-16-0636.1>.
- Manatsa, D., T. Mushore, and A. Lenouo, 2017: Improved predictability of droughts over southern Africa using the standardized precipitation evapotranspiration index and ENSO. *Theor. Appl. Climatol.*, **127**, 259–274, <https://doi.org/10.1007/s00704-015-1632-6>.
- Maúre, G., and Coauthors, 2018: The southern African climate under 1.5°C and 2°C of global warming as simulated by CORDEX regional climate models. *Environ. Res. Lett.*, **13**, 065002, <https://doi.org/10.1088/1748-9326/aab190>.
- Nangombe, S., T. Zhou, W. Zhang, B. Wu, S. Hu, L. Zou, and D. Li, 2018: Record-breaking climate extremes in Africa under stabilized 1.5°C and 2°C global warming scenarios. *Nat. Climate Change*, **8**, 375–380, <https://doi.org/10.1038/s41558-018-0145-6>.
- Niang, I., O. C., Ruppel, M. A., Abdrabo, C. Essel, J. Padgham, P. Urquhart, K. K. E. Abdrabo, 2014: *Africa. Climate Change 2014: Impacts, Adaptation and Vulnerability*, Cambridge University Press, 1199–1265, <https://doi.org/10.1017/CBO9781107415386.002>.
- Nikulin, G., and Coauthors, 2018: The effects of 1.5 and 2 degrees of global warming on Africa in the CORDEX ensemble. *Environ. Res. Lett.*, **13**, 065003, <https://doi.org/10.1088/1748-9326/aab1b1>.
- Otto, F. E. L., R. G. Jones, K. Halladay, and M. R. Allen, 2013: Attribution of changes in precipitation patterns in African rainforests. *Philos. Trans. Roy. Soc. London*, **B368**, 20120299 <https://doi.org/10.1098/rstb.2012.0299>.
- Poli, P., and Coauthors, 2016: ERA-20C: An atmospheric reanalysis of the twentieth century. *J. Climate*, **29**, 4083–4097, <https://doi.org/10.1175/JCLI-D-15-0556.1>.
- Richman, M. B., and L. M. Leslie, 2018: The 2015–2017 Cape Town drought: Attribution and prediction using machine learning. *Procedia Comput. Sci.*, **140**, 248–257, <https://doi.org/10.1016/j.procs.2018.10.323>.
- Risser, M. D., and Coauthors, 2018: A basis set for exploration of sensitivity to prescribed ocean conditions for estimating human contributions to extreme weather in CAM5.1-1degree. *Wea. Climate Extremes*, **19**, 10–19, <https://doi.org/10.1016/j.wace.2017.12.003>.
- SADC CSC, 2018: Early warning bulletin update of the 2018-19 southern Africa regional rainfall season. Southern African Development Community Climate Services Centre, 14 pp., https://www.sadc.int/files/8715/4764/2883/SADC_CSC_Regional_Early_Warning_Bulletin_Update_for_2018-19_Rainfall_Season.pdf.
- SADC RVAA, 2019: Synthesis report on the state of food and nutrition security and vulnerability in southern Africa 2019. Southern African Development Community Regional Vulnerability Assessment and Analysis Programme, 21 pp., https://reliefweb.int/sites/reliefweb.int/files/resources/SADC_2019_Synthesis_Report_on_the_State_of_Food_and_Nutrition_Security_and_Vulnerability_in_Southern_Africa.pdf.
- Schneider, U., and Coauthors, 2015: GPCC full data reanalysis version 7.0 at 0.5°: Monthly land-surface precipitation from rain-gauges built on GTS-based and historic data. Global Precipitation Climatology Centre, accessed 14 April 2019, https://doi.org/10.5676/DWD_GPCC/FD_M_V6_050.
- Stone, D. A., and Coauthors, 2019: Experiment design of the international CLIVAR C20C+ detection and attribution project. *Wea. Climate Extremes*, **24**, 100206 <https://doi.org/10.1016/j.wace.2019.100206>.
- Stott, P. A., D. A. Stone, and M. R. Allen, 2004: Human contribution to the European heatwave of 2003. *Nature*, **432**, 610–614, <https://doi.org/10.1038/nature03089>.
- Wellmer, F. W., 1998: The normal distribution. *Statistical Evaluations in Exploration for Mineral Deposits*, Springer, 27–30, https://doi.org/10.1007/978-3-642-60262-7_4.
- Yuan, X., L. Wang, and E. F. Wood, 2018: Anthropogenic intensification of southern African flash droughts as exemplified by the 2015/16 season. *Bull. Amer. Meteor. Soc.*, **99**, S86–S90, <https://doi.org/10.1175/BAMS-D-17-0077.1>.