EXPLAINING EXTREME EVENTS OF 2018

From a Climate Perspective

Special Supplement to the
Bulletin of the American Meteorological Society
Vol. 101, No. 1, January 2020

EXPLAINING EXTREME EVENTS OF 2018 FROM A CLIMATE PERSPECTIVE

Editors

Stephanie C. Herring, Nikolaos Christidis, Andrew Hoell,
Martin P. Hoerling, and Peter A. Stott

BAMS Special Editors for Climate

Andrew King, Thomas Knutson,
John Nielsen-Gammon, and Friederike Otto

Special Supplement to the

Bulletin of the American Meteorological Society

Vol. 101, No. 1, January 2020

AMERICAN METEOROLOGICAL SOCIETY

CORRESPONDING EDITOR:

Stephanie C. Herring, PhD
NOAA National Centers for Environmental Information
325 Broadway, E/CC23, Rm 1B-131
Boulder, CO, 80305-3328
E-mail: stephanie.herring@noaa.gov

COVER CREDIT: iStock.com/Alena Kravchenko—River Thames receded during a heatwave in summer 2018 in London, United Kingdom.

HOW TO CITE THIS DOCUMENT

Citing the complete report:

Herring, S. C., N. Christidis, A. Hoell, M. P. Hoerling, and P. A. Stott, Eds., 2020: Explaining Extreme Events of 2018 from a Climate Perspective. *Bull. Amer. Meteor. Soc.*, **101** (1), S1–S128, doi:10.1175/BAMS-ExplainingExtremeEvents2018.1.

Citing a section (example):

Mahoney, K., 2020: Extreme Hail Storms and Climate Change: Foretelling the Future in Tiny, Turbulent Crystal Balls? [in “Explaining Extremes of 2018 from a Climate Perspective”]. *Bull. Amer. Meteor. Soc.*, **101** (1), S17–S22, doi:10.1175/BAMS-D-19-0233.1.

TABLE OF CONTENTS

1. The Extreme 2018 Northern California Fire Season	1
2. Anthropogenic Impacts on the Exceptional Precipitation of 2018 in the Mid-Atlantic United States	5
3. Quantifying Human-Induced Temperature Impacts on the 2018 United States Four Corners Hydrologic and Agro-Pastoral Drought	11
4. Extreme Hail Storms and Climate Change: Foretelling the Future in Tiny, Turbulent Crystal Balls?	17
5. The Extremely Cold Start of the Spring of 2018 in the United Kingdom.....	23
6. The Exceptional Iberian Heatwave of Summer 2018	29
7. Analyses of the Northern European Summer Heatwave of 2018.....	35
8. Anthropogenic Influence on the 2018 Summer Warm Spell in Europe: The Impact of Different Spatio-Temporal Scales	41
9. On High Precipitation in Mozambique, Zimbabwe and Zambia in February 2018.....	47
10. The Record Low Bering Sea Ice Extent in 2018: Context, Impacts, and an Assessment of the Role of Anthropogenic Climate Change	53
11. The Late Spring Drought of 2018 in South China.	59
12. Anthropogenic Influence on 2018 Summer Persistent Heavy Rainfall in Central Western China	65
13. Conditional Attribution of the 2018 Summer Extreme Heat over Northeast China: Roles of Urbanization, Global Warming, and Warming-Induced Circulation Changes	71
14. Effects of Anthropogenic Forcing and Natural Variability on the 2018 Heatwave in Northeast Asia.	77
15. Anthropogenic Influences on the Persistent Night-Time Heat Wave in Summer 2018 over Northeast China	83
16. Anthropogenic Contributions to the 2018 Extreme Flooding over the Upper Yellow River Basin in China	89
17. Attribution of the Record-Breaking Consecutive Dry Days in Winter 2017/18 in Beijing	95
18. Quantifying Human Impact on the 2018 Summer Longest Heat Wave in South Korea	103
19. The Heavy Rain Event of July 2018 in Japan Enhanced by Historical Warming.	109
20. Deconstructing Factors Contributing to the 2018 Fire Weather in Queensland, Australia	115
21. A 1-Day Extreme Rainfall Event in Tasmania: Process Evaluation and Long Tail Attribution	123

THE NORTHERN CALIFORNIA 2018 EXTREME FIRE SEASON

TIMOTHY BROWN, STEVE LEACH, BRENT WACHTER, AND BILLY GARDUNIO

The fire season of 2018 was the most extreme on record in Northern California in terms of the number of fatalities (95), over 22,000 structures destroyed, and over 600,000 ha burned (https://www.fire.ca.gov/media/5511/top20_destruction.pdf; accessed 24 November 2019). The most deadly and destructive fire in California history, the Camp Fire, occurred in Butte County in the Sierra Nevada foothills in early November, and caused 85 fatalities and destroyed nearly 19,000 structures. The largest fire complex in state history, the Mendocino Complex, which included the Ranch fire, the largest single fire in state history, burned nearly 186,000 ha. It occurred in July and August, killing one firefighter. In western Shasta County nearly 138,000 ha burned from July through September in the Carr, Hirz, and Delta Fires. These fires caused multiple closures of Interstate 5 and exhibited some of the most extreme fire behavior ever observed in California. The Carr Fire caused eight fatalities, including two firefighters and two workers supporting firefighting efforts, burned over 1,100 homes in west Redding, caused the evacuation of one-third of the city, and produced an extreme fire vortex with an Enhanced Fujita scale rating between 136 to 165 mph, making it arguably the strongest tornado type event in state history, and one of the strongest documented cases in the world (Lareau et al. 2018).

Actual weather during the onset of these large fires was not outside of what would normally be expected during periods of higher fire danger that occur during most fire seasons, as opposed to the extreme offshore wind event that led to the October 2017 wine country fires (Nauslar et al. 2018). The Tubbs fire during

that period caused 22 fatalities and the loss of 5,636 structures, and at the time was California's most destructive fire on record dating back to the early 1900s. Instead, a series of climate and fuels conditions aligned to create an exceptional and explosive fire environment between June to November 2018 in naturally fire-prone areas with a substantial wildland–urban interface—a zone where human development meets and intermingles with wildland fuels.

CLIMATE AND FUELS. Climate enables fire and weather drives fire. The 2018 Northern California fire season origins began during the winter of 2016/17. Precipitation, which was much above normal, exceeded the 90th percentile ranking across much of the state at low to middle elevations. Some of the higher elevations in the Sierra Nevada range were the wettest on record. November through April is climatologically the wet season in northern California. This particular wet winter ended the statewide hydrologic drought from 2011–17. The significance of this wet season was that it led to the second consecutive spring with above average fine fuel (grasses) crop across much of Northern California.

The 2017–18 Northern California wet season was drier than normal, and temperatures were generally above normal, with the warmest conditions from the Cascade–Sierra foothills through the Sacramento Valley and into the northern Coast Range. A lack of stormy weather during this period allowed above normal standing carry-over fine fuels from the previous wet year. April 2018 was warmer and wetter than normal. This was perfectly timed to produce above normal spring brush growth and grow a record fine fuel crop at elevations below 1,200 m, despite the overall wet season rainfall deficit (Fig. 1). This was the third straight spring with an above average fine fuel crop based on data taken at the University of California Sierra Foothill Research and Extension Center. The Brown's Valley, California, research station forage production data have been reported continuously since the 1979/80 growing season. Three straight above normal growing seasons had not occurred in the dataset prior to the 2017/18 growing season. Peak forage production toward the end of spring was 150% of normal (<https://ucanr>

AFFILIATIONS: BROWN—Western Regional Climate Center, Desert Research Institute, Reno, Nevada; LEACH—Bureau of Land Management, Redding, California; WACHTER AND GARDUNIO—USDA Forest Service, Redding, California
CORRESPONDING AUTHOR: Timothy Brown, tim.brown@dri.edu

DOI:10.1175/BAMS-D-19-0275.1

© 2020 American Meteorological Society
For information regarding reuse of this content and general copyright information, consult the [AMS Copyright Policy](#).

FIG. 1. Photo of fine fuel grass loading taken on 12 Sep 2018 about 60 miles northwest of the Camp Fire location. Left of the fence line shows a continuous bed of fine fuel. The paddock to the right of the fence line has less fuel due to grazing. Photo: Brent Wachter, Predictive Services.

[.shinyapps.io/SFREC_Web_Application/](https://shinyapps.io/SFREC_Web_Application/); accessed 24 November 2019) and the maximum value during the 40-yr history. As a testament to the abundant and continuous fine fuel growth, firefighters during 2018 often described arriving to a larger fire compared to what was normally observed during initial attack situations, thus making the fires harder to contain (Northern California Predictive Services, personal communication, 24 July 2019).

The summer of 2018 in Northern California had two notable climate anomalies—lack of rain and warmer than average temperatures. This led to a rapid curing of the annual fine fuels and drying of live fuels. For the most part, rain shut off in late April and did not return until late November. In the Sacramento Valley and foothills only two light rain events produced approximately 15 mm each, one in late May and the other in late September to early October. The lack of precipitation from late spring through the middle of November caused fuels to dry out to record levels.

The summer of 2018 was warmer than average in Northern California. Afternoon maximum temperatures were not extraordinarily high compared to 2017. Rather, it was the warm morning minimum temperatures that pushed the overall average temperature in July 2018 to record high levels. Record high lows for California were observed in July, continuing a positive trend of nighttime warming since 1895, (<https://www.climate.gov/news-features/event-tracker/extreme>

[-overnight-heat-california-and-great-basin-july-2018](#); accessed 24 November 2019). These warm temperatures led to more nights of poor overnight humidity recovery, which has direct implications on the flammability of fuel. The poor recoveries led to unusually long active burn periods during the overnights and set the stage for earlier active burning during the daytime hours. This created more hours of active to extreme fire behavior and provided less opportunity for direct attack by firefighters (Northern California Predictive Services, personal communication, 24 July 2019).

WEATHER. Long periods dominated by strong high pressure occurred in July, which hastened the summer drying of fuels. The hottest day of the year in the Sacramento Valley was 26 July, setting all-time records for the date, and was a key element in the conditions that rapidly brought the Carr Fire into Redding, California. Other large-scale fires started that week as well, including the Mendocino Complex. Interestingly, temperatures were slightly cooler than normal during August and September in areas where thick smoke from nearby wildfires persisted for unusually long periods. Northern California had a very low occurrence of lightning during the 2018 fire season. Roughly 12,000 lightning strikes were recorded during the fire season, compared to the long-term average of more than 30,000 strikes (in Northern California lightning accounts for the majority of fire

ignitions, while in Southern California human causes dominate ignitions; Balch et al. 2017). Dry offshore wind events, which typically increase in occurrence after the middle of September, occurred with fuels at extreme and record dry levels. The Camp Fire on 8 November was the result of an ignition occurring in extremely dry fuels during an enhanced strong and dry downslope wind pattern.

FIRE ENVIRONMENT. The precipitation deficit at the close of the 2017/18 rainy season combined with drier and warmer than normal summer conditions caused an expansion and intensification of short-term drought across northern California. The Evaporative Demand Drought Index (EDDI; Hobbins et al. 2016; McEvoy et al. 2016) clearly illustrates the intensity of short-term drought conditions during late July when significant fires started (Fig. 2a), and even more so in November as dry northeast/offshore wind patterns became more frequent and the Camp Fire broke out (Fig. 2b). Evaporative demand relates to the thirst of the atmosphere which directly corresponds to fuel stress (higher values equate to more stress). There is a close correlation between EDDI values and fire danger (McEvoy et al. 2019) including extreme fire behavior and spread rates. EDDI comprises inputs of temperature, humidity, wind speed, and solar

radiation. These are also key factors in determining dead fuel moisture calculated in the newly updated National Fire Danger Rating System (https://gacc.nifc.gov/eacc/predictive_services/fuels_fire-danger/documents/Overview%20of%20NFDRS2016%20and%20Implementation%20and%20Evaluation.pdf; accessed 24 November 2019). Starting in late July, fire danger indicators in northern California such as the energy release component (the amount of energy at the head of fire's flaming front) approached all-time record high values accentuating the impact of the anomalously warm and dry summer on fuels. In effect, fuel regimes found at all elevations (sea level to over 3,000 m) had a large fire potential by July.

SUMMARY. Both 2017 and 2018 were extreme fire years in California. While this paper has focused on Northern California in 2018, Southern California that year also saw its share of extreme fire including the Ferguson (two firefighter fatalities, 19 structures destroyed, and approximately 39,000 ha burned), Cranston, and Holy Fires. On the same day as the Camp Fire, 8 November, two large fires developed in Ventura and Los Angeles County, including the Woolsey Fire (approximately 39,000 ha burned and over 1,600 structures destroyed). Following the unusually late season December 2017 Thomas Fire in

FIG. 2. Evaporative Demand Drought Index 2-month categories for (a) 26 Jul 2018, and (b) 8 Nov 2018. Triangles represent the Mendocino (lower), Carr (upper), and Camp fire locations, respectively.

the Ventura and Santa Barbara county area, a post-fire debris flow event occurred over the burn area on 9 January resulting in 23 fatalities and 246 structures destroyed (Oakley et al. 2018).

Climate, fuels, and people are the three confluent factors for California's recent destructive wildfires. The wildland–urban interface has been growing extensively within the state during the past few decades (e.g., Radeloff et al. 2018), placing people in fire-prone areas. Climate is an enabler of wildfire by providing seasonal moisture to grow fuels, and seasonal warming and drying that increases fuel flammability. Increasing temperature trends enable longer and more extreme fire seasons. California's annual temperatures have been increasing substantially during the past four decades and are expected to continue warming this century (California's Fourth Climate Assessment; Bedsworth et al. 2018). Abatzoglou and Williams (2016) have shown that California (as well as all of the West) has had significantly enhanced fuel aridity due to anthropogenic increases in temperature and vapor pressure deficit over the past several decades. This can also be seen in the increasing number of days of fire weather season length based on fire danger indicators (e.g., Jolly et al. 2015). Nighttime temperature trends especially may be playing an important role in more extreme fire behavior. Research is currently underway to examine the specific relationship of this warming to nighttime fuel drying and subsequent extreme fire behavior that was observed during the California 2018 fire season.

An estimated 54% of California ecosystems are fire dependent and most of the rest are fire adaptive (Pyne 2016). California has always had fire given its climate, topography, and distinctive varieties of combustible vegetation. Today it is a state of nearly 40 million people, and one in four Californians live in a “high risk” wildfire area (<https://www.latimes.com/local/lanow/la-me-california-braces-for-new-wildfires-20190614-story.html>; accessed 24 November 2019). California insured losses in 2018 from wildfire topped \$13 billion (<https://www.insurancejournal.com/news/west/2019/05/08/525930.htm>; accessed 24 November 2019). The year 2018 now holds the record for the most destructive wildfire, the largest wildfire, and the costliest wildfire season in California state history.

REFERENCES

- Abatzoglou, J. T., and A. P. Williams, 2016: Impact of anthropogenic climate change on wildfire across western US forests. *Proc. Natl. Acad. Sci. USA*, **113**, 11 770–11 775, <https://doi.org/10.1073/pnas.1607171113>.
- Balch, J. K., B. A. Bradley, J. T. Abatzoglou, R. C. Nagya, E. J. Fusco, and A. L. Mahood, 2017: Human-started wildfires expand the fire niche across the United States. *Proc. Natl. Acad. Sci. USA*, **114**, 2946–2951, <https://doi.org/10.1073/pnas.1617394114>.
- Bedsworth, L., D. Cayan, G. Franco, L. Fisher, and S. Ziaja, 2018: Statewide Summary Report. California's Fourth Climate Change Assessment, California Public Utilities Commission Publication SUM- CCCA4-2018-013, 133 pp.
- Hobbins, M., A. Wood, D. McEvoy, J. Huntington, C. Morton, M. Anderson, and C. Hain, 2016: The evaporative demand drought index. Part I: Linking drought evolution to variations in evaporative demand. *J. Hydrometeor.*, **17**, 1745–1761, <https://doi.org/10.1175/JHM-D-15-0121.1>.
- Jolly, W. M., M. A. Cochrane, P. H. Freeborn, Z. A. Holden, T. J. Brown, G. J. Williamson, and D. M. J. S. Bowman, 2015: Climate-induced variations in global wildfire danger from 1979 to 2013. *Nat. Commun.*, **6**, 7537, <https://doi.org/10.1038/ncomms8537>.
- Lareau, N. P., N. J. Nauslar, and J. T. Abatzoglou, 2018: The Carr fire vortex: A case of pyrotornadogenesis? *Geophys. Res. Lett.*, **45**, 13 107–13 115, <https://doi.org/10.1029/2018GL080667>.
- McEvoy, D. J., J. L. Huntington, M. T. Hobbins, A. Wood, C. Morton, M. Anderson, and C. Hain, 2016: The evaporative demand drought index. Part II: CONUS-wide assessment against common drought indicators. *J. Hydrometeor.*, **17**, 1763–1779, <https://doi.org/10.1175/JHM-D-15-0122.1>.
- , M. Hobbins, T. J. Brown, K. VanderMolen, T. Wall, J. L. Huntington, and M. Svoboda, 2019: Establishing relationships between drought indices and wildfire danger outputs: A test case for the California–Nevada drought early warning system. *Climate*, **7**, 52, <https://doi.org/10.3390/cli7040052>.
- Nauslar, N. J., J. T. Abatzoglou, and P. T. Marsh, 2018: The 2017 North Bay and Southern California fires: A case study. *Fire*, **1** (1), 18, <https://doi.org/10.3390/fire1010018>.
- Oakley, N. S., F. Cannon, R. Munroe, J. T. Lancaster, D. Gomberg, and F. M. Ralph, 2018: Brief communication: Meteorological and climatological conditions associated with the 9 January 2018 post-fire debris flows in Montecito and Carpinteria, California, USA. *Nat. Hazards Earth Syst. Sci.*, **18**, 3037–3043, <https://doi.org/10.5194/nhess-18-3037-2018>.
- Pyne, S. J., 2016: *California: A Fire Survey*. The University of Arizona Press, 203 pp.
- Radeloff, V. C., and Coauthors, 2018: Rapid growth of the US wildland–urban interface raises wildfire risk. *Proc. Natl. Acad. Sci. USA*, **115**, 3314–3319, <https://doi.org/10.1073/pnas.1718850115>.