The Global Drought and Flood Catalogue

A Complex Relation of Hydrology and Impact

Adapted from "A Global Drought and Flood Catalog from 1950 to 2016," by Xiaogang He, Ming Pan, Zhongwang Wei, Eric F. Wood, and Justin Sheffield. Published online in *BAMS*, May 2020. For the full, citable article, see: [DOI:10.1175/BAMS-D-18-0269.1](https://doi.org/10.1175/BAMS-D-18-0269.1).

To improve assessment of drought and flood risk, we need to improve our understanding of the current risks and how they may change with climate change and human activities. However, current drought and flood risk estimates are often short term and inconsistent, limited to developed nations, and often associated with large uncertainties. Developing a catalogue of hydrological extremes and their characteristics can address these challenges and enable more robust risk quantification.

Drought and flood risk should be quantified in a consistent way to enable a comprehensive understanding of both extremes. Although numerous studies exist on drought and flood risk, most of them treat drought and flood separately. A multivariate risk assessment framework is needed to improve understanding of the combined impact of these extremes.

We aim to develop the first Global Drought and Flood Catalogue (GDFC) based on long-term (1950–2016) meteorological data and improved land surface and river simulations. Development of the GDFC enables the study of the commonalities and differences between these two types of extremes in a comprehensive and systematic way. The GDFC facilitates analysis of worldwide trends in large-scale droughts and floods and therefore provides a key reference for climate change studies, climate model evaluations, and risk analysis.

The GDFC builds on existing models and datasets in several respects. For example, it extends the existing long-term global meteorological dataset [(Princeton Global Forcings (PGF))] from 1950 to near present (2016), with enhanced spatial resolution (0.25°) through statistical downscaling. This dataset drives an

updated version of the Variable Infiltration Capacity (VIC) land surface model to improve soil moisture and runoff variability estimates. The GDFC uses a new Catchment-Based Macro-Scale Floodplain model (CaMa-Flood) to explicitly characterize flood stage and river discharge. Indices of hydrological variables identify drought and pluvial events at the pixel level. A joint spatial-temporal analysis is performed to investigate how droughts and pluvials propagate, merge, or break up. We utilize a severity–area–duration (SAD) clustering algorithm to identify spatially contiguous droughts and pluvials and estimate their time-varying frequency. Finally, characteristics such as frequency, area, and severity are synthesized into the catalogue with a particular focus on

characterizing the long-term variability in risk from both a univariate and (primarily) multi-variate perspective.

Droughts and pluvials around the world

Globally, 453 short-duration (4–6 month) meteorological droughts and 476 pluvials (with a contiguous area larger than 375,000 km²) are documented in the GDFC from 1950 to 2016. There are far fewer medium-duration (7–12 month) events (200), as prolonged hydrological extremes tend to break into short-term events during their evolution.

Based on a three-month standardized precipitation index (SPI3), there has been a statistically significant decrease in the global frequency of

short-term droughts, from more than eight events per year in the 1950s to roughly six events per year in the 1990s; the number then stabilizes afterward. No similar trend has been observed for short-term pluvial frequency, which peaked around eight times per year during the 1980s, followed by a subsequent steep decline.

Continental-scale trends differ. For example, over Africa the robust increasing occurrence of medium-duration drought coincides with the decreased frequency in medium-duration pluvials. This is consistent with previous findings that Africa is moving toward a drier climate.

The SPI3 findings are complemented by the SMPct (soil moisture percentile) analysis: reduced decadal variability and a more synchronous and coherent temporal evolution between droughts and pluvials. Africa shows vulnerability to both dry and wet extremes. We find that, at the global scale, significantly greater frequency of short-duration extremes estimated from SMPct compared to that estimated from SPI3. There are consistencies between SPI3 and SMPct, highlighting that changing precipitation leads to more frequent meteorological extremes, which can translate to agricultural extremes.

For regions lacking such consistency between SPI3 and SMPct, the complexity of hydrologic processes, and how they respond to long-term changes in precipitation, need further investigation. The geographically variable trends may not be consistent or statistically robust, depending on what index is used, highlighting the need to improve our understanding of how prolonged events persist or break up under changing atmospheric conditions and changing hydrological processes.

Droughts and pluvials in Asia generally cover less than 20% of the continent, whereas extremes that occurred in Oceania can cover more than 80% of the total area. As droughts/pluvials become larger, the dependence between severity and area decreases due to increased spatial variability of soil moisture. Furthermore, risks of meteorological and agricultural extremes differ even for events

Timeline of the top five longest drought and pluvial episodes

.....
The GDFC is
on a publicly
accessible data
portal with
web interface
(<http://hydrology.princeton.edu/data/hexg/GDFC/>)
.....

with the same severity, area, and duration, and agricultural extremes have smaller return periods than meteorological extremes given the same magnitude of severity and area.

Frequency analysis thus should consider relationships between multiple characteristics (i.e., severity, area, and duration). It is not necessary for all characteristics of droughts/pluvials to be extreme for the impact to be extreme, but not accounting for the multivariate nature of these extremes can lead to an underestimation of combined impact.

Next challenges

The GDFC can be used in decision-making and impact assessment, and risk reduction, including examining engineering design—for example, evaluation of reservoir storage could guide government's investment decisions to reduce future impacts. To promote actionable science and decision-making, the GDFC needs to be integrated in the form of scalable and policy-relevant catalogue databases accounting for human influences.

Continued efforts are needed to incorporate the dynamic nature of drought and flood risk (e.g., how risks evolve over time) into the current static GDFC. The current GDFC, which only provides deterministic information on drought and flood risk, would benefit from incorporating uncertainties so that risk information can be utilized in strategies for a wide range of scenarios. ••

DISASTERS DON'T
PLAN AHEAD.

YOU CAN.

DON'T WAIT. COMMUNICATE.

Talk to your loved ones about how you are
going to be ready in an emergency.

[VISIT READY.GOV/PLAN.](https://www.ready.gov/plan)