Compound Climate Events and Extremes in the Midlatitudes

Dynamics, Simulation, and Statistical Characterization

Gabriele Messori, Emanuele Bevacqua, Rodrigo Caballero, Dim Coumou, Paolo De Luca, Davide Faranda, Kai Kornhuber, Olivia Martius, Flavio Pons, Colin Raymond, Kunhui Ye, Pascal Yiou, and Jakob Zscheischler

Compound Climate Events and Extremes in the Midlatitudes

What: The workshop, conducted virtually due to travel restrictions related to COVID-19, gathered scientists from six countries and focused on the mechanistic understanding, statistical characterization, and modeling of societally relevant compound climate events and extremes in the midlatitudes. These ranged from co-occurring hot–humid or wet–windy extremes, to spatially compounding wet and dry extremes, to temporally compounding hot–wet events and more. The aim was to bring together selected experts studying a diverse range of compound climate events and extremes to present their ongoing work and outline challenges and future developments in this societally relevant field of research.

When: 7–9 September 2020

Where: Held digitally

KEYWORDS: Atmospheric circulation; Extreme events; Statistical techniques

<https://doi.org/10.1175/BAMS-D-20-0289.1>

Corresponding author: Gabriele Messori, gabriele.messori@geo.uu.se

In final form 9 November 2020

©2021 American Meteorological Society

For information regarding reuse of this content and general copyright information, consult the [AMS Copyright Policy](#).

This article is licensed under a [Creative Commons Attribution 4.0 license](#).

AFFILIATIONS: **Messori**—Department of Earth Sciences, and Centre of Natural Hazards and Disaster Science, Uppsala University, Uppsala, and Department of Meteorology, and Bolin Centre for Climate Research, Stockholm University, Stockholm, Sweden; **Bevacqua**—Department of Meteorology, University of Reading, Reading, United Kingdom; **Caballero**—Department of Meteorology, and Bolin Centre for Climate Research, Stockholm University, Stockholm, Sweden; **Coumou**—Department of Water and Climate Risk, Institute for Environmental Studies (IVM), Vrije Universiteit Amsterdam, Amsterdam, and Royal Netherlands Meteorological Institute (KNMI), De Bilt, Netherlands; **De Luca**—Department of Water and Climate Risk, Institute for Environmental Studies (IVM), Vrije Universiteit Amsterdam, Amsterdam, Netherlands; **Faranda**—Laboratoire des Sciences du Climat et de l’Environnement, UMR8212, CNRS–CEA–UVSQ, Université Paris Saclay, IPSL, Gif-sur-Yvette, and LMD–École Normale Supérieure, Paris, France, and London Mathematical Laboratory, London, United Kingdom; **Kornhuber**—Earth Institute, and Lamont-Doherty Earth Observatory, Columbia University, New York, New York; **Martius**—Institute of Geography, and Oeschger Centre for Climate Change Research, University of Bern, Bern, Switzerland; **Pons and Yiou**—Laboratoire des Sciences du Climat et de l’Environnement, UMR8212, CNRS–CEA–UVSQ, Université Paris Saclay, IPSL, Gif-sur-Yvette, France; **Raymond**—Jet Propulsion Laboratory, California Institute of Technology, Pasadena, California; **Ye**—Department of Earth Sciences, Uppsala University, Uppsala, Sweden; **Zscheischler**—Oeschger Centre for Climate Change Research, and Climate and Environmental Physics, Physics Institute, University of Bern, Bern, Switzerland, and Department of Computational Hydrosystems, Helmholtz Centre for Environmental Research–UFZ, Leipzig, Germany

Compound climate events result from the combination of multiple drivers and/or hazards, which collectively lead to a socioeconomic and/or environmental risk (e.g., Zscheischler et al. 2018). The term *compound extreme* refers to the specific case where several or all of the compounding elements are themselves extreme events. A univariate perspective hampers the process understanding of climate events and extremes, and may ultimately lead to an underestimation of the associated risk (Zscheischler et al. 2020). At the same time, a multivariate perspective brings its own unique challenges, both conceptual and methodological. Multivariate statistical models are typically complex and data intensive (e.g., Bevacqua et al. 2017), yet compound events are by definition less sampled than their univariate counterparts. This is particularly evident for studies on compound extremes, which often rely on very limited datasets. Process studies face analogous challenges, since the components of a compound event may be controlled by one or a multitude of drivers.

The workshop on “Compound climate events and extremes in the midlatitudes” was convened to outline promising ongoing and future research developments in this field and foster discussions on cross-disciplinary methodological approaches. A specific focus was to delineate challenges hindering our understanding of compound events, which may require a concerted community effort to be overcome.

We summarize the workshop’s outcomes around the following themes: drivers of multivariate compound events, spatially or temporally compounding events, and novel cross-disciplinary methodologies for the study of compound events. We conclude by reporting the challenges and future perspectives in the field, issued from roundtable discussions that closed each of the three days of the workshop.

Day 1: Drivers of multivariate compound events

Multivariate compound climate events are the archetype of a compound event, and occur when relevant climate drivers and/or hazards co-occur in the same geographical region.

Rodrigo Caballero and Colin Raymond opened the workshop with a discussion of concurrent hot–humid extremes. These are understood to drive heat stress extremes, often diagnosed

through wet-bulb temperature, with potentially lethal effects on humans (Sherwood and Huber 2010). A comprehensive understanding of hot–humid extremes requires complementary approaches, embracing global-scale statistical analyses and process-oriented studies at the regional-to-local scale. Colin highlighted strong observed trends in combined hot–humid extremes at the global scale, driven by warming air and sea surface temperatures (Raymond et al. 2020). The balance between temperature and moisture contributions to hot–humid extremes results from processes that affect water vapor and atmospheric stability—for example, surface radiative heating, evapotranspiration, and deep convection. However, the lack of detailed and comprehensive datasets at process-relevant time scales (~hourly), as well as regional geographic and meteorological variations, remain outstanding challenges. At the regional scale, a notable hot–humid hotspot is Pakistan’s Indus Valley. Rodrigo explained that events in the region are driven by advection of moist air masses from the Arabian Sea onto land, where they are further heated and moistened through surface fluxes as they pass over the highly irrigated Indus Valley region (Monteiro and Caballero 2019). A comprehensive evaluation of current and future hot–humid extremes thus requires both full-complexity global climate models and high-resolution regional simulations, able to resolve small-scale processes such as land surface evaporation.

The focus next shifted to the role of moist oceanic air masses—and more broadly storm-track activity—in favoring compound coastal flooding from co-occurring extreme precipitation and storm surge. Emanuele Bevacqua analyzed the co-occurrence probability of these two meteorological drivers in the present climate and in a future scenario under high anthropogenic greenhouse gas emissions. Storm-track variability explains to a large extent the spatial variation of the co-occurrence probability. In a future warmer climate, a combination of thermodynamic and dynamic atmospheric changes will likely increase the occurrence of compound flooding, particularly in the midlatitudes (Bevacqua et al. 2020a,b). These findings indicate that the detected changes should be considered in future risk assessments, as they may aggravate the hazard caused by mean sea level rise.

Storm-track variability also modulates other midlatitude compound events, in both summer and winter. Summer can see persistent hot–dry or cold–wet weather, with significant impacts on agriculture, health, the energy sector, and the environment. In Europe, such summer weather anomalies are associated with specific states of jet stream, i.e., respectively, the dominance of blocked flows or a persistent zonal jet. Dim Coumou presented a dynamical systems perspective on the topic, which supports Lorenz’s hypothesis that summer climate is largely intransitive (Lorenz 1990) and suggests that persistent blocked or zonal states may be characterized by long memory (~40 days) and governed by different attractors. If intransitivity were indeed a fundamental property of summer atmospheric dynamics, this would have major implications for predictability on meteorological to interannual time scales. However, there are still critical gaps in our knowledge of the dynamics underlying jet-stream variability in the summer season. This is a pressing issue, as the summer season is when the bulk of the agricultural production takes place, and thus present and future climate risks are likely most severe in this season.

In winter, storm-track variability modulates concurrent wet–windy and hot–windy extremes in Europe (e.g., Messori et al. 2019; De Luca et al. 2020a). Gabriele Messori discussed the causal chain leading to these extremes. The simultaneous occurrence of anticyclonic planetary wave breaking to the south of the North Atlantic jet stream and cyclonic wave breaking to the north of the jet stream leads to a very zonal and intense large-scale flow. This, in turn, favors a heightened frequency of explosive cyclones in the Atlantic basin and destructive windstorms over western and continental Europe (Messori and Caballero 2015). A zonal, intense jet also leads to heavy precipitation, associated with the Atlantic cyclones, and favors the penetration of warm, moist airmasses deep into the European continent. The

above process chain is recovered when studying multidecadal or longer variability in long climate model integrations (Messori et al. 2019). This has important implications for the interpretation of dynamical changes in climate projections and their relevance for high-impact compound climate extremes in Europe.

Day 2: Spatially or temporally compounding events

Spatially compounding climate events occur when multiple locations are affected by climate hazards within a short time window. Temporally compounding events result from a temporal sequence of hazards at a given geographical location (Zscheischler et al. 2020).

Kai Kornhuber opened the day with an analysis of concurrent summertime heatwaves across different Northern Hemisphere breadbasket regions. This co-occurrence can be ascribed to amplified planetary waves, which induce large meanders in the jet stream in turn favoring the spatial compounding of the heatwaves. Specifically, planetary waves with zonal wavenumbers 5 and 7 are recurrent in summer, and have a favored phase position, making them very effective drivers of synchronized regional heatwaves (Kornhuber et al. 2020). While no scientific consensus on future changes in these wave patterns has been established so far, their impacts are expected to become more severe due to thermodynamic factors alone, possibly enhancing crop production volatility and imperiling global food security.

Food security was also one of the motivations underlying Paolo De Luca's study of co-occurring wet and dry extreme events on a global scale (De Luca et al. 2020b). Paolo highlighted periods in the observational record when large parts of the global land surface were simultaneously affected by flooding and drought, and then proceeded to evaluate the correlations between these extremes and leading modes of climate variability. El Niño–Southern Oscillation, the Pacific decadal oscillation, and the Atlantic multidecadal oscillation show regionally significant correlations with wet and dry extremes, highlighting how their interplay may lead to extensive co-occurrence episodes. This result puts the spotlight on global multihazard maps as a key tool to evaluate worst-case hydrological scenarios.

Large-scale variability patterns were also highlighted by Kunhui Ye as key to the spatial compounding of extremes. Kunhui discussed the large-scale dynamics underlying the warm Arctic–cold Eurasia (WACE) pattern observed in recent decades, and highlighted two winter-time circulation modes of variability. Decadal trends in these modes explain a large part of the observed WACE pattern and additionally lead to a heightened frequency of temperature extremes over northern Europe and northern Eurasia (Ye and Messori 2020).

The focus next shifted to temporally compounding events. Olivia Martius revisited several methods that have been proposed to characterize and quantify temporal (serial) clustering—including the index of dispersion, Ripley's K , and the Cox regression—with a specific focus on the serial clustering of heavy precipitation. The latter has implications on seasonal-to-yearly time scales for insurance contracts and the broader economy (Priestley et al. 2018), and on subseasonal time scales for flooding (Barton et al. 2016). Olivia highlighted how most of the available statistical approaches quantify the significance of serial clustering and allow to include covariates, but are unable to identify specific clustering episodes and require a predefined event set.

Day 3: Cross-disciplinary methodologies for the study of compound events

The final day of the workshop was dedicated to methodological innovations for the study of compound climate events and extremes, with a focus on cross-pollination from fields beyond climate science.

Flavio Pons opened the day by presenting a novel bias correction approach for snowfall in climate models. Accurate estimation of snowfall is crucial to correctly describing compound hydrometeorological events in winter, such as rain-on-snow episodes, yet climate models

often display large biases in this respect (Frei et al. 2018). Flavio argued that, by combining breakpoint search algorithms with cubic spline logit-linear regression outputs, it is possible to improve on past bias-correction methods for snowfall by relying only on temperature and precipitation data. The method provided encouraging results when applied to the ERA5 data (Hersbach et al. 2020) and to a high-resolution numerical climate simulation over Europe, and offers a feasible approach to reconstruct snowfall without requiring multivariate or conditional bias correction, nor stochastic generation of unobserved events.

The issue of model biases, and of how to evaluate the ability of climate models to simulate compound events, is indeed crucial to gaining a robust understanding of the latter. Compound extremes are particularly challenging in this respect, since many commonly used model evaluation metrics for multivariate distributions, such as correlation coefficients, are not suitable for isolating extreme occurrences. Jakob Zscheischler proposed a new metric, based on the Kullback–Leibler divergence concept from information theory, that measures whether the tails of different bivariate distributions show a similar dependence structure. Such metric was then used to compare compound precipitation and wind extremes in the Alpine region across different datasets (Zscheischler et al. 2021). Boundary conditions appear to be a key factor in correctly modeling compound extremes, while external forcings are a second-order effect. The proposed metric allows to evaluate climate model simulations with respect to compound extremes, and to outline key requirements for future model development.

The difficulty of modeling compound extremes, and more broadly low-probability events, was also the topic of Pascal Yiou's presentation. Pascal presented a methodology combining importance sampling—often used in statistical mechanics—with a stochastic weather generator based on circulation analogs. This allows to simulate physically plausible, but unprecedented, long-lasting events such as warm or wet seasons (Yiou and Jézéquel 2020). In this context, analogs are days displaying similar large-scale atmospheric states over a chosen geographical domain. The methodology was then applied to simulate temporally compounding warm winters and wet springs in France. These events have a major detrimental effect on French wheat production: in 2016, a record-low yield was reported due to warm winter temperatures followed by heavy precipitation in May. The stochastic importance sampling tool shows that the present-day climatic conditions in principle allow for a compound warm winter–wet spring event even more devastating than that of 2016 (Pfleiderer et al. 2021). This type of approach thus allows investigating the properties of unprecedented events linked to the large-scale atmospheric circulation.

Closely related to the study of unprecedented compound events is the question of how anthropogenic emissions may impact hazardous climate events. This is often studied under the assumption that the atmospheric circulation associated to such events is not itself affected by climate change. However, the compound events paradigm demands for new methodologies in this respect. Davide Faranda presented an approach grounded in dynamical systems theory, that allows diagnosing the role of the atmospheric circulation during hazardous extreme events (Faranda et al. 2017). The approach is based on embedding the circulation patterns observed during selected extremes into historical climate simulations and future projections. When applied to the latter, it highlights major changes in the probability, predictability and persistence of large-scale atmospheric patterns leading to hazardous extremes such as European heatwaves and cold spells (Faranda et al. 2020). These results highlight that dynamical changes in the atmosphere must be taken into account when performing attribution studies for compound events.

Future perspectives and challenges in the field

The results presented during the workshop highlighted the breadth of climate drivers and hazards falling under the umbrella term *compound events*, even when focusing on the

midlatitudes. At the same time, the presentations underscored some pivotal ideas—both theoretical and methodological—that may be leveraged as unifying paradigms for advancing our understanding of the topic. Recurrent planetary-scale atmospheric anomalies, such as those associated with planetary wave patterns and the closely related storm-track variability, provide an effective lens for interpreting different categories of compound events, from multivariate extremes such as wet and windy episodes (e.g., De Luca et al. 2020a) to spatially compounding heatwaves (e.g., Kornhuber et al. 2019, 2020) to compound long-lasting anomalies (e.g., Messori et al. 2019). Persistence was also highlighted as an overarching concept relevant to a diverse set of compound events. Often, multivariate or spatially compounding extremes are associated with unusually persistent large-scale atmospheric anomalies. Similarly, persistent structures can be key for driving temporally compounding events. The workshop also pointed to the added value of cross-pollination with tools drawn from other fields of science, such as dynamical systems theory, statistical physics, or information theory. These tools provide, for example, a means of simulating unprecedented compound events (Yiou and Jézéquel 2020) or evaluating the ability of climate models to reproduce compound extremes (Zscheischler et al. 2021). They also provide a framework to diagnose the role of changes in atmospheric dynamics in modulating the frequency of occurrence and physical properties of hazardous extremes in future climates (Faranda et al. 2020). These advances do not preclude the relevance of more conventional analyses based on atmospheric dynamics, and indeed partly rely on the latter through, for example, the calculation of circulation analogs for compound events.

During the daily roundtable discussions, the workshop participants also distilled some outstanding challenges faced by the compound climate events community. These pertain to both the seamless integration of the different perspectives discussed above and longer-standing interpretational issues in atmospheric dynamics. A first, fundamental challenge is to develop a clear terminology to distinguish between different compound events and extremes. A decisive step in this direction was recently made with the publication of a compound events typology (Zscheischler et al. 2020), but a continued collective effort by the community is needed in this respect.

A similar problem extends to the integration of perspectives issued from different fields of science, where the same term may be used with different meanings. *Persistence* is emblematic in this respect: on the one hand, it was identified as a concept of relevance to a broad range of compound events; on the other hand, discussions on persistence highlighted how the meaning of the term is highly context specific. In dynamical systems theory, persistence relates to how long the system being studied resides in a given neighborhood in an appropriately defined phase space (e.g., Faranda et al. 2017). In the atmospheric sciences, persistence is used very broadly: from the number of consecutive days the atmosphere spends within a given cluster of large-scale configurations (often referred to as *weather regimes*), to the duration of propagating structures, such as planetary wave patterns. Each definition has advantages and shortcomings, and reflects different aspects of the evolution of a given system. A concerted effort is required to compare and reconcile the different viewpoints and identify those most relevant to specific classes of compound events.

Other challenges reflect long-standing open questions within the atmospheric sciences. The continuum of spatial and temporal scales within the climate system complicates separating the drivers of specific compound events. For example, the interplay between planetary waves and storm-track variability often results in a chicken-and-egg problem of what drives what. Similarly, energy transfers across scales—resulting in direct and inverse turbulent energy cascades—are key for triggering atmospheric waves, yet they can be complex to diagnose and quantify (e.g., Faranda et al. 2018). These challenges feed into the broader difficulty of understanding the role of atmospheric dynamics in climate change (Shepherd 2014), and particularly in modulating the frequency and nature of future compound events.

Finally, a challenge unique to this year—albeit common to all fields of research—was the need to hold a virtual meeting. On the one hand, this enabled the participation of colleagues who may have been unable to travel to a physical meeting even under normal circumstances. On the other hand, it required a careful tailoring of the workshop's format to enable effective discussions and brainstorming. For example, a deliberate choice was made to reduce the audience to a minimum, and to shorten the time allotted to individual presentations. While the informal and often extremely useful “coffee-break interactions” are difficult to reproduce in a virtual setting, the authors of this report were overall satisfied with the ease and quality of the virtual discussions.

In summary, the workshop's outcomes motivate a cautious optimism and the view that neither the scientific nor the logistic challenges facing research on compound events are unsurmountable. For example, dynamical systems theory and statistical physics can help to diagnose how future changes in atmospheric dynamics may affect compound events. Similarly, a joint application of different concepts of persistence to temporally compounding precipitation extremes was discussed during the workshop, and no a priori major theoretical or technical hindrances were identified. Ultimately, the workshop helped to identify and describe key challenges in the study of compound climate events and extremes, which is an important first step toward developing concerted initiatives to tackle them.

Acknowledgments. The workshop was convened by G. Messori, and was held virtually due to the COVID-19 pandemic. However, the Stockholm International Meteorological Institute had made funding available should a physical meeting have taken place.

References

- Barton, Y., P. Giannakaki, H. von Waldow, C. Chevalier, S. Pfahl, and O. Martius, 2016: Clustering of regional-scale extreme precipitation events in southern Switzerland. *Mon. Wea. Rev.*, **144**, 347–369, <https://doi.org/10.1175/MWR-D-15-0205.1>.
- Bevacqua, E., D. Maraun, I. Hobæk Haff, M. Widmann, and M. Vrac, 2017: Multivariate statistical modelling of compound events via pair-copula constructions: Analysis of floods in Ravenna (Italy). *Hydrol. Earth Syst. Sci.*, **21**, 2701–2723, <https://doi.org/10.5194/hess-21-2701-2017>.
- , M. I. Voudoukas, G. Zappa, K. Hodges, T. G. Shepherd, D. Maraun, L. Mentaschi, and L. Feyen, 2020a: More meteorological events that drive compound coastal flooding are projected under climate change. *Commun. Earth Environ.*, **1**, 47, <https://doi.org/10.1038/s43247-020-00044-z>.
- , T. G. Shepherd, and M. Vrac, 2020b: Brief communication: The role of using precipitation or river discharge data when assessing global coastal compound flooding. *Nat. Hazards Earth Syst. Sci.*, **20**, 1765–1782, <https://doi.org/10.5194/nhess-20-1765-2020>.
- De Luca, P., G. Messori, F. M. E. Pons, and D. Faranda, 2020a: Dynamical systems theory sheds new light on compound climate extremes in Europe and eastern North America. *Quart. J. Roy. Meteor. Soc.*, **146**, 1636–1650, <https://doi.org/10.1002/qj.3757>.
- , R. L. Wilby, M. Mazzoleni, and G. Di Baldassarre, 2020b: Concurrent wet and dry hydrological extremes at the global scale. *Earth Syst. Dyn.*, **11**, 251–266, <https://doi.org/10.5194/esd-11-251-2020>.
- Faranda, D., G. Messori, and P. Yiou, 2017: Dynamical proxies of North Atlantic predictability and extremes. *Sci. Rep.*, **7**, 41278, <https://doi.org/10.1038/srep41278>.
- , V. Lembo, M. Iyer, D. Kuzay, S. Chibbaro, F. Daviaud, and B. Dubrulle, 2018: Computation and characterization of local subfilter-scale energy transfers in atmospheric flows. *J. Atmos. Sci.*, **75**, 2175–2186, <https://doi.org/10.1175/JAS-D-17-0114.1>.
- , M. Vrac, P. Yiou, A. Jézéquel, and S. Thao, 2020: Changes in future synoptic circulation patterns: Consequences for extreme event attribution. *Geophys. Res. Lett.*, **47**, e2020GL088002, <https://doi.org/10.1029/2020GL088002>.
- Frei, P., S. Kotlarski, M. A. Liniger, and C. Schar, 2018: Future snowfall in the Alps: Projections based on the EURO-CORDEX regional climate models. *Cryosphere*, **12**, 1–24, <https://doi.org/10.5194/tc-12-1-2018>.
- Hersbach, H., and Coauthors, 2020: The ERA5 global reanalysis. *Quart. J. Roy. Meteor. Soc.*, **146**, 1999–2049, <https://doi.org/10.1002/qj.3803>.
- Kornhuber, K., S. Osprey, D. Coumou, S. Petri, V. Petoukhov, S. Rahmstorf, and L. Gray, 2019: Extreme weather events in early summer 2018 connected by a recurrent hemispheric wave-7 pattern. *Environ. Res. Lett.*, **14**, 054002, <https://doi.org/10.1088/1748-9326/ab13bf>.
- , D. Coumou, E. Vogel, C. Lesk, J. F. Donges, J. Lehmann, and R. M. Horton, 2020: Amplified Rossby waves enhance risk of concurrent heatwaves in major breadbasket regions. *Nat. Climate Change*, **10**, 48–53, <https://doi.org/10.1038/s41558-019-0637-z>.
- Lorenz, E. N., 1990: Can chaos and intransitivity lead to interannual variability? *Tellus*, **42A**, 378–389, <https://doi.org/10.3402/tellusa.v42i3.11884>.
- Messori, G., and R. Caballero, 2015: On double Rossby wave breaking in the North Atlantic. *J. Geophys. Res. Atmos.*, **120**, 11 129–11 150, <https://doi.org/10.1002/2015JD023854>.
- , P. Davini, M. C. Alvarez-Castro, F. S. R. Pausata, P. Yiou, and R. Caballero, 2019: On the low-frequency variability of wintertime Euro-Atlantic planetary wave-breaking. *Climate Dyn.*, **52**, 2431–2450, <https://doi.org/10.1007/s00382-018-4373-2>.
- Monteiro, J. M., and R. Caballero, 2019: Characterization of extreme wet-bulb temperature events in southern Pakistan. *Geophys. Res. Lett.*, **46**, 10 659–10 668, <https://doi.org/10.1029/2019GL084711>.
- Pfleiderer, P., A. Jézéquel, J. Legrand, N. Legrix, I. Markantonis, E. Vignotto, and P. Yiou, 2021: Simulating compound weather extremes responsible for critical crop failure with stochastic weather generators. *Earth Syst. Dyn.*, **12**, 103–120, <https://doi.org/10.5194/esd-12-103-2021>.
- Priestley, M. D. K., H. F. Dacre, L. C. Shaffrey, K. I. Hodges, and J. G. Pinto, 2018: The role of serial European windstorm clustering for extreme seasonal losses as determined from multi-centennial simulations of high resolution global climate model data. *Nat. Hazards Earth Syst. Sci.*, **18**, 2991–3006, <https://doi.org/10.5194/nhess-18-2991-2018>.
- Raymond, C., T. Matthews, and R. M. Horton, 2020: The emergence of heat and humidity too severe for human tolerance. *Sci. Adv.*, **6**, eaaw1838, <https://doi.org/10.1126/sciadv.aaw1838>.
- Shepherd, T. G., 2014: Atmospheric circulation as a source of uncertainty in climate change projections. *Nat. Geosci.*, **7**, 703–708, <https://doi.org/10.1038/ngeo2253>.
- Sherwood, S. C., and M. Huber, 2010: An adaptability limit to climate change due to heat stress. *Proc. Natl. Acad. Sci. USA*, **107**, 9552–9555, <https://doi.org/10.1073/pnas.0913352107>.
- Ye, K., and G. Messori, 2020: Two leading modes of wintertime atmospheric circulation drive the recent warm Arctic–cold Eurasia temperature pattern. *J. Climate*, **33**, 5565–5587, <https://doi.org/10.1175/JCLI-D-19-0403.1>.
- Yiou, P., and A. Jézéquel, 2020: Simulation of extreme heat waves with empirical importance sampling. *Geosci. Model Dev.*, **13**, 763–781, <https://doi.org/10.5194/gmd-13-763-2020>.
- Zscheischler, J., and Coauthors, 2018: Future climate risk from compound events. *Nat. Climate Change*, **8**, 469–477, <https://doi.org/10.1038/s41558-018-0156-3>.
- , and Coauthors, 2020: A typology of compound weather and climate events. *Nat. Rev. Earth Environ.*, **1**, 333–347, <https://doi.org/10.1038/s43017-020-0060-z>.
- , P. Naveau, O. Martius, S. Engelke, and C. C. Raible, 2021: Evaluating the dependence structure of compound precipitation and wind speed extremes. *Earth Syst. Dyn.*, **12**, 1–16, <https://doi.org/10.5194/esd-12-1-2021>.