A photograph of a snowy park path, likely in Central Park, New York. The path is covered in a thick layer of snow, with footprints visible. On either side of the path are ornate blue and gold lampposts. In the background, a large brick building is visible. The sky is overcast and it appears to be snowing.

EXPLAINING EXTREME EVENTS OF 2014

From A Climate Perspective

Special Supplement to the
Bulletin of the American Meteorological Society
Vol. 96, No. 12, December 2015

EXPLAINING EXTREME EVENTS OF 2014 FROM A CLIMATE PERSPECTIVE

Editors

Stephanie C. Herring, Martin P. Hoerling, James P. Kossin, Thomas C. Peterson, and Peter A. Stott

Special Supplement to the

Bulletin of the American Meteorological Society

Vol. 96, No. 12, December 2015

AMERICAN METEOROLOGICAL SOCIETY

CORRESPONDING EDITOR:

Stephanie C. Herring, PhD
NOAA National Centers for Environmental Information
325 Broadway, E/CC23, Rm 1B-131
Boulder, CO 80305-3328
E-mail: stephanie.herring@noaa.gov

COVER CREDITS:

FRONT: ©iStockphotos.com/coleong—Winter snow, Boston, Massachusetts, United States.

BACK: ©iStockphotos.com/nathanphoto—Legget, California, United States – August 13, 2014: CAL FIRE helicopter surveys a part of the Lodge Fire, Mendocino County.

HOW TO CITE THIS DOCUMENT

Citing the complete report:

Herring, S. C., M. P. Hoerling, J. P. Kossin, T. C. Peterson, and P. A. Stott, Eds., 2015: Explaining Extreme Events of 2014 from a Climate Perspective. *Bull. Amer. Meteor. Soc.*, **96** (12), S1–S172.

Citing a section (example):

Yoon, J. H., S.-Y. S. Wang, R. R. Gillies, L. Hipps, B. Kravitz, and P. J. Rasch, 2015: Extreme fire season in California: A glimpse into the future? [in “Explaining Extremes of 2014 from a Climate Perspective”]. *Bull. Amer. Meteor. Soc.*, **96** (12), S5–S9.

EDITORIAL AND PRODUCTION TEAM

Riddle, Deborah B., Lead Graphics Production, NOAA/NESDIS National Centers for Environmental Information, Asheville, NC

Love-Brotak, S. Elizabeth, Graphics Support, NOAA/NESDIS National Centers for Environmental Information, Asheville, NC

Veasey, Sara W., Visual Communications Team Lead, NOAA/NESDIS National Centers for Environmental Information, Asheville, NC

Griffin, Jessica, Graphics Support, Cooperative Institute for Climate and Satellites-NC, North Carolina State University, Asheville, NC

Maycock, Tom, Editorial Support, Cooperative Institute for Climate and Satellites-NC, North Carolina State University, Asheville, NC

Misch, Deborah J., Graphics Support, LMI Consulting, Inc., NOAA/NESDIS National Centers for Environmental Information, Asheville, NC

Osborne, Susan, Editorial Support, LMI Consulting, Inc., NOAA/NESDIS National Centers for Environmental Information, Asheville, NC

Schreck, Carl, Editorial Support, Cooperative Institute for Climate and Satellites-NC, North Carolina State University, and NOAA/NESDIS National Centers for Environmental Information, Asheville, NC

Sprain, Mara, Editorial Support, LAC Group, NOAA/NESDIS National Centers for Environmental Information, Asheville, NC

Young, Teresa, Graphics Support, STG, Inc., NOAA/NESDIS National Centers for Environmental Information, Asheville, NC

TABLE OF CONTENTS

Abstract.....	ii
1. Introduction to Explaining Extreme Events of 2014 from a Climate Perspective	1
2. Extreme Fire Season in California: A Glimpse Into the Future?	5
3. How Unusual was the Cold Winter of 2013/14 in the Upper Midwest?.....	10
4. Was the Cold Eastern Us Winter of 2014 Due to Increased Variability?	15
5. The 2014 Extreme Flood on the Southeastern Canadian Prairies	20
6. Extreme North America Winter Storm Season of 2013/14: Roles of Radiative Forcing and the Global Warming Hiatus.....	25
7. Was the Extreme Storm Season in Winter 2013/14 Over the North Atlantic and the United Kingdom Triggered by Changes in the West Pacific Warm Pool?	29
8. Factors Other Than Climate Change, Main Drivers of 2014/15 Water Shortage in Southeast Brazil.....	35
9. Causal Influence of Anthropogenic Forcings on the Argentinian Heat Wave of December 2013	41
10. Extreme Rainfall in the United Kingdom During Winter 2013/14: The Role of Atmospheric Circulation and Climate Change.....	46
11. Hurricane Gonzalo and its Extratropical Transition to a Strong European Storm.....	51
12. Extreme Fall 2014 Precipitation in the Cévennes Mountains	56
13. Record Annual Mean Warmth Over Europe, the Northeast Pacific, and the Northwest Atlantic During 2014: Assessment of Anthropogenic Influence.....	61
14. The Contribution of Human-Induced Climate Change to the Drought of 2014 in the Southern Levant Region.....	66
15. Drought in the Middle East and Central–Southwest Asia During Winter 2013/14.....	71
16. Assessing the Contributions of East African and West Pacific Warming to the 2014 Boreal Spring East African Drought	77
17. The 2014 Drought in the Horn of Africa: Attribution of Meteorological Drivers.....	83
18. The Deadly Himalayan Snowstorm of October 2014: Synoptic Conditions and Associated Trends	89
19. Anthropogenic Influence on the 2014 Record-Hot Spring in Korea	95
20. Human Contribution to the 2014 Record High Sea Surface Temperatures Over the Western Tropical And Northeast Pacific Ocean	100
21. The 2014 Hot, Dry Summer in Northeast Asia.....	105
22. Role of Anthropogenic Forcing in 2014 Hot Spring in Northern China	111
23. Investigating the Influence of Anthropogenic Forcing and Natural Variability on the 2014 Hawaiian Hurricane Season.	115
24. Anomalous Tropical Cyclone Activity in the Western North Pacific in August 2014	120
25. The 2014 Record Dry Spell at Singapore: An Intertropical Convergence Zone (ITCZ) Drought.....	126
26. Trends in High-Daily Precipitation Events in Jakarta and the Flooding of January 2014	131
27. Extreme Rainfall in Early July 2014 in Northland, New Zealand—Was There an Anthropogenic Influence?	136
28. Increased Likelihood of Brisbane, Australia, G20 Heat Event Due to Anthropogenic Climate Change	141
29. The Contribution of Anthropogenic Forcing to the Adelaide and Melbourne, Australia, Heat Waves of January 2014	145
30. Contributors to the Record High Temperatures Across Australia in Late Spring 2014	149
31. Increased Risk of the 2014 Australian May Heatwave Due to Anthropogenic Activity.....	154
32. Attribution of Exceptional Mean Sea Level Pressure Anomalies South of Australia in August 2014	158
33. The 2014 High Record of Antarctic Sea Ice Extent.....	163
34. Summary and Broader Context.....	168

Understanding how long-term global change affects the intensity and likelihood of extreme weather events is a frontier science challenge. This fourth edition of explaining extreme events of the previous year (2014) from a climate perspective is the most extensive yet with 33 different research groups exploring the causes of 29 different events that occurred in 2014. A number of this year's studies indicate that human-caused climate change greatly increased the likelihood and intensity for extreme heat waves in 2014 over various regions. For other types of extreme events, such as droughts, heavy rains, and winter storms, a climate change influence was found in some instances and not in others. This year's report also included many different types of extreme events. The tropical cyclones that impacted Hawaii were made more likely due to human-caused climate change. Climate change also decreased the Antarctic sea ice extent in 2014 and increased the strength and likelihood of high sea surface temperatures in both the Atlantic and Pacific Oceans. For western U.S. wildfires, no link to the individual events in 2014 could be detected, but the overall probability of western U.S. wildfires has increased due to human impacts on the climate.

Challenges that attribution assessments face include the often limited observational record and inability of models to reproduce some extreme events well. In general, when attribution assessments fail to find anthropogenic signals this alone does not prove anthropogenic climate change did not influence the event. The failure to find a human fingerprint could be due to insufficient data or poor models and not the absence of anthropogenic effects.

This year researchers also considered other human-caused drivers of extreme events beyond the usual radiative drivers. For example, flooding in the Canadian prairies was found to be more likely because of human land-use changes that affect drainage mechanisms. Similarly, the Jakarta floods may have been compounded by land-use change via urban development and associated land subsidence. These types of mechanical factors re-emphasize the various pathways beyond climate change by which human activity can increase regional risk of extreme events.

14. THE CONTRIBUTION OF HUMAN-INDUCED CLIMATE CHANGE TO THE DROUGHT OF 2014 IN THE SOUTHERN LEVANT REGION

K. BERGAOUI, D. MITCHELL, R. ZAABOUL, R. McDONNELL, F. OTTO, AND M. ALLEN

A combined modeling and observational study suggests that the persistent rainfall deficit during the 2014 rainy season in southern Levant was made more likely due to anthropogenic climate change.

Introduction. While the extent to which the 2007/08 drought in the Levant region destabilized the Syrian government continues to be debated, there is no questioning the enormous toll this extreme event took on the region's population. The movement of refugees from both the drought and war affected regions into Jordan and Lebanon ensured that the anomalously low precipitation in the winter of 2013/14 amplified impacts on already complex water and food provisions.

It is hypothesized that droughts over the Levant region can be linked to three types of synoptic regimes (Saaroni et al. 2014). The regimes include 1) an expansion of the subtropical high over the majority of the Mediterranean Basin, 2) a pronounced stagnant ridge/block, and 3) an intrusion of lower-level continental polar air. The 2014 drought, affecting parts of Jordan, Lebanon, Palestine, and Israel, was characterized by extremes in low rainfall, the extent of the long dry periods, and three exceptional rainfall events that interspersed these. The drought itself was thought to be due to a large-scale winter blocking event that prevented weather systems from reaching the region (Udasin 2014).

The motivation for this study is twofold: 1) due to the instability in the Levant region, stresses such as meteorological extremes can often lead to enhanced

public and political tensions. 2) There is a clear bias in the literature for looking at extreme events over industrialized countries (e.g., Herring et al. 2014)—this study provides an example in an alternative geographic region.

Data and Methods. In studying the 2014 drought, one of the challenges to overcome is accessing data in this extremely politically charged region. As such, we only concentrate on precipitation changes here and discuss their relevance for wider impacts.

A spatially and temporally coherent set of station precipitation data is available from the Israeli Meteorological Service (<http://data.gov.il/ims>), and it follows the World Meteorological Organization quality control guidelines. The data span the period 1980–2014. Additional data for the Levant region was obtained from the Jordanian Ministry of Water and Irrigation and the Palestinian Water Authority. All station locations are plotted on Fig. 14.1a, and these stations provide the longest reliable record available for this region. In addition, we use the gridded E-OBS dataset (Haylock et al. 2008) to put the analysis in the context of longer-term climate variability.

Often direct observational inferences of extreme events are highly uncertain due to small sample sizes (e.g., Sipple et al. 2015). This is especially true in the Levant region. One well-established technique to circumvent this issue is using climate models to run many thousands of simulations of “possible” climate, thereby sampling internal climate variability. Here, we make use of the weather@home project (Massey et al. 2015), which allows us to run many thousands of simulations for 2014 over the region of interest. We use the atmosphere-only model, HadAM3P, to drive a regional model (HadRM3P) that simulates climate for mid-northern Africa and the southern Levant

AFFILIATIONS: BERGAOUI—International Centre for Biosaline Agriculture, Dubai, United Arab Emirates, and Institut National de la Météorologie, Tunis, Tunisia; MITCHELL, OTTO, AND ALLEN—Environmental Change Institute, Oxford University, Oxford, United Kingdom; ZAABOUL—International Centre for Biosaline Agriculture, Dubai, United Arab Emirates; McDONNELL—International Centre for Biosaline Agriculture, Dubai, United Arab Emirates, and Environmental Change Institute, Oxford University, Oxford, United Kingdom

DOI:10.1175/BAMS-D-15-00129.1

A supplement to this article is available online (10.1175/BAMS-D-15-00129.2)

FIG. 14.1. (a) A map of the southern Levant region, showing the topography, political boundaries, and station locations. The region is bounded by 46°N–14°S and 25°W–63°E. **(b)** The cumulative rainy season (JF) precipitation totals for each year of the observational record averaged over the whole domain. The dots show stations, and the blue line shows E-OBS. The correlation coefficient between these datasets is ~ 0.94 . **(c)** The monthly observed climatology (black line) of domain-averaged precipitation from 1980 to 2014. The 2013/14 values are plotted as black circles. Box-and-whisker plots show the median, interquartile range, and range for the biased corrected model data for actual conditions (blue) and natural conditions (red). **(d)** Return level plots of cumulative precipitation for Jan and Feb during the rainy season for natural conditions (red) and actual conditions (blue). Gray line shows the 65-year return level. Individual natural SST patterns are shown with thin pink lines, and the pooled SST pattern is shown with a thick orange line.

at 50-km resolution. The details are the same as the European model described in Massey et al. 2015.

To perform the 2014 event attribution, we simulate two possible climates under 2014 conditions: 1) an experiment with all known natural and anthropogenic climate forcings (“actual conditions”) and 2) an experiment with only natural climate forcings (“natural conditions”). All anthropogenic and natural conditions follow the recommendations outlined by the Intergovernmental Panel on Climate Change (IPCC; Solomon et al. 2007). For the actual conditions experiment, the sea surface temperatures (SSTs) are taken from Operational Sea Surface Temperature and

Sea Ice Analysis (OSTIA) observations (Stark et al. 2007). For the natural conditions experiment, the anthropogenic signal is taken out of the OSTIA SST patterns to leave only the naturalized SST pattern. The anthropogenic fingerprint is calculated by differencing the Coupled Model Intercomparison Project Phase 5 (CMIP5) “historical” and “historicalNat” simulations from 11 GCMs that performed these experiments and then subtracting this change in SST pattern from the OSTIA SSTs (see Schaller et al. 2014 for details). The use of 11 different models gives us 1 potential naturalized SST pattern.

The observed perspective. Figure 14.1b shows the extreme persistence of this two-month precipitation deficit event is unprecedented in the observational record. The cumulative JF precipitation for 2014 is clearly an outlier over the 1950–present period. Figure 14.1c shows the monthly climatology of observed precipitation along with the specific 2013/14 observed values. In December, the precipitation was in the upper quartile of the observed range, and

most of that was due to a 10-day extreme downpour midway into the month (not shown). Following that wet extreme event, both January and February had the lowest rainfall since observations began. While the wet and dry events balance out in the seasonal average, the extreme persistence of this two-month precipitation deficit event is unprecedented.

The modeled perspective. To understand how anthropogenic climate change increased the likelihood of such a persistent drought, we use climate simulations to sample the internal climate variability during the rainy season of 2013/14. A well-known issue

is the large dry bias in precipitation over this region, which may be due to any combination of 1) the varied land cover/topography and relationship to convection in the region, 2) the representation of Mediterranean storm tracks in models (Anagnostopoulou et al. 2006), or 3) capturing upslope flow of moist air masses (Black 2009). In general, models have trouble capturing all the complex factors contributing to precipitation in this region. Here, we make the plausible assumption that the statistics of observed precipitation are quasi stationary over the 30-year period, and we use this as a climatology to evaluate the regional model. Comparing the cumulative distribution functions (CDFs) of the observations and the model simulations, it is also found that there is a dry bias in the model. In Supplementary Fig. S14.1, we show the CDFs for grid boxes with the lowest and highest biases, respectively. The bias was found to be linearly changing, with no significant difference between the “actual” and “natural” simulations, suggesting that it was not a result of misrepresentation of large-scale climate modes. To correct the bias, we use a nonparametric quantile mapping technique, matching estimated empirical quantiles between the observations and the “actual conditions” model scenario (Boe et al. 2007). The same bias correction is then applied to the natural model scenario. Figure 14.1c shows the bias corrected actual conditions and natural conditions model data plotted as box-and-whisker plots. There is a tendency for the actual conditions scenario to have less extreme precipitation with a persistently drier mean state (at least for January–March). Return level plots of these data averaged over the whole domain are shown in Fig. 14.1d, and the natural forcings scenario shows

wetter conditions than the actual conditions throughout.

Figure 14.2a expands on this by showing the region averaged, January–February averaged, and 65-year return levels for the actual scenario and natural scenarios based on the individual estimates of naturalized SSTs. In general, meteorological drought conditions are more frequent in the actual conditions scenario, and this holds true for 9 out of 11 of the estimated naturalized experiments. For instance, the average return level in the natural scenarios is estimated to have around ~10 mm more precipitation than the actual conditions scenario.

To understand these differences in more detail, we calculate the Fraction of Attributable Risk (FAR; Allen 2003), defined as $FAR = 1 - (p_{nat} / p_{act})$, where p_{nat} is the probability of a rainfall deficit as low as or lower than the observed 2014 January–February

FIG. 14.2. (a) 65-year return levels over the entire region for the actual conditions (blue) and the individual estimates of natural conditions (red). (b) As in (a), but for estimates of the FAR. The uncertainty estimates are calculated by randomly resampling (with replacement) the underlying distribution 10 000 times. The bars show the median, interquartile range, and range of uncertainty.

average in the individual natural conditions scenarios, and p_{act} is the same but for the actual conditions. From Fig. 14.2b, it is shown that the persistent drought of 2014 was made more likely because of the anthropogenic climate change signal, indicating that patterns of precipitation are shifting to a drier state in this region. The mean FAR is ~ 0.45 , indicating that anthropogenic climate change made the event $\sim 45\%$ more likely. This is consistent with the predicted weakening of the Mediterranean storm tracks in a warming climate (Hatzaki et al. 2009), a response that is captured in our model (Anagnostopoulou et al. 2006). The increased SSTs between our actual and natural simulations are relatively uniform across the globe (e.g., Schaller et al. 2015, manuscript submitted to *Nat. Climate Change*, their Fig. S3), and Hoerling et al. 2012 showed that this would induce an Eastern Mediterranean drying through circulation changes, also consistent with our detected response.

Discussion. In this study, we have used local station data to understand the uniquely persistent drought that occurred in the southern Levant rainy season of 2014. We show that the event was unprecedented for the critical January–February period in the observational record, and through modeling the event, we showed that anthropogenic climate change made it $\sim 45\%$ more likely. However, our model is dry-biased in the region of interest, most likely due to a lack of very intense cyclones in the Mediterranean region (Anagnostopoulou et al. 2006). If changes in these processes are nonlinear under human-induced emissions, our model is unlikely to capture them fully. This wet season is when reservoirs and groundwater systems are recharged and snow pack accumulates to support summer streamflows. The consequent external stresses that came with this drought, such as crop failures, degraded grazing land, and overpumping of nonrenewable groundwater, suggest that water and agriculture authorities in the southern Levant region should have additional fail-safes in place going into the future.

ACKNOWLEDGMENTS. This project was funded by NERC ACE Africa, and by USAID Office of the Middle East Programme’s MAWRED project. We would like to thank the Met Office Hadley Centre PRECIS team and our colleagues at the Oxford eResearch Centre for their technical and scientific support for the development and application of Weather@Home. Finally, we thank all of the

volunteers who have donated their computing time to Weather@Home.

REFERENCES

- Allen, M., 2003: Liability for climate change. *Nature*, **421**, 891–892.
- Anagnostopoulou, Chr., K. Tolika, H. Flocas, and P. Maheras, 2006: Cyclones in the Mediterranean region: present and future climate scenarios derived from a general circulation model (HadAM3P). *Adv. Geosci.*, **7**, 9–14.
- Black, E., 2009: The impact of climate change on daily precipitation statistics for Jordan and Israel. *Atmos. Sci. Lett.*, **10**, 192–200, doi:10.1002/asl.233.
- Boe, J., L. Terray, F. Habets, and E. Martin, 2007: Statistical and dynamical downscaling of the Seine basin climate for hydro-meteorological studies. *Int. J. Climatol.*, **27**, 1643–1655, doi:10.1002/joc.1602.
- Hatzaki, M., H. A. Flocas, C. Giannakopoulos, and P. Maheras, 2009: The impact of the eastern Mediterranean teleconnection pattern on the Mediterranean climate. *J. Climate*, **22**, 977–992.
- Haylock, M. R., N. Hofstra, A. M. G. Klein Tank, E. J. Klok, P. D. Jones, and M. New, 2008: A European daily high-resolution gridded dataset of surface temperature and precipitation. *J. Geophys. Res.*, **113**, D20119, doi:10.1029/2008JD10201.
- Herring, S. C., M. P. Hoerlin, T. C. Peterson, and P. A. Stott, Eds., 2014: Explaining extreme events of 2013 from a climate perspective. *Bull. Amer. Meteor. Soc.*, **95** (9), S1–S104.
- Hoerling, M., J. Eischeid, J. Perlwitz, X. Quan, T. Zhang, and P. Pegion, 2012: On the increased frequency of Mediterranean drought. *J. Climate*, **25**, 2146–2161, doi:10.1175/JCLI-D-11-00296.1.
- Massey, N., and Coauthors, 2015: weather@ home—development and validation of a very large ensemble modelling system for probabilistic event attribution. *Quart. J. Roy. Meteor. Soc.*, **141**, 1528–1545, doi:10.1002/qj.2455.
- Saaroni, H., B. Ziv, J. Lempert, Y. Gazit, and E. Morin, 2014: Prolonged dry spells in the Levant region: Climatologic-synoptic analysis. *Int. J. Climatol.*, **35**, 2223–2236, doi:10.1002/joc.4143.
- Schaller, N., F. E. L. Otto, G. J. van Oldenborgh, N. R. Massy, S. Sparrow, and M. R. Allen, 2014: The heavy precipitation event of May–June 2013 in the upper Danube and Elbe basins [in “Explaining Extreme Events of 2013 from a Climate Perspective”]. *Bull. Amer. Meteor. Soc.*, **95** (9), S69–S7.

- Sipple, S., D. Mitchell, M. T. Black, A. J. Dittus, L. Harrington, N. Schaller, and F. E. L. Otto, 2015: Combining large model ensembles with extreme value statistics to improve attribution statements of rare events. *Wea. Climate Extremes*, **9**, 25-35, doi:10.1016/j.wace.2015.06.004.
- Solomon, S., D. Qin, M. Manning, Z. Chen, M. Marquis, K. Averyt, M. Tignor, and H. L. Miller Jr., Eds., 2007: *Climate Change 2007: The Physical Science Basis*. Cambridge University Press, 996 pp.
- Stark, J. D., C. J. Donlon, M. J. Martin, and M. E. McCulloch, 2007: OSTIA: An operational, high resolution, real time, global sea surface temperature analysis system. *Proc., Oceans 2007, Marine Challenges: Coastline to Deep Sea*, Aberdeen, Scotland, IEEE/OES, 1-4, doi:10.1109/OCEANSE.2007.4302251.
- Udasin, S., 2014: Israel experiencing unprecedented drought conditions. *Jerusalem Post*, 10 February 2014. [Available online at www.jpost.com/Enviro-Tech/Israel-experiencing-unprecedented-drought-conditions-341003.]

Table 34.I. ANTHROPOGENIC INFLUENCE

ON EVENT STRENGTH †

	INCREASE	DECREASE	NOT FOUND OR UNCERTAIN
Heat	Australia (Ch. 31) Europe (Ch.13) S. Korea (Ch. 19)		Australia, Adelaide & Melbourne (Ch. 29) Australia, Brisbane (Ch.28)
Cold		Upper Midwest (Ch.3)	
Winter Storms and Snow			Eastern U.S. (Ch. 4) N. America (Ch. 6) N. Atlantic (Ch. 7)
Heavy Precipitation	Canada** (Ch. 5)		Jakarta**** (Ch. 26) United Kingdom*** (Ch. 10) New Zealand (Ch. 27)
Drought	E. Africa (Ch. 16) E. Africa* (Ch. 17) S. Levant (Ch. 14)		Middle East and S.W. Asia (Ch. 15) N.E. Asia (Ch. 21) Singapore (Ch. 25)
Tropical Cyclones			Gonzalo (Ch. 11) W. Pacific (Ch. 24)
Wildfires			California (Ch. 2)
Sea Surface Temperature	W. Tropical & N.E. Pacific (Ch. 20) N.W. Atlantic & N.E. Pacific (Ch. 13)		
Sea Level Pressure	S. Australia (Ch. 32)		
Sea Ice Extent			Antarctica (Ch. 33)

† Papers that did not investigate strength are not listed.

†† Papers that did not investigate likelihood are not listed.

* No influence on the likelihood of low rainfall, but human influences did result in higher temperatures and increased net incoming radiation at the surface over the region most affected by the drought.

** An increase in spring rainfall as well as extensive artificial pond drainage increased the risk of more frequent severe floods from the enhanced rainfall.

*** Evidence for human influence was found for greater risk of UK extreme rainfall during winter 2013/14 with time scales of 10 days

**** The study of Jakarta rainfall event of 2014 found a statistically significant increase in the probability of such rains over the last 115 years, though the study did not establish a cause.

	ON EVENT LIKELIHOOD ††			Total Number of Papers
	INCREASE	DECREASE	NOT FOUND OR UNCERTAIN	
Heat	Argentina (Ch. 9) Australia (Ch. 30, Ch. 31) Australia, Adelaide (Ch. 29) Australia, Brisbane (Ch. 28) Europe (Ch. 13) S. Korea (Ch. 19) China (Ch. 22)		Melbourne, Australia (Ch. 29)	7
Cold		Upper Midwest (Ch.3)		1
Winter Storms and Snow	Nepal (Ch. 18)		Eastern U.S. (Ch. 4) N. America (Ch. 6) N. Atlantic (Ch. 7)	4
Heavy Precipitation	Canada** (Ch. 5) New Zealand (Ch. 27)		Jakarta**** (Ch. 26) United Kingdom*** (Ch. 10) S. France (Ch. 12)	5
Drought	E. Africa (Ch. 16) S. Levant (Ch. 14)		Middle East and S.W. Asia (Ch. 15) E. Africa* (Ch. 17) N.E. Asia (Ch. 21) S. E. Brazil (Ch. 8) Singapore (Ch. 25)	7
Tropical Cyclones	Hawaii (Ch. 23)		Gonzalo (Ch. 11) W. Pacific (Ch. 24)	3
Wildfires	California (Ch. 2)			1
Sea Surface Temperature	W. Tropical & N.E. Pacific (Ch. 20) N.W. Atlantic & N.E. Pacific (Ch. 13)			2
Sea Level Pressure	S. Australia (Ch. 32)			1
Sea Ice Extent			Antarctica (Ch. 33)	1
TOTAL				32

† Papers that did not investigate strength are not listed.

†† Papers that did not investigate likelihood are not listed.

* No influence on the likelihood of low rainfall, but human influences did result in higher temperatures and increased net incoming radiation at the surface over the region most affected by the drought.

** An increase in spring rainfall as well as extensive artificial pond drainage increased the risk of more frequent severe floods from the enhanced rainfall.

*** Evidence for human influence was found for greater risk of UK extreme rainfall during winter 2013/14 with time scales of 10 days

**** The study of Jakarta rainfall event of 2014 found a statistically significant increase in the probability of such rains over the last 115 years, though the study did not establish a cause.