On the Drivers and Predictability of Seasonal-to-Interannual Variations in Regional Sea Level[Ⓐ]

C. D. ROBERTS, D. CALVERT, N. DUNSTONE, L. HERMANSON, M. D. PALMER, AND D. SMITH

Met Office Hadley Centre, Exeter, United Kingdom

(Manuscript received 3 December 2015, in final form 20 July 2016)

ABSTRACT

Observations and eddy-permitting ocean model simulations are used to evaluate the drivers of sea level variability associated with 15 modes of climate variability covering the Atlantic, Pacific, Indian, and Southern Oceans. Sea level signals are decomposed into barotropic, steric, and inverted barometer contributions. Forcings are decomposed into surface winds, buoyancy fluxes, and Ekman pumping. Seasonal-to-interannual sea level variability in the low latitudes is governed almost entirely by the thermosteric response to wind forcing associated with tropical modes of climate variability. In the extratropics, changes to dynamic sea level associated with atmospheric modes of variability include a substantial barotropic response to wind forcing, particularly over the continental shelf seas. However, wind-driven steric changes are also important in some locations. On interannual time scales, wind-forced steric changes dominate, although heat and freshwater fluxes are important in the northwest Atlantic, where low-frequency sea level variations are associated with changes in the Atlantic meridional overturning circulation. Using the version 3 of the Met Office Decadal Prediction System (DePreSys3), the predictability of large-scale dynamic sea level anomalies on seasonal-to-interannual time scales is evaluated. For the first year of the hindcast simulations, DePreSys3 exhibits skill exceeding persistence over large regions of the Pacific, Atlantic, and Indian Oceans. Skill is particularly high in the tropical Indo-Pacific because of the accurate initialization and propagation of thermocline depth anomalies associated with baroclinic adjustments to remote wind forcing. Skill in the extratropics is hindered by the limited predictability of wind anomalies associated with modes of atmospheric variability that dominate local and/or barotropic responses.

1. Introduction

a. Background and motivation

The timing and magnitude of coastal sea level extremes are largely governed by the phase of astronomical tides and the sea level response to meteorological phenomena (e.g., storm surges, ocean swell, surface waves, and wave setup; Wong et al. 2014). However, changes to local time-mean sea level due to long-term trends and natural modes of climate variability exert a significant influence on the background risk of such extremes (Nicholls et al. 1999). The physical consequences of extreme sea level events include coastal submergence

and flooding, saltwater intrusion into ground and surface waters, and coastal erosion, all of which can have negative social and economic impacts (Wong et al. 2014; Nicholls and Cazenave 2010).

Over the period 1993–2009, global mean sea level (GMSL) increased at a rate of $3.2 \pm 0.4 \text{ mm yr}^{-1}$ because of thermal expansion and the transfer of mass from the continents to the oceans (Church and White 2011). However, on monthly-to-decadal time scales, regional sea level anomalies associated with large-scale modes of internal climate variability can dominate long-term changes. For example, sea level anomalies in the tropical Pacific associated with the year-to-year changes in the El Niño–Southern Oscillation (ENSO) can exceed 30 cm (Becker et al. 2012), which is larger than the 21-cm increase in GMSL that occurred between 1880 and 2009 (Church and White 2011). There are also substantial geographic variations in rates of sea level rise on decadal time scales due to the influence of climate variability (Zhang and Church 2012). For example, rates of sea level rise in the western tropical Pacific during the last decade have been 2–3 times

[Ⓐ] Supplemental information related to this paper is available at the Journals Online website: <http://dx.doi.org/10.1175/JCLI-D-15-0886.s1>.

Corresponding author address: C. D. Roberts, Met Office Hadley Centre, Fitzroy Road, Exeter EX1 3P, United Kingdom.
E-mail: chris.roberts@metoffice.gov.uk

higher than the global mean (Stammer et al. 2013). Skillful and timely prediction of such anomalies will allow coastal communities to better assess the risk of extreme coastal flooding events and therefore inform strategies to mitigate the associated impacts.

b. Dynamic sea level

In this study, we focus on the seasonal-to-interannual variations in dynamic sea level (DSL) associated with leading modes of climate variability. We define DSL as regional deviations from GMSL associated with local variations in ocean mass and density arising from changes in the barotropic and baroclinic ocean circulation, respectively. Following Gill and Niller (1973), DSL anomalies can be expressed as

$$\text{DSL} = \eta_{\text{barotropic}} + \eta_{\text{steric}} = \frac{p_b}{g\rho_0} - \frac{1}{\rho_0} \int_{-H}^0 \Delta\rho(T, S, p) dz, \quad (1)$$

where g is acceleration due to gravity, T is ocean potential temperature, S is salinity, p is pressure, p_b is pressure at the ocean bottom, ρ_0 is a reference density, and $\Delta\rho$ is potential density relative to a reference value. By focusing on the dynamic component of regional sea level, we neglect contributions from global thermosteric sea level rise, vertical land motion, changes to Earth's geoid, and loss or addition of mass to the ocean. In addition, our definition of DSL does not include contributions from the inverted barometer (IB) effect (i.e., the change in sea level induced by changes in local atmospheric pressure). Observational estimates of DSL are derived from satellite altimeter data and are adjusted to remove IB effects. However, where specified, we separately evaluate contributions from the IB effect to compare with changes in DSL.

c. Drivers of seasonal-to-interannual sea level variability

To understand the origins and limits of DSL predictability, it is instructive to consider the processes that drive variability. These can include local and remote atmospheric forcings and intrinsic (i.e., unforced) ocean variability arising from nonlinear processes. DSL signals associated with local atmospheric forcings and rapid barotropic adjustments present a particular challenge because they require accurate prediction of the evolving atmospheric state, which has significantly less "memory" than the ocean. In contrast, ocean advective processes and baroclinic responses to remote forcings can be captured through accurate initialization of the ocean dynamics without requiring predictability of the atmospheric forcing. Table 1 summarizes some of the previously described processes responsible for DSL variability along with their

typical time scales. These dynamic sources of variability are often linked to large-scale modes of climate variability such as ENSO and the North Atlantic Oscillation (NAO; Stammer et al. 2013), though others have argued that a substantial fraction of interannual-to-decadal time-scale variability can be explained by the baroclinic response to stochastic wind forcing (e.g., Frankignoul et al. 1997).

Local sea level responses to atmospheric forcing include steric changes due to surface buoyancy forcings (e.g., Gill and Niller 1973; Ferry et al. 2000; Cabanes et al. 2006), Ekman pumping in the presence of stratification (Cabanes et al. 2006; Piecuch and Ponte 2011; Cummins and Lagerloef 2004), and along-equator heave of the thermocline balanced by changes to zonal wind stress (Merrifield and Maltrud 2011; Timmermann et al. 2010). Remote sea level responses to atmospheric forcings include changes associated with baroclinic and barotropic wave adjustments to wind forcing (Wyrtki 1975; Forget and Ponte 2015; Cabanes et al. 2006; Qiu 2002; Frankignoul et al. 1997; Hakkinen 2001; Sturges and Hong 1995; Qiu and Chen 2006), advective anomalies in response to buoyancy-forced water mass transformations (e.g., Hakkinen 2001; Köhl 2014), and time-dependent Sverdrup dynamics associated with zonally integrated wind stress curl (Cabanes et al. 2006; Qiu 2002). Unforced intrinsic variability also contributes to changes in sea level and arises because of nonlinear ocean processes such as barotropic and baroclinic instability (e.g., Chelton et al. 2007; Penduff et al. 2011).

Barotropic redistribution of mass is important for DSL variability on intra-annual time scales (Fukumori et al. 1998; Vinogradova et al. 2007). On interannual time scales, barotropic changes play a role in the high latitudes and over shallow shelf seas (Forget and Ponte 2015; Piecuch et al. 2013), but in most other regions DSL variability is largely steric in nature (Cazenave and Llovel 2010; Forget and Ponte 2015; Köhl 2014). In the low and midlatitudes, steric signals are almost entirely thermosteric and driven by the adiabatic displacement of isopycnals driven by wind stress (Köhl 2014; Forget and Ponte 2015; Stammer et al. 2013). In contrast, sea level variability in the North Atlantic western boundary and subpolar gyre has been linked to the diabatic response to basin-scale thermal forcing and changes in the overturning circulation (e.g., Hakkinen 2001). Salinity-driven (i.e., halosteric) changes make an important contribution in the Arctic, but elsewhere in the ocean they are mostly offset by density-compensating thermosteric changes (Antonov et al. 2002; Köhl 2014).

d. Recent studies of sea level predictability

Several studies have demonstrated the ability of statistical techniques to make useful predictions of sea

TABLE 1. Processes contributing to seasonal-to-interannual dynamic sea level variability.

Process	Driver	Approx time scale	References
Remotely forced Baroclinic Rossby waves	Wind stress	Months to decades	Cabanes et al. (2006) Cummins and Lagerloef (2004) Forget and Ponte (2015) Frankignoul et al. (1997) McGregor et al. (2007) Qiu (2002) Qiu and Chen (2006) Qiu and Chen (2010) Sturges and Hong (1995) Vivier et al. (1999)
Boundary-forced Rossby waves	Perturbations at an eastern boundary	Months to decades	Cabanes et al. (2006) Jacobs et al. (1994) Qiu and Chen (2006)
Kelvin waves	Wind stress	Months to years	Enfield and Allen (1980) Forget and Ponte (2015) McGregor et al. (2007) Wyrski (1975)
Time-dependent Sverdrup balance	Zonal integral of wind stress curl	Months to years	Cabanes et al. (2006) Qiu (2002) Stammer (1997) Vivier et al. (1999)
Steric (advection)	Advection of temperature/salinity anomalies	Years to decades	Cabanes et al. (2006) Forget and Ponte (2015) Köhl (2014) Piecuch and Ponte (2011)
Buoyancy-forced circulation changes	Heat and freshwater fluxes in regions of water mass formation	Years to decades	Hakkinen (2001)
Barotropic adjustments	Wind stress and mass fluxes; steric changes over sloping topography	Days to years	Hakkinen and Rhines (2004) Fukumori et al. (1998) Vinogradova et al. (2007)
Locally forced Ekman pumping	Local wind stress curl	Months to years	Cabanes et al. (2006) Hakkinen (2001) Kelly et al. (1993) Köhl (2014) Vivier et al. (1999)
Steric (local)	Local air–sea buoyancy fluxes	Months to decades	Cabanes et al. (2006) Ferry et al. (2000) Gill and Niller (1973) Köhl (2014) Piecuch and Ponte (2011) Stammer (1997) Vivier et al. (1999)
Equatorial thermocline heave	Zonal wind stress	Months to decades	Merrifield and Maltrud (2011) Timmermann et al. (2010)
Unforced (i.e., intrinsic) Eddies	Baroclinic–barotropic instability	Days to years	Cabanes et al. (2006) Chelton et al. (2007)
Mixing	Temperature and salinity gradients	Years to decades	Piecuch and Ponte (2011)

level variability in the tropical Pacific at lead times of several months (e.g., Xue and Leetmaa 2000; Chowdhury et al. 2007, 2014). These methods rely on historical relationships between sea surface temperature (SST), surface winds, and sea level to make predictions of variability at selected tide gauge locations. Seasonal outlooks based on these statistical methods

are currently made available online by the National Oceanic and Atmospheric Administration (NOAA) for the U.S.-affiliated Pacific Islands (<http://www.weather.gov/peac/sealevel>).

Although statistical predictions are useful, they are reliant on the use of a training dataset such that their application is limited to locations with sufficiently long

TABLE 2. Indices of climate variability used in this study.

Mode of variability	Abbreviation	Type of index	Reference
Antarctic Oscillation	AAO	Atmospheric pressure	Mo (2000)
Atlantic meridional mode	AMM	SST and surface winds	Chiang and Vimont (2004)
Atlantic Niño-1 index	ANINO	SST	Lutz et al. (2013)
East Atlantic pattern	EA	Atmospheric pressure	Barnston and Livezey (1987)
East Atlantic–Western Russia pattern (Eurasia-2)	EAWR	Atmospheric pressure	Barnston and Livezey (1987)
East Pacific–North Pacific pattern	EPNP	Atmospheric pressure	Barnston and Livezey (1987)
El Niño Modoki index	EMI	SST	Ashok et al. (2007)
Indian Ocean dipole	IOD	SST	Saji et al. (1999)
North Atlantic Oscillation	NAO	Atmospheric pressure	Barnston and Livezey (1987)
North Pacific index	NP	Atmospheric pressure	Trenberth and Hurrell (1994)
Pacific Niño-3.4 index	ENSO	SST	Trenberth (1997)
Pacific–North American pattern	PNA	Atmospheric pressure	Barnston and Livezey (1987)
Polar–Eurasia pattern	POL	Atmospheric pressure	Barnston and Livezey (1987)
Scandinavian pattern (Eurasia-1)	SCA	Atmospheric pressure	Barnston and Livezey (1987)
West Pacific pattern	WP	Atmospheric pressure	Barnston and Livezey (1987)

records of reliable and well-documented tide gauge observations. In contrast, global climate models that include sophisticated numerical representations of the dynamics and thermodynamics of the oceans and atmosphere can, in principle, provide a forecast for any coastal or open-ocean location. Such models have recently been applied to investigate the predictability of sea level on seasonal (Miles et al. 2014; McIntosh et al. 2015) and decadal (Polkova et al. 2014, 2015) time scales.

Miles et al. (2014) used a dynamical general circulation model (GCM) to predict regional sea level anomalies over the global ocean at lead times of up to 7 months. Although using an ocean model with a relatively coarse horizontal resolution ($\sim 2^\circ$), Miles et al. (2014) found skill in the prediction of sea level anomalies in the tropical Pacific. In a follow-up study, McIntosh et al. (2015) extended the analysis to coastal locations in the tropical Indian and Pacific Oceans and emphasized the value of dynamical GCMs by demonstrating skill in the prediction of coastal sea level that exceeded existing statistical approaches. Polkova et al. (2015) used a GCM-based decadal prediction system and found predictability of steric sea level in the subtropics for a lead time of 2–5 yr. They attributed this skill to westward-propagating baroclinic Rossby waves. However, this skill was difficult to separate from persistence at high latitudes, where Rossby waves can take decades to cross a basin.

e. This study

Here, we combine observations and experiments with eddy-permitting GCMs to evaluate the drivers and predictability of DSL on seasonal-to-interannual time scales. In particular, we focus on the response of DSL to 15 leading modes of climate variability covering the

Atlantic, Indian, Pacific, and Southern Oceans. Section 2 describes the data, models, and methods we use in our analysis. Section 3 evaluates the drivers of DSL associated with modes of climate variability on seasonal-to-interannual time scales. Section 4 evaluates the predictability of large-scale DSL variability in a state-of-the-art coupled forecasting system. Our main conclusions are summarized in section 5.

2. Data and model configurations

a. Satellite altimeter data

We use satellite altimeter data from a multimission reanalysis produced by the European Space Agency (ESA) Climate Change Initiative (CCI; <http://www.esa-sealevel-cci.org>). This product is described in detail by Ablain et al. (2015). Additional preprocessing steps applied for this study include bilinear interpolation to a regular $1^\circ \times 1^\circ$ grid and removal of the global trend by subtraction of the area-weighted global mean from each monthly field. Monthly anomalies are calculated by subtracting the climatological seasonal cycle for the period 1993–2012.

b. Climate mode indices

Here, we evaluate the drivers of sea level anomalies associated with 15 large-scale modes of climate variability (see Table 2) chosen for their coverage of the major ocean basins and importance for seasonal-to-interannual climate variability, including some that generally receive less attention in the sea level literature. It is well-known that some modes of climate variability, such as ENSO and the NAO, are associated with basin-scale changes in sea level (Wyrski 1975; Enfield

and Allen 1980; Chelton and Davis 1982; Zhang and Church 2012; Goddard et al. 2015; Tsimplis et al. 2006; Yan et al. 2004; Woolf et al. 2003; Woodworth et al. 2007). However, less well known modes of climate variability also have a significant influence on regional sea level and the risk of extreme events (Menéndez and Woodworth 2010). For example, Fagherazzi et al. (2005) found that variations in the east Atlantic–western Russia (EAWR) and polar–Eurasian (POL) indices were linked to the frequency of flooding in Venice.

Monthly indices are retrieved from NOAA (<http://www.esrl.noaa.gov/psd/data/climateindices/list/>) and the KNMI climate explorer (<https://climexp.knmi.nl/>). In addition, the El Niño Modoki index (EMI) and Atlantic Niño-1 index are based on data from Ashok et al. (2007) and Lutz et al. (2013), respectively. Each index is detrended and anomalized relative to its mean seasonal cycle for the period 1993–2012 in order to isolate changes on seasonal-to-interannual time scales.

c. Ocean model sensitivity experiments

To complement our analysis of observational data, we perform sensitivity experiments with an eddy-permitting ocean model. Our experiments are designed to isolate the historical variations in DSL attributable to the following processes: 1) mechanical forcing from surface winds and 2) buoyancy fluxes that arise because of variations in the overlying atmospheric state. We use version 3.4 of the Nucleus for European Models of the Ocean (NEMO) global ocean model (Madec 2008) configured with 75 vertical levels and an eddy-permitting horizontal resolution of $\sim 0.25^\circ$ coupled to the Los Alamos National Laboratory sea ice model (CICE), version 4.1 (Hunke and Lipscomb 2010). Surface boundary conditions are specified using a mixture of time-varying and climatological “normal year” versions of the updated Coordinated Ocean–Ice Reference Experiments, version 2 (CORE-II) dataset (Large and Yeager 2009). Turbulent air–sea fluxes are calculated using the bulk formulae of Large and Yeager (2004). The normal year forcing consists of a single annual cycle that is representative of climatological conditions but also includes realistic propagation of synoptic weather patterns such that the ocean is excited by high-frequency forcings (Large and Yeager 2004). Previous studies have demonstrated that eddy-permitting configurations of the NEMO model are useful tools for simulating variability of sea level (Penduff et al. 2010) and large-scale ocean dynamics (Roberts et al. 2013). The specific version of NEMO we use is designated Global Ocean 5 (GO5) and is described in detail by Megann et al. (2014).

The four sensitivity experiments used in this study can be summarized as follows:

- 1) CTRL includes interannual variability in all atmospheric boundary conditions used to calculate air–sea fluxes of heat, freshwater, and momentum.
- 2) CTRL_NYF is run parallel to CTRL but with atmospheric boundary conditions constrained to follow a repeating normal year cycle such that interannual variability can only arise as a consequence of adjustments to the initial conditions and unforced intrinsic ocean variability. This experiment is used to correct for multidecadal adjustments in our ocean experiments that are unrelated to interannual variability in the atmosphere (i.e., drift).
- 3) WIND is as CTRL, but with normal year atmospheric boundary conditions used to calculate all air–sea exchanges other than momentum fluxes. Note that this experiment still includes some interannual variations in fluxes of sensible/latent heat and outgoing longwave radiation that arise in response to wind-induced changes in SST. This contribution to heat flux variability is dominant in regions of equatorial upwelling, but because the feedback is negative (i.e., negative SST anomalies induce a heat flux anomaly that warms the ocean), it acts to damp ocean variability and we therefore do not consider it a buoyancy forcing.
- 4) BUOY includes interannual variability in all atmospheric boundary conditions that affect buoyancy fluxes (i.e., precipitation, downward radiation fluxes, air temperature, humidity, and wind speed) but uses normal year winds to calculate air–sea momentum fluxes. This experiment therefore includes variations in heat and freshwater fluxes that originate from the atmosphere and can be considered to be driving the ocean.

A more complete description of these experiments, including our drift-correction strategy, is included as supplementary material.

d. DePreSys3

To evaluate the predictability of DSL on seasonal-to-interannual time scales, we use version 3 of the Met Office Decadal Prediction System (DePreSys3; Dunstone et al. 2016). DePreSys3 uses the Hadley Centre Global Environmental Model version 3 at the global coupled model 2.0 (HadGEM3-GC2) configuration (Williams et al. 2015). It has an atmospheric resolution of 0.83° longitude $\times$ 0.55° latitude (about 60 km at midlatitudes), 85 atmospheric levels, and an upper boundary at 85 km near the mesopause. The ocean model used is the NEMO GO5 configuration described in section 2c. DePreSys3 uses a data assimilation scheme that nudges toward observed analyses in the atmosphere, ocean, and sea ice

TABLE 3. Dominant drivers of sea level variability associated with different modes of climate variability. When more than one type of sea level response or atmospheric driver is involved, they are listed in order of importance.

Mode	Region	Sea level responses	Atmospheric forcings
AAO	Southern Ocean	IB, barotropic	Winds
AMM	Tropical Atlantic	Thermosteric	Winds, buoyancy fluxes
ANINO	Tropical Atlantic	Thermosteric	Winds
EA	North Atlantic	IB, thermosteric, barotropic, halosteric	Winds, buoyancy fluxes, Ekman pumping
EAWR	Northeast Atlantic	IB, barotropic	Winds
EMI	Indo-Pacific	Thermosteric, IB, halosteric	Winds, buoyancy fluxes
ENSO	Indo-Pacific	Thermosteric, IB, halosteric	Winds, buoyancy fluxes
EPNP	North Pacific	IB, thermosteric, halosteric, barotropic	Winds, buoyancy fluxes, Ekman pumping
IOD	Indian Ocean	Thermosteric, halosteric, IB	Winds, buoyancy fluxes
NAO	North Atlantic	IB, barotropic, thermosteric, halosteric	Winds, buoyancy fluxes, Ekman pumping
NP	North Pacific	IB, barotropic, thermosteric	Winds, Ekman pumping
PNA	North Pacific	IB, barotropic, thermosteric	Winds, Ekman pumping
POL	North Atlantic	Thermosteric, halosteric, barotropic, IB	Winds, buoyancy
SCA	Northeast Atlantic	IB, thermosteric, barotropic	Winds
WP	North Pacific	IB, barotropic, thermosteric	Winds, buoyancy

(Knight et al. 2014) during an assimilation run of the coupled model. The atmosphere is nudged to 6-hourly fields of ERA-Interim (Dee et al. 2011) temperatures and winds with a 6-hourly relaxation time scale. The ocean is nudged with a 10-day relaxation time scale toward monthly mean fields of 3D temperature and salinity from a statistical ocean analysis based on global covariances (Smith and Murphy 2007; Smith et al. 2015). Sea ice concentration is nudged to monthly mean fields from the HadISST dataset (Rayner et al. 2003) using a 24-h relaxation time scale. In contrast to previous versions, DePreSys3 assimilates absolute values of temperature and salinity rather than adding observed anomalies onto the model climatology. This approach is sometimes referred to as “full field” initialization. No direct observations of sea level are used to constrain the DePreSys3 analysis.

Hindcasts (retrospective forecasts) cover the period 1960–2014 with a total of 25 start dates spaced 2–3 yr apart between 1960 and 2008 and 1 yr apart from 2008 onward. Each hindcast consists of 10 ensemble members that are started from the initial conditions provided by the assimilation run on 1 November. The ensemble is created by using different seeds to a stochastic physics scheme (MacLachlan et al. 2015). The assimilation run and hindcasts have full knowledge of external forcing datasets (e.g., greenhouse gases, aerosols, ozone, and solar and volcanic forcings) as per the CMIP5 protocol (Taylor et al. 2012) and follow the representative concentration pathway 4.5 (RCP4.5) after 2005 (Meinshausen et al. 2011).

Forecasts from dynamic models using full-field initialization have a tendency to drift toward a “preferred” model climatology, owing to systematic errors in the model simulation. These biases can be accounted for using a lead-time-dependent correction diagnosed

from a set of hindcasts (Smith et al. 2013). This correction is necessary if time series from individual hindcasts are being evaluated or if forecast skill is calculated using a measure of deviation (e.g., root-mean-square errors). In contrast, Pearson correlation coefficients calculated using data averaged over a specified forecast period are insensitive to model biases and therefore do not require a lead-time-dependent correction (Smith et al. 2013). For this reason, we evaluate predictive skill for each lead time using correlation coefficients between hindcast ensemble means and equivalent periods in the data assimilating model analysis. The model analysis of DSL is used to assess predictive skill in DePreSys3 because of the limited length of the satellite altimeter record. For this reason, our estimates of skill do not take into account uncertainties in the initial ocean state.

3. Results

a. Observed and simulated dynamic sea level variability

Many previous studies have discussed the dynamical origins and complex geographical distribution of processes responsible for sea level variability (see Table 1). Here, we complement the existing literature by investigating the sea level response associated with 15 leading modes of climate variability using a combination of satellite data and sensitivity experiments with an eddy-permitting ocean model. A summary of the main results from this section is presented in Table 3.

To begin, we calculate the spatial fingerprints of DSL associated with each mode of variability. Fingerprints are defined as the differences in DSL between positive and negative phases of the associated climate index and

are calculated using composite means of the 50 most positive and 50 most negative events during the period 1993–2012 (Fig. 1). As expected, there are large changes in DSL associated with tropical modes of climate variability such as ENSO. These signals are particularly evident in the Pacific and Indian Oceans, but less so in the Atlantic. The role for wind-forced baroclinic waves in tropical modes of variability such as ENSO has been understood for several decades (e.g., Wyrtki 1975; Enfield and Allen 1980). Propagating Rossby and Kelvin waves are particularly clear in the tropics and along coastal–equatorial waveguides when composite patterns of DSL are calculated using different lags (see supplementary Fig. 1 in supplementary material).

Extratropical modes of climate variability are also linked to distinct changes in DSL. For example, the NAO is associated with a clear tripole pattern of DSL that exhibits particularly large signals in continental shelf regions such as the Baltic and Mediterranean. However, the DSL signal associated with extratropical modes of variability is typically of a similar (or smaller) magnitude to changes because of the IB effect (Fig. 2). Moreover, in many cases the IB effect and dynamic changes compensate one another [e.g., the central North Pacific during North Pacific index (NP) events]. In other instances, the IB and DSL changes are additive and are therefore more likely to have significant coastal impacts (e.g., the NAO impact on the northeastern United States and the EA impact on the west coast of the United Kingdom and Ireland). This comparison emphasizes that skillful prediction of sea level in the extratropics requires accurate predictions of both the ocean and atmospheric state.

We recalculate the same sea level fingerprints using output from the eddy-permitting ocean model experiment, CTRL (Fig. 1). Each model composite is calculated using the 50 most positive (or negative) months of the associated climate index within the 1993–2009 period of overlap between the altimeter data and our model experiments. In general, the model CTRL simulation does an excellent job of reproducing the DSL signal associated with modes of seasonal-to-interannual climate variability. However, there are some deficiencies worth noting, including the tendency for anomalies in the tropical Pacific associated with ENSO and EMI to extend too far north. Lead-lag analysis of model composite patterns (not shown) reveals this particular bias is a result of Rossby waves emitted from the eastern Pacific boundary at $\sim 10^\circ\text{N}$ that propagate westward too rapidly compared to observations, suggesting a stratification bias in this region. Other deficiencies include a tendency for the model to underestimate the response of the North Atlantic marginal

seas to NAO variability and inaccurate responses to the EA in parts of the North Sea, Baltic Sea, and Mediterranean Sea. Despite these shortcomings, the overall agreement with the observed signals justifies the use of our model sensitivity experiments to understand the processes driving DSL variability.

b. Isolating the drivers of dynamic sea level variability

Separation of the local barotropic and steric contributions to changes in DSL in CTRL (Fig. 3) reveals that a substantial fraction of the DSL signal associated with high-latitude modes of climate variability is barotropic in nature, particularly on the continental shelves. Steric signals are also important for extratropical modes of variability in the North Atlantic (EA and NAO) and North Pacific (PNA and EPNP), but less so in the Southern Ocean (AAO). These results are consistent with previous studies that have emphasized the role of barotropic variability in the high latitudes and over shallow shelf seas (Vinogradova et al. 2007). In contrast, changes in DSL associated with tropical modes of variability in the Indian, Pacific, and Atlantic Oceans are almost entirely steric.

To investigate the role of temperature and salinity in sea level variability, we decompose the steric signal into thermosteric and halosteric components as follows

$$\eta_{\text{thermosteric}} = -\frac{1}{\rho_0} \int_{-H}^0 \Delta\rho(T, \bar{S}, p) dz \quad \text{and} \quad (2)$$

$$\eta_{\text{halosteric}} = -\frac{1}{\rho_0} \int_{-H}^0 \Delta\rho(\bar{T}, S, p) dz, \quad (3)$$

where $\bar{T}$ and $\bar{S}$ are climatological values for temperature and salinity, respectively. This decomposition reveals that steric changes associated with the modes of climate variability in Fig. 3 are almost exclusively due to changes in temperature, with changes in salinity mostly acting to offset a fraction of the thermosteric signal (Fig. 4). An exception is the EPNP pattern in the northeast Pacific, where halosteric effects appear to be important.

To identify the relative role of the winds and buoyancy forcings in driving DSL changes associated with each mode of climate variability, we calculate the same composite patterns using the model simulations WIND and BUOY (Fig. 5). These experiments demonstrate that changes in DSL associated with seasonal-to-interannual modes of climate variability are largely explained by the response to wind forcing. Buoyancy forcings also make a contribution to steric changes associated with some modes, notably in the tropics (e.g., AMM, ENSO, and EMI) and the North Atlantic (NAO, EA, and POL). Our results for the NAO are consistent

FIG. 1. Dynamic sea level (mm) fingerprints associated with 15 modes of climate variability from (top) altimeter data and (bottom) experiment CTRL. Fingerprints are defined as the difference between composite means of the 50 most positive and 50 most negative months of the associated climate index.

FIG. 2. As in Fig. 1, but for (top) the IB effect (mm) defined as $\eta_{IB} = -p_{\text{surface}}\rho_0/g$ and (bottom) Ekman forcing of sea level (mm month⁻¹). Fingerprints are calculated using surface pressure data and 10-m wind velocities from the NCEP reanalysis (Kalnay et al. 1996).

FIG. 3. As in Fig. 1, but for (top) steric and (bottom) barotropic sea level (mm) in experiment CTRL.

FIG. 4. As in Fig. 1, but for (top) thermosteric and (bottom) halosteric sea level (mm) in experiment CTRL.

FIG. 5. As in Fig. 1, but for experiments (top) WIND and (bottom) BUOY. Regions with climatological sea ice cover exceeding 10% are masked in white as the forcing decomposition was not made for air–ice or ice–ocean fluxes. See the methods in the supplementary material for further description of the flux decomposition in our ocean model sensitivity experiments.

with previous work that has emphasized the dominance of the winds for interannual variability of steric sea level in the North Atlantic (Esselborn and Eden 2001).

Wind forcing tends to dominate barotropic sea level variability, and previous studies have emphasized the importance of the winds in driving baroclinic waves in the tropics (e.g., Vivier et al. 1999; Wyrski 1975). However, the specific role for the winds in driving steric variability associated with extratropical modes of variability such as the NAO and EA is less obvious. One obvious candidate is the contemporaneous steric response to local Ekman pumping that occurs in phase with changes to wind stress curl. To evaluate the importance of this process, we produce composite patterns of Ekman sea level forcing $\partial_t \eta_{\text{Ek}}$ following Gill and Niller (1973) and Cabanes et al. (2006):

$$\partial_t \eta_{\text{Ek}} = \left(\frac{\rho_{\text{surface}} - \langle \rho \rangle}{\rho_0} \right) \times \text{curl} \left(\frac{\boldsymbol{\tau}}{f} \right), \quad (4)$$

where f is the Coriolis parameter, $\boldsymbol{\tau}$ is the wind stress vector estimated from 10-m wind velocities, ρ_{surface} is potential density at the ocean surface, and $\langle \rho \rangle$ is potential density averaged over the depth of the ocean. Note that this definition for $\partial_t \eta_{\text{Ek}}$ is an algebraic simplification of the form presented by Cabanes et al. (2006). Our composites of $\partial_t \eta_{\text{Ek}}$ (Fig. 2) indicate that although the steric response to Ekman pumping may contribute in some locations (e.g., in the subtropical North Atlantic during NAO events), in general the instantaneous Ekman forcing is weak compared to the magnitude of extratropical steric anomalies, even when integrated over several months. This result is consistent with previous work that found a limited role for local Ekman forcing in the North Atlantic (Cabanes et al. 2006) and indicates that other dynamic responses to wind forcing, such as changes in advection or baroclinic adjustments, must be important for steric changes associated with extratropical modes of climate variability.

To evaluate the drivers of variability on interannual time scales (defined as variations in annual means during the period 1960–2009), we calculate the fraction of variance F explained by each process (Fig. 6), which is defined as

$$F = 1 - \frac{\text{Var}(\eta_{\text{signal}} - \eta_{\text{process}})}{\text{Var}(\eta_{\text{signal}})}, \quad (5)$$

where η_{signal} is the time-varying sea level signal and η_{process} is the variability attributable to a specific process. A perfect score ($F = 1$) occurs when η_{signal} and η_{process} are perfectly correlated and have the same variance.

Consistent with previous studies (Fukumori et al. 1998; Vinogradova et al. 2007), we find that DSL variability on interannual time scales is largely steric in nature, although barotropic variations are important in the high latitudes and over shallow marginal seas (Figs. 6a,b). The reduced importance of steric variability over shallow water is a simple consequence of the limited depth over which density anomalies can be integrated. However, offshore steric variability can also play an indirect role in coastal sea level variability by inducing barotropic signals on the continental shelf associated with adjustments to density changes over sloping topography (Bingham and Hughes 2012). Locations with low values of F for both steric and barotropic signals are indicative of compensation between the two processes. Separation of steric variations into thermosteric and halosteric contributions reveals that changes in temperature dominate over most of the oceans, except in the high latitudes (particularly in the Arctic and northernmost Pacific Ocean) and on the continental shelf where changes in salinity are important (Figs. 6c,d). In some regions (e.g., the subtropics in the Atlantic and South Pacific), neither thermosteric or halosteric changes alone are a good measure of the total steric signal. This indicates an important role for density compensation in these regions, a process that occurs as a result of the advection of spiciness anomalies along isopycnals (Köhl 2014).

Using the ocean model sensitivity experiments described in section 2, we also evaluate the relative importance of wind and buoyancy forcings for interannual variations in steric sea level (Figs. 6e,f). Between 20°S and 20°N, almost all interannual variability of steric sea level in CTRL can be attributed to the wind forcing. This result is not surprising given the importance of wind-driven baroclinic waves associated with tropical modes of climate variability (e.g., Wyrski 1975). It is also in agreement with previous work that has linked decadal changes in tropical Pacific sea level with long-term variations in the trade winds (e.g., Merrifield et al. 2012; Merrifield and Maltrud 2011; England et al. 2014). Our experiments also show the importance of wind-driven interannual variability in the extratropics and high latitudes. This is consistent with previous works that have highlighted the role of winds on interannual time scales via their impact on baroclinic waves, quasi-steady Sverdrup balance, and Ekman pumping (e.g., Cabanes et al. 2006; Sturges et al. 1998; Piecuch and Ponte 2011). Buoyancy forcings have a limited impact on sea level in the low latitudes, but are particularly important for steric changes in the North Atlantic linked to variability of the overturning circulation (Figs. 6f,h). This result is consistent with earlier work that has linked low-frequency sea

FIG. 6. Fraction of dynamic sea level variance in CTRL explained by (a) steric and (b) barotropic signals. Fraction of steric sea level variance in CTRL explained by (c) temperature and (d) salinity. Fraction of steric sea level variance in CTRL explained by (e) WIND, (f) BUOY, and (g) WIND + BUOY. (h) Coefficient of determination r^2 for the linear relationship between steric sea level and the AMOC at 26°N in BUOY. Skill scores are calculated using annual mean data covering the duration of the model experiments (1960–2009). In plots (e)–(h), regions with climatological sea ice cover exceeding 10% are masked in white as the forcing decomposition was not made for air–ice or ice–ocean fluxes.

level variability in the Gulf Stream and North Atlantic subpolar gyre with the ocean circulation response to basin-scale modes of buoyancy flux variability (Hakkinen 2001; Hakkinen and Rhines 2004).

We also linearly combine anomalies from WIND and BUOY and calculate their skill in reproducing steric anomalies in CTRL (Fig. 6g). Regions of low skill are

indicative of a substantial role for intrinsic variability and/or nonlinear interactions between the response to buoyancy and wind forcings. Alternatively, regions of low skill could indicate deficiencies in our experimental design such that the combined forcings in WIND and BUOY are not exactly equal to those in CTRL (see methods in the supplementary material for further

FIG. 7. Correlations for the period 1993–2013 calculated using annual mean dynamic sea level anomalies from the DePreSys3 analysis and satellite altimeter data. Hatched areas indicate regions with $|r| \geq 0.433$, the critical value corresponding to $p = 0.05$ for a two-sided test assuming $N - 2$ degrees of freedom, where $N = 21$ is the number of years of data.

details). The regions with the lowest skill ($<20\%$ variance explained) are located in eddy-active regions such as the Southern Ocean and western boundary current extensions, consistent with the locations of substantial intrinsic variability identified by Penduff et al. (2011). The significant role for intrinsic variability in these regions explains why there are regions of low skill in WIND on interannual time scales, despite wind-forcing being the dominant contributor to sea level anomalies associated with modes of climate variability in the same location.

4. Predictability of dynamic sea level

In this section, we evaluate the predictability of DSL variability on seasonal-to-interannual time scales in a state-of-the-art coupled forecast system. Correlations of annual mean DSL anomalies from the DePreSys3 analysis and satellite altimeter data demonstrate that large-scale DSL variability is generally well captured in the state estimate we use to initialize our hindcasts (Fig. 7). As we noted earlier, no direct observations of sea level are used to constrain this analysis, so the good agreement must originate from the assimilation of temperature, salinity, and wind data. Correlations are particularly high in the tropical Pacific and Indian oceans, but generally lower in shallow enclosed seas and regions of substantial intrinsic variability. Although DePreSys3 incorporates an eddy-permitting OGCM, mesoscale variability is unable to fully develop in the observationally constrained analysis. This is because the analysis is strongly determined by the 10-day relaxation

time scale used to restore DePreSys3 to a reconstruction of large-scale temperature and salinity anomalies (Smith and Murphy 2007; Smith et al. 2015). The restoring used in the DePreSys3 analysis does not preclude the development of eddies in individual hindcasts, but it means the mesoscale eddy field is not initialized in hindcasts or predicted as part of the ensemble mean. For this reason, we are implicitly evaluating only the ability of DePreSys3 to predict large-scale patterns of sea level variability.

To assess the skill of forecasts, correlation coefficients from DePreSys3 are compared to those calculated using a simple model of persistence. For each location, a persistence forecast is specified as the mean anomaly in the season or year preceding the initialization date. For example, persistence forecast estimates for November–January are based on the mean anomaly for August–October. For the first season following initialization (i.e., months 1–3), DePreSys3 exhibits high skill for DSL and outperforms persistence over most of the oceans (Fig. 8). Skill in DePreSys3 also exceeds persistence over most of the Pacific and large regions of the Atlantic and Indian Oceans when anomalies are averaged over the first year. Predictions are particularly good in the equatorial and tropical Pacific, which we attribute to skillful predictions of ENSO in DePreSys3 (Dunstone et al. 2016).

At lead times of 3–12 months, there is a strong reduction in the skill of persistence forecasts in the tropical Pacific and Indian Oceans. In contrast, forecasts from DePreSys3 show a less dramatic decline in skill and continue to show significant skill in parts of the tropical Pacific for the second winter since initialization (i.e.,

FIG. 8. Correlations for the period 1960–2013 between the DePreSys3 hindcast ensemble mean and (left) the data assimilating DePreSys3 analysis compared with (right) persistence forecasts: (top)–(bottom) months 1–3, 4–6, 7–9, 10–12, and 13–15, and year 1. Hatching in the left panels indicates areas where correlations from DePreSys3 are greater than those for persistence and $|r| \geq 0.396$, the critical value corresponding to $p = 0.05$ for a two-sided test assuming $N - 2$ degrees of freedom, where $N = 25$ is the number of start dates.

months 13–15). The success of DePreSys3 is a consequence of its ability to more realistically propagate thermocline depth anomalies associated with baroclinic Rossby and Kelvin waves. This is evident from the skill in predictions of the 18°C isotherm depth (Fig. 9). At

extratropical latitudes, although DePreSys3 generally outperforms persistence when anomalies are averaged over the first year, the difference is less distinct. This is likely due to the decreasing speed of baroclinic Rossby waves with latitude that makes it harder to distinguish

FIG. 9. Correlations for the period 1960–2013 between the DePreSys3 hindcast ensemble mean and the data assimilating DePreSys3 analysis calculated for (left) the depth of the 18°C isotherm and surface (center) zonal and (right) meridional wind stress for: (top)–(bottom) months 1–3, 4–6, 7–9, 10–12, and 13–15. Hatching indicates areas where correlations from DePreSys3 are greater than those for persistence and $|r| \geq 0.396$, the critical value corresponding to $p = 0.05$ for a two-sided test assuming $N - 2$ degrees of freedom, where $N = 25$ is the number of start dates.

between slowly propagating features and stationary anomalies (Polkova et al. 2015).

In the previous section, we emphasized the dominant role of the winds for DSL change on seasonal-to-interannual time scales, particularly for DSL anomalies associated with leading modes of climate variability. Given this knowledge, it is interesting to consider two distinct ways it is possible to predict wind-driven variability using a coupled climate model. First, accurate predictions of wind-driven signals can occur via prediction of the atmospheric state and the associated ocean response. On seasonal-to-interannual time scales, this scenario is most relevant when ocean–atmosphere variability is tightly coupled such that surface winds respond to the ocean state and vice versa. Alternatively, predictive skill can emerge by accurately initializing the

ocean to contain the imprint of a previous wind-forcing (e.g., a depth anomaly in the thermocline) that is then propagated by the ocean model dynamics in the absence of skill in the atmosphere (also see Köhl 2014). In this scenario, only the slowly evolving baroclinic contributions to sea level variability will be predictable as the local (e.g., Ekman pumping) and rapid (i.e., barotropic) responses to the simulated wind field will be erroneous.

DePreSys3 has some skill in the predictability of zonal and meridional wind stress at lead times of several months, particularly in the tropical Pacific where air–sea interactions and predictability are strongly coupled (Fig. 9). However, at lead times of 6 months or more, skill in wind stress is much lower than skill for either DSL or 18°C isotherm depth anomalies. Based on these comparisons, it is evident that much of the DSL skill

demonstrated by DePreSys3 must originate from the initialization and propagation of DSL anomalies by the ocean dynamics rather than the prediction of the atmospheric drivers. In the tropics, this translates to accurate predictions of DSL many months in advance, with predictability ultimately limited by the basin transit times for baroclinic adjustments. In the extratropics, although baroclinic wave speeds are much slower, the fraction of variability that can be attributed to remotely forced adjustments is also smaller. This is because local forcing from winds (e.g., Cabanes et al. 2006; Vivier et al. 1999) and rapid barotropic adjustments in response to atmospheric modes of variability (e.g., Fig. 3) are also important. Furthermore, the IB effect is nonnegligible in the extratropics. Based on this evidence, we emphasize that prediction of seasonal-to-interannual sea level variability in the extratropics will be governed to a large extent by the predictability of surface wind stress and modes of atmospheric variability.

5. Summary and conclusions

We have presented an analysis of satellite observations and eddy-permitting model simulations and evaluated the drivers of sea level variability associated with 15 leading modes of climate variability covering the Atlantic, Pacific, Indian, and Southern Oceans. Based on our model experiments, and consistent with previous research (e.g., Köhl 2014; Forget and Ponte 2015; Stammer et al. 2013), we find that seasonal-to-interannual sea level variations in the low latitudes are dominated by the thermosteric response to wind-driven baroclinic waves associated with tropical modes of climate variability. In the extratropics, changes to dynamic sea level associated with atmospheric modes of variability include a substantial barotropic response to wind forcing, particularly over the continental shelf seas. However, wind-driven steric changes are also important for some modes (e.g., NAO, EPNP, and EA; see Table 3). Changes in steric sea level associated with heat and freshwater forcings are generally less important, except on interannual time scales in the northwest Atlantic. In this region, low-frequency sea level variations are associated with buoyancy-forced changes in the Atlantic meridional overturning circulation (AMOC).

Using a state-of-the-art coupled forecast system (DePreSys3), we find skill in seasonal-to-interannual predictions of dynamic sea level that exceeds persistence over large regions of the Pacific, Atlantic, and Indian Oceans. Skill is particularly high in the tropical Indo-Pacific because of the accurate initialization and propagation of thermocline depth anomalies associated with baroclinic waves. Although we also find some skill in the

extratropics that previous authors have linked to slow-moving baroclinic Rossby waves (Polkova et al. 2015), we also emphasize the role of nonsteric signals associated with modes of atmospheric variability. These additional contributions to sea level variability (including rapid barotropic adjustments, local responses to wind forcing, and the IB effect) limit the skill of extratropical sea level forecasts as they require predictability of the atmospheric state, including surface wind and pressure anomalies.

Finally, our ocean model sensitivity experiments also show the important role for intrinsic ocean variability, particularly in boundary currents and the Southern Ocean. The degree of interannual sea level predictability in these regions will thus depend on the time scales on which intrinsic variability becomes chaotic and the accuracy with which the ocean mesoscale can be initialized.

Acknowledgments. This work was supported by the Joint DECC/Defra Met Office Hadley Centre Climate Programme (GA01101) and the EU FP7 SPECS project. We thank three anonymous reviewers for useful comments that helped us to improve an earlier version of this manuscript.

REFERENCES

- Ablain, M., and Coauthors, 2015: Improved sea level record over the satellite altimetry era (1993–2010) from the Climate Change Initiative Project. *Ocean Sci.*, **11**, 67–82, doi:10.5194/os-11-67-2015.
- Antonov, J. I., S. Levitus, and T. P. Boyer, 2002: Steric sea level variations during 1957–1994: Importance of salinity. *J. Geophys. Res.*, **107**, 8013, doi:10.1029/2001JC000964.
- Ashok, K., S. K. Behera, S. A. Rao, H. Weng, and T. Yamagata, 2007: El Niño Modoki and its possible teleconnection. *J. Geophys. Res.*, **112**, C11, doi:10.1029/2006JC003798.
- Barnston, A. G., and R. E. Livezey, 1987: Classification, seasonality and persistence of low-frequency atmospheric circulation patterns. *Mon. Wea. Rev.*, **115**, 1083–1126, doi:10.1175/1520-0493(1987)115<1083:CSAPOL>2.0.CO;2.
- Becker, M., B. Meyssignac, C. Letetrel, W. Llovel, A. Cazenave, and T. Delcroix, 2012: Sea level variations at tropical Pacific islands since 1950. *Global Planet. Change*, **80–81**, 85–98, doi:10.1016/j.gloplacha.2011.09.004.
- Bingham, R. J., and C. W. Hughes, 2012: Local diagnostics to estimate density-induced sea level variations over topography and along coastlines. *J. Geophys. Res.*, **117**, C01013, doi:10.1029/2011JC007276.
- Cabanes, C., T. Huck, and A. Colin de Verdière, 2006: Contributions of wind forcing and surface heating to interannual sea level variations in the Atlantic Ocean. *J. Phys. Oceanogr.*, **36**, 1739–1750, doi:10.1175/JPO2935.1.
- Cazenave, A., and W. Llovel, 2010: Contemporary sea level rise. *Annu. Rev. Mar. Sci.*, **2**, 145–173, doi:10.1146/annurev-marine-120308-081105.
- Chelton, D. B., and R. E. Davis, 1982: Monthly mean sea-level variability along the west coast of North America. *J. Phys.*

- Oceanogr.*, **12**, 757–784, doi:[10.1175/1520-0485\(1982\)012<0757:MMSLVA>2.0.CO;2](https://doi.org/10.1175/1520-0485(1982)012<0757:MMSLVA>2.0.CO;2).
- , M. G. Schlax, R. M. Samelson, and R. A. de Szoeke, 2007: Global observations of large oceanic eddies. *Geophys. Res. Lett.*, **34**, L15606, doi:[10.1029/2007GL030812](https://doi.org/10.1029/2007GL030812).
- Chiang, J. C. H., and D. J. Vimont, 2004: Analogous meridional modes of atmosphere–ocean variability in the tropical Pacific and tropical Atlantic. *J. Climate*, **17**, 4143–4158, doi:[10.1175/JCLI4953.1](https://doi.org/10.1175/JCLI4953.1).
- Chowdhury, M. R., P.-S. Chu, T. Schroeder, and N. Colasacco, 2007: Seasonal sea-level forecasts by canonical correlation analysis—An operational scheme for the US-affiliated Pacific Islands. *Int. J. Climatol.*, **27**, 1389–1402, doi:[10.1002/joc.1474](https://doi.org/10.1002/joc.1474).
- , —, and C. C. Guard, 2014: An improved sea level forecasting scheme for hazards management in the US-affiliated Pacific Islands. *Int. J. Climatol.*, **34**, 2320–2329, doi:[10.1002/joc.3841](https://doi.org/10.1002/joc.3841).
- Church, J. A., and N. J. White, 2011: Sea-level rise from the late 19th to the early 21st century. *Surv. Geophys.*, **32**, 585–602, doi:[10.1007/s10712-011-9119-1](https://doi.org/10.1007/s10712-011-9119-1).
- Cummins, P. F., and G. S. E. Lagerloef, 2004: Wind-driven interannual variability over the northeast Pacific Ocean. *Deep-Sea Res. I*, **51**, 2105–2121, doi:[10.1016/j.dsr.2004.08.004](https://doi.org/10.1016/j.dsr.2004.08.004).
- Dee, D. P., and Coauthors, 2011: The ERA-Interim reanalysis: Configuration and performance of the data assimilation system. *Quart. J. Roy. Meteor. Soc.*, **137**, 553–507, doi:[10.1002/qj.828](https://doi.org/10.1002/qj.828).
- Dunstone, N., D. Smith, A. Scaife, L. Hermanson, R. Ease, N. Robinson, M. Andrews, and J. Knight, 2016: Skillful predictions of the winter North Atlantic oscillation one year ahead. *Nature Geosci.*, in press.
- Enfield, D. B., and J. S. Allen, 1980: On the structure and dynamics of monthly mean sea level anomalies along the Pacific coast of North and South America. *J. Phys. Oceanogr.*, **10**, 557–578, doi:[10.1175/1520-0485\(1980\)010<0557:OTSADO>2.0.CO;2](https://doi.org/10.1175/1520-0485(1980)010<0557:OTSADO>2.0.CO;2).
- England, M. H., and Coauthors, 2014: Recent intensification of wind-driven circulation in the Pacific and the ongoing warming hiatus. *Nat. Climate Change*, **4**, 222–227, doi:[10.1038/nclimate2106](https://doi.org/10.1038/nclimate2106).
- Esselborn, S., and C. Eden, 2001: Sea surface height changes in the North Atlantic Ocean related to the North Atlantic Oscillation. *Geophys. Res. Lett.*, **28**, 3473–3476, doi:[10.1029/2001GL012863](https://doi.org/10.1029/2001GL012863).
- Fagherazzi, S., G. Fossler, L. D’Alpaos, and P. D’Odorico, 2005: Climatic oscillations influence the flooding of Venice. *Geophys. Res. Lett.*, **32**, L19710, doi:[10.1029/2005GL023758](https://doi.org/10.1029/2005GL023758).
- Ferry, N., G. Reverdin, and A. Oschlies, 2000: Seasonal sea surface height variability in the North Atlantic Ocean. *J. Geophys. Res.*, **105**, 6307–6326, doi:[10.1029/1999JC900296](https://doi.org/10.1029/1999JC900296).
- Forget, G., and R. M. Ponte, 2015: The partition of regional sea level variability. *Prog. Oceanogr.*, **137**, 173–195, doi:[10.1016/j.pocean.2015.06.002](https://doi.org/10.1016/j.pocean.2015.06.002).
- Frankignoul, C., P. Müller, and E. Zorita, 1997: A simple model of the decadal response of the ocean to stochastic wind forcing. *J. Phys. Oceanogr.*, **27**, 1533–1546, doi:[10.1175/1520-0485\(1997\)027<1533:ASMOTD>2.0.CO;2](https://doi.org/10.1175/1520-0485(1997)027<1533:ASMOTD>2.0.CO;2).
- Fukumori, I., R. Raghunath, and L.-L. Fu, 1998: Nature of global large-scale sea level variability in relation to atmospheric forcing: A modeling study. *J. Geophys. Res.*, **103**, 5493–5512, doi:[10.1029/97JC02907](https://doi.org/10.1029/97JC02907).
- Gill, A. E., and P. P. Niller, 1973: The theory of the seasonal variability in the ocean. *Deep-Sea Res. Oceanogr. Abstr.*, **20**, 141–177, doi:[10.1016/0011-7471\(73\)90049-1](https://doi.org/10.1016/0011-7471(73)90049-1).
- Goddard, P. B., J. Yin, S. M. Griffies, and S. Zhang, 2015: An extreme event of sea-level rise along the Northeast coast of North America in 2009–2010. *Nat. Commun.*, **6**, 6346, doi:[10.1038/ncomms7346](https://doi.org/10.1038/ncomms7346).
- Häkkinen, S., 2001: Variability in sea surface height: A qualitative measure for the meridional overturning in the North Atlantic. *J. Geophys. Res.*, **106**, 13 837–13 848, doi:[10.1029/1999JC000155](https://doi.org/10.1029/1999JC000155).
- , and P. B. Rhines, 2004: Decline of subpolar North Atlantic circulation during the 1990s. *Science*, **304**, 555–559, doi:[10.1126/science.1094917](https://doi.org/10.1126/science.1094917).
- Hunke, E. C., and W. H. Lipscomb, 2010: CICE: The Los Alamos Sea Ice Model documentation and software user’s manual, version 4.1. Doc. LA-CC-06-012, 76 pp. [Available online at http://csdms.colorado.edu/w/images/CICE_documentation_and_software_user's_manual.pdf.]
- Jacobs, G. A., H. E. Hurlburt, J. C. Kindle, E. J. Metzger, J. L. Mitchell, W. J. Teague, and A. J. Wallcraft, 1994: Decade-scale trans-Pacific propagation and warming effects of an El-Niño anomaly. *Nature*, **370**, 360–363, doi:[10.1038/370360a0](https://doi.org/10.1038/370360a0).
- Kalnay, E., and Coauthors, 1996: The NCEP/NCAR 40-Year Reanalysis Project. *Bull. Amer. Meteor. Soc.*, **77**, 437–471, doi:[10.1175/1520-0477\(1996\)077<0437:TNYRP>2.0.CO;2](https://doi.org/10.1175/1520-0477(1996)077<0437:TNYRP>2.0.CO;2).
- Kelly, K. A., M. J. Caruso, and J. A. Austin, 1993: Wind-forced variations in sea surface height in the northeast Pacific Ocean. *J. Phys. Oceanogr.*, **23**, 2392–2411, doi:[10.1175/1520-0485\(1993\)023<2392:WFWISS>2.0.CO;2](https://doi.org/10.1175/1520-0485(1993)023<2392:WFWISS>2.0.CO;2).
- Knight, J. F., and Coauthors, 2014: Predictions of climate several years ahead using an improved decadal prediction system. *J. Climate*, **27**, 7550–7567, doi:[10.1175/JCLI-D-14-00069.1](https://doi.org/10.1175/JCLI-D-14-00069.1).
- Köhl, A., 2014: Detecting processes contributing to interannual halosteric and thermosteric sea level variability. *J. Climate*, **27**, 2417–2426, doi:[10.1175/JCLI-D-13-00412.1](https://doi.org/10.1175/JCLI-D-13-00412.1).
- Large, W. G., and S. G. Yeager, 2004: Diurnal to decadal global forcing for ocean and sea-ice models: The data sets and flux climatologies. NCAR Tech. Note NCAR/TN-460+STR, 105 pp., doi:[10.5065/D6KK98Q6](https://doi.org/10.5065/D6KK98Q6).
- , and —, 2009: The global climatology of an interannually varying air–sea flux data set. *Climate Dyn.*, **33**, 341–364, doi:[10.1007/s00382-008-0441-3](https://doi.org/10.1007/s00382-008-0441-3).
- Lutz, K., J. Rathmann, and J. Jacobbeit, 2013: Classification of warm and cold water events in the eastern tropical Atlantic Ocean. *Atmos. Sci. Lett.*, **14**, 102–106, doi:[10.1002/asl2.424](https://doi.org/10.1002/asl2.424).
- MacLachlan, C., and Coauthors, 2015: Global Seasonal forecast system version 5 (GloSea5): A high-resolution seasonal forecast system. *Quart. J. Roy. Meteor. Soc.*, **141**, 1072–1084, doi:[10.1002/qj.2396](https://doi.org/10.1002/qj.2396).
- Madec, G., 2008: NEMO ocean engine, version 3.0. IPSL Pole Modelization Note 27, 209 pp. [Available online at http://www.nemo-ocean.eu/content/download/5302/31828/file/NEMO_book.pdf.]
- McGregor, S., N. J. Holbrook, and S. B. Power, 2007: Interdecadal sea surface temperature variability in the equatorial Pacific Ocean. Part I: The role of off-equatorial wind stresses and oceanic Rossby waves. *J. Climate*, **20**, 2643–2658, doi:[10.1175/JCLI4145.1](https://doi.org/10.1175/JCLI4145.1).
- McIntosh, P. C., J. A. Church, E. R. Miles, K. Ridgway, and C. M. Spillman, 2015: Seasonal coastal sea level prediction using a dynamical model. *Geophys. Res. Lett.*, **42**, 6747–6753, doi:[10.1002/2015GL065091](https://doi.org/10.1002/2015GL065091).
- Megann, A., and Coauthors, 2014: GO5.0: The joint NERC-Met Office NEMO global ocean model for use in coupled and

- forced applications. *Geosci. Model Dev.*, **7**, 1069–1092, doi:10.5194/gmd-7-1069-2014.
- Meinshausen, M., and Coauthors, 2011: The RCP greenhouse gas concentrations and their extensions from 1765 to 2300. *Climatic Change*, **109**, 213–241, doi:10.1007/s10584-011-0156-z.
- Menéndez, M., and P. L. Woodworth, 2010: Changes in extreme high water levels based on a quasi-global tide-gauge data set. *J. Geophys. Res.*, **115**, C10011, doi:10.1029/2009JC005997.
- Merrifield, M. A., and M. E. Maltrud, 2011: Regional sea level trends due to a Pacific trade wind intensification. *Geophys. Res. Lett.*, **38**, L21605, doi:10.1029/2011GL049576.
- , P. R. Thompson, and M. Lander, 2012: Multidecadal sea level anomalies and trends in the western tropical Pacific. *Geophys. Res. Lett.*, **39**, L13602, doi:10.1029/2012GL052032.
- Miles, E. R., C. M. Spillman, J. A. Church, and P. C. McIntosh, 2014: Seasonal prediction of global sea level anomalies using an ocean–atmosphere dynamical model. *Climate Dyn.*, **43**, 2131–2145, doi:10.1007/s00382-013-2039-7.
- Mo, K. C., 2000: Relationships between low-frequency variability in the Southern Hemisphere and sea surface temperature anomalies. *J. Climate*, **13**, 3599–3610, doi:10.1175/1520-0442(2000)013<3599:RBLFVI>2.0.CO;2.
- Nicholls, R. J., and A. Cazenave, 2010: Sea-level rise and its impact on coastal zones. *Science*, **328**, 1517–1520, doi:10.1126/science.1185782.
- , F. M. J. Hoozemans, and M. Marchand, 1999: Increasing flood risk and wetland losses due to global sea-level rise: Regional and global analyses. *Global Environ. Change*, **9**, S69–S87, doi:10.1016/S0959-3780(99)00019-9.
- Penduff, T., M. Juza, L. Brodeau, G. C. Smith, B. Barnier, J.-M. Molines, A.-M. Treguier, and G. Madec, 2010: Impact of global ocean model resolution on sea-level variability with emphasis on interannual time scales. *Ocean Sci.*, **6**, 269–284, doi:10.5194/os-6-269-2010.
- , B. Barnier, J. Zika, W. K. Dewar, A.-M. Treguier, J.-M. Molines, and N. Audiffren, 2011: Sea level expression of intrinsic and forced ocean variabilities at interannual time scales. *J. Climate*, **24**, 5652–5670, doi:10.1175/JCLI-D-11-00077.1.
- Piecuch, C. G., and R. M. Ponte, 2011: Mechanisms of interannual steric sea level variability. *Geophys. Res. Lett.*, **38**, L15605, doi:10.1029/2011GL048440.
- , K. J. Quinn, and R. M. Ponte, 2013: Satellite-derived interannual ocean bottom pressure variability and its relation to sea level. *Geophys. Res. Lett.*, **40**, 3106–3110, doi:10.1002/grl.50549.
- Polkova, I., A. Köhl, and D. Stammer, 2014: Impact of initialization procedures on the predictive skill of a coupled ocean–atmosphere model. *Climate Dyn.*, **42**, 3151–3169, doi:10.1007/s00382-013-1969-4.
- , —, and —, 2015: Predictive skill for regional interannual steric sea level and mechanisms for predictability. *J. Climate*, **28**, 7407–7419, doi:10.1175/JCLI-D-14-00811.1.
- Qiu, B., 2002: Large-scale variability in the midlatitude subtropical and subpolar North Pacific Ocean: Observations and causes. *J. Phys. Oceanogr.*, **32**, 353–375, doi:10.1175/1520-0485(2002)032<0353:LSVTIM>2.0.CO;2.
- , and S. Chen, 2006: Decadal variability in the large-scale sea surface height field of the South Pacific Ocean: Observations and causes. *J. Phys. Oceanogr.*, **36**, 1751–1762, doi:10.1175/JPO2943.1.
- , and —, 2010: Interannual-to-decadal variability in the bifurcation of the North Equatorial Current off the Philippines. *J. Phys. Oceanogr.*, **40**, 2525–2538, doi:10.1175/2010JPO4462.1.
- Rayner, N. A., D. E. Parker, E. B. Horton, C. K. Folland, L. V. Alexander, D. P. Rowell, E. C. Kent, and A. Kaplan, 2003: Global analyses of sea surface temperature, sea ice, and night marine air temperature since the late nineteenth century. *J. Geophys. Res.*, **108**, 4407, doi:10.1029/2002JD002670.
- Roberts, C. D., and Coauthors, 2013: Atmosphere drives recent interannual variability of the Atlantic meridional overturning circulation at 26.5°N. *Geophys. Res. Lett.*, **40**, 5164–5170, doi:10.1002/grl.50930.
- Saji, N. H., B. N. Goswami, P. N. Vinayachandran, and T. Yamagata, 1999: A dipole mode in the tropical Indian Ocean. *Nature*, **401** (6751), 360–363.
- Smith, D. M., and J. M. Murphy, 2007: An objective ocean temperature and salinity analysis using covariances from a global climate model. *J. Geophys. Res.*, **112**, C02022, doi:10.1029/2006JD007574.
- , R. Eade, and H. Pohlmann, 2013: A comparison of full-field and anomaly initialization for seasonal to decadal climate prediction. *Climate Dyn.*, **41**, 3325–3338, doi:10.1007/s00382-013-1683-2.
- , and Coauthors, 2015: Earth’s energy imbalance since 1960 in observations and CMIP5 models. *Geophys. Res. Lett.*, **42**, 1205–1213, doi:10.1002/2014GL062669.
- Stammer, D., 1997: Steric and wind-induced changes in TOPEX/POSEIDON large-scale sea surface topography observations. *J. Geophys. Res.*, **102**, 20 987–21 009, doi:10.1029/97JC01475.
- , A. Cazenave, R. M. Ponte, and M. E. Tamisiea, 2013: Causes for contemporary regional sea level changes. *Annu. Rev. Mar. Sci.*, **5**, 21–46, doi:10.1146/annurev-marine-121211-172406.
- Sturges, W., and B. G. Hong, 1995: Wind forcing of the Atlantic thermocline along 32°N at low frequencies. *J. Phys. Oceanogr.*, **25**, 1706–1715, doi:10.1175/1520-0485(1995)025<1706:WFOTAT>2.0.CO;2.
- , —, and A. J. Clarke, 1998: Decadal wind forcing of the North Atlantic subtropical gyre. *J. Phys. Oceanogr.*, **28**, 659–668, doi:10.1175/1520-0485(1998)028<0659:DWFOTN>2.0.CO;2.
- Taylor, K. E., R. J. Stouffer, and G. A. Meehl, 2012: An overview of CMIP5 and the experiment design. *Bull. Amer. Meteor. Soc.*, **93**, 485–498, doi:10.1175/BAMS-D-11-00094.1.
- Timmermann, A., S. McGregor, and F.-F. Jin, 2010: Wind effects on past and future regional sea level trends in the southern Indo-Pacific. *J. Climate*, **23**, 4429–4437, doi:10.1175/2010JCLI3519.1.
- Trenberth, K. E., 1997: The definition of El Niño. *Bull. Amer. Meteor. Soc.*, **78**, 2771–2777, doi:10.1175/1520-0477(1997)078<2771:TDOENO>2.0.CO;2.
- , and J. W. Hurrell, 1994: Decadal atmosphere–ocean variations in the Pacific. *Climate Dyn.*, **9**, 303–319, doi:10.1007/BF00204745.
- Tsimplis, M. N., A. G. Shaw, R. A. Flather, and D. K. Woolf, 2006: The influence of the North Atlantic Oscillation on the sea-level around the northern European coasts reconsidered: The thermosteric effects. *Philos. Trans. Roy. Soc. London*, **A364**, 845–856, doi:10.1098/rsta.2006.1740.
- Vinogradova, N. T., R. M. Ponte, and D. Stammer, 2007: Relation between sea level and bottom pressure and the vertical dependence of oceanic variability. *Geophys. Res. Lett.*, **34**, L03608, doi:10.1029/2006GL028588.
- Vivier, F., K. A. Kelly, and L. Thompson, 1999: Contributions of wind forcing, waves, and surface heating to sea surface height observations in the Pacific Ocean. *J. Geophys. Res.*, **104**, 20 767–20 788, doi:10.1029/1999JC900096.

- Williams, K., and Coauthors, 2015: The Met Office Global Coupled model 2.0 (GC2) configuration. *Geosci. Model Dev.*, **8**, 1509–1524, doi:[10.5194/gmd-8-1509-2015](https://doi.org/10.5194/gmd-8-1509-2015).
- Wong, P. P., I. J. Losada, J.-P. Gattuso, J. Hinkel, A. Khattabi, K. L. McInnes, Y. Saito, and A. Sallenger, 2014: Coastal systems and low-lying areas. *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects*. C. B. Field et al., Eds., Cambridge University Press, 361–410.
- Woodworth, P. L., R. A. Flather, J. A. Williams, S. L. Wakelin, and S. Jevrejeva, 2007: The dependence of UK extreme sea levels and storm surges on the North Atlantic Oscillation. *Cont. Shelf Res.*, **27**, 935–946, doi:[10.1016/j.csr.2006.12.007](https://doi.org/10.1016/j.csr.2006.12.007).
- Woolf, D. K., A. G. P. Shaw, and M. N. Tsimplis, 2003: The influence of the North Atlantic Oscillation on sea-level variability in the North Atlantic region. *J. Atmos. Ocean Sci.*, **9**, 145–167, doi:[10.1080/10236730310001633803](https://doi.org/10.1080/10236730310001633803).
- Wyrtki, K., 1975: El Niño—The dynamic response of the equatorial Pacific ocean to atmospheric forcing. *J. Phys. Oceanogr.*, **5**, 572–584, doi:[10.1175/1520-0485\(1975\)005<0572:ENTDRO>2.0.CO;2](https://doi.org/10.1175/1520-0485(1975)005<0572:ENTDRO>2.0.CO;2).
- Xue, Y., and A. Leetmaa, 2000: Forecasts of tropical Pacific SST and sea level using a Markov model. *Geophys. Res. Lett.*, **27**, 2701–2704, doi:[10.1029/1999GL011107](https://doi.org/10.1029/1999GL011107).
- Yan, Z., M. N. Tsimplis, and D. Woolf, 2004: Analysis of the relationship between the North Atlantic oscillation and sea-level changes in northwest Europe. *Int. J. Climatol.*, **24**, 743–758, doi:[10.1002/joc.1035](https://doi.org/10.1002/joc.1035).
- Zhang, X., and J. A. Church, 2012: Sea level trends, interannual and decadal variability in the Pacific Ocean. *Geophys. Res. Lett.*, **39**, L21701, doi:[10.1029/2012GL053240](https://doi.org/10.1029/2012GL053240).