Wave Climate Patterns from Spatial Tracking of Global Long-Term Ocean Wave Spectra

HAOYU JIANG

College of Marine Science and Technology, China University of Geosciences, Wuhan, and Laboratory for Regional Oceanography and Numerical Modeling, Qingdao National Laboratory for Marine Science and Technology, Qingdao, and Shenzhen Research Institute, China University of Geosciences, Shenzhen, China

(Manuscript received 27 September 2019, in final form 20 January 2020)

ABSTRACT

Long-term wave spectral statistics can provide a better description of wave climate than integrated wave parameters because several wave climate systems (WCSs) generated by different wind climate systems can coexist at the same location. In this study, global wave climate patterns are presented by spatially tracking point-wise long-term wave spectra (probability density distributions of wave spectral partitions) from a WAVEWATCH III hindcast, providing new insights into global wave climate. Tens of well-defined WCSs, which are generated in different source regions by different wind systems, including prevailing westerlies, polar easterlies, trade winds, and monsoons, were identified. These WCSs are independent of each other because wave systems from different origins travel independently. The spatial distributions of these WCSs can illustrate the entire life cycle of ocean waves, from being generated as dominant wind-seas to becoming less dominant swells in far fields, from a climatic point of view. The mean wave directions in WCS patterns, especially those in westerlies-generated WCSs, are generally in agreement with the great circles on Earth's surface, which display the propagation routes of ocean swells.

1. Introduction

Wind-generated surface gravity waves (simply called waves henceforth) are a fundamental and ubiquitous phenomenon at the air–sea interface. They have an impact on many aspects of human life (e.g., seafaring, coastal engineering, port operations) and play an important role in many geophysical processes (e.g., momentum exchange at the air–sea boundary layer). Studies of global and regional wave climate are important for both social and scientific perspectives and thus are widely conducted using the data obtained from various sources including numerical model outputs and observations from satellites and voluntary observing ships (e.g., Young 1999; Cox and Swail 2001; Chen et al. 2002; Gulev et al. 2003; Gulev and Grigorieva 2004; Alves 2006; Hemer et al. 2010, 2013; Semedo et al. 2011, 2013; Young et al. 2011, 2012; Fan et al. 2012, 2013, 2014; Aarnes et al. 2015).

Most of these studies focused on important integrated wave parameters such as significant wave height H_s , mean wave period T_m , and mean wave direction θ_m . Because these parameters are not sufficient to describe mixed sea states with both wind-seas and swells, the H_s , T_m , and θ_m values of wind-seas (H_s^w , T_m^w , and θ_m^w , respectively) and swells (H_s^s , T_m^s , and θ_m^s , respectively) are also computed separately in some of the aforementioned studies. These widely used integrated wave parameters are very useful for describing wave climate. However, they also have some disadvantages. For example, sometimes it might be difficult to find a clear objective criterion to distinguish “old” wind-seas and “young” swells because their distinction is often not obvious (Portilla et al. 2009). More importantly, multiple swell systems can coexist at the same time and location in the open ocean (e.g., Portilla et al. 2009), leading to “crossing” swells (e.g., Jiang et al. 2017) or “opposing” swells. A sea state with $H_s^s = 2$ m, $T_m^s = 12$ s, and $\theta_m^s = 90^\circ$ might be the superposition of two swell systems with $H_s^{s1} = H_s^{s2} = \sqrt{2}$ m, $T_m^{s1} = 10$ s, $\theta_m^{s1} = 45^\circ$, and $\theta_m^{s2} = 135^\circ$. In a long-term-averaged ocean wave spectrum at a given location, multiple wind-sea climate systems can also coexist. For example, monsoons can generate two wind-sea climate systems with

Supplemental information related to this paper is available at the Journals Online website: <https://doi.org/10.1175/JCLI-D-19-0729.s1>.

Corresponding author: Haoyu Jiang, haoyujiang@cug.edu.cn

different characteristics at the same location. When the energy of one wave (climate) system is higher than other wave (climate) systems at a location, the bulk wave parameters (H_s , T_m , and θ_m) can only represent the system with the highest energy. Because multimodal sea states and wave climates are very common in the open ocean, the bulk integrated wave parameters even after the separation of wind-sea climate and swell climate can still provide only a limited (and sometimes misleading) description of the wave climate.

The wave climate system (WCS) generated by the same wind climate system (e.g., westerlies, trade winds, and polar easterlies in both hemispheres) have different properties from those generated by other wind climate systems. Distinguishing different WCSs generated by different wind climate systems is helpful to have a more comprehensive description of the global wave climate. [Alves \(2006\)](#) presented a technique of simulating the wave climates generated by the wind in different regions separately, which can isolate the contribution of different WCSs to the global wave climate to some extent. However, the method presented by [Alves \(2006\)](#) has four small shortcomings: 1) The way in which the swell generation areas are separated is subjective to some extent. 2) The presented method cannot separate different WCSs generated in the same area (i.e., there might be more than one WCS generated in the same region, e.g., in the monsoon regions). 3) Some physical processes such as swell dissipation are not taken into account. 4) Separating wind forces in different regions may reduce the fetches of some developing waves, which induces error in mean H_s . Notwithstanding these problems, the work of [Alves \(2006\)](#) successfully provides some patterns associated with WCSs generated by different wind climate systems.

Another way of isolating the contribution of different wave systems is to use the wave spectral information. [Echevarria et al. \(2019\)](#) presented the global spectral wave climate using a relatively coarse resolution of $10^\circ \times 10^\circ$, and they applied the empirical orthogonal function (EOF) to these wave spectra to obtain the seasonal cycle of the spectral wave climate. The long-term wave spectra are directly analyzed in [Echevarria et al. \(2019\)](#), but a major shortcoming of their work is the adoption of a coarse resolution. [Portilla \(2018\)](#) developed an atlas of global spectral wave climate (GLOSWAC) based on the concept of long-term wave spectra and their spectral partitioning. In GLOSWAC, the long-term wave spectrum at a certain geographical location is “partitioned” into different wave systems to classify waves originating from the different wind climate systems. Hereafter, the point-wise wave systems obtained by partitioning long-term wave spectra are

referred to as climatic wave partitions (CWPs), and a WCS consists of CWPs at different spatial locations originating from the same wind climate. The mean H_s , T_m , θ_m , and some other characteristic parameters can be computed separately for each CWP at any point in the ocean.

The mean H_s , T_m , and θ_m for different CWPs can provide a better description of wave climate than bulk integrated parameters (H_s , T_m , θ_m , H_s^w , T_m^w , θ_m^w , H_s^s , T_m^s , θ_m^s) ([Portilla 2018](#)). The analysis of GLOSWAC is preliminary point-wise, whereas the global distributions of wave parameters for different CWPs are helpful for a better understanding of the wave climate. This is particularly helpful for the understanding of the swell climate in the region where several CWPs coexist, because the CWPs with relatively low energy (mainly swell-related CWPs) cannot be represented by bulk wave parameters in this situation. The long-term wave spectra are usually similar at two locations close to each other ([Portilla et al. 2015](#)), which means that the spatial gradients of mean H_s , T_m , and θ_m for the CWPs originating from the same source should also be low in the ocean (except at some discontinuous edges such as islands). However, such spatial coherence of two CWPs at two geographical locations close to each other is currently not guaranteed because of the coexistence of several CWPs at the same location, making it difficult to present the global distributions of WCSs. [Jiang \(2019\)](#) developed a technique to track the wave partitions generated by the same meteorological event (e.g., a storm) from model outputs. This technique is based on the feature that the spatial gradients of H_s , T_m , and θ_m for the wave partitions generated by the same meteorological event are small. Therefore, this technique should also work for spatially tracking CWPs, and the aim of this study is to present a global view of WCSs using this technique.

The rest of the paper is organized as follows. [Section 2](#) introduces the numerical wave model data used in this study and the method of tracking WCSs from CWPs. In [section 3](#), an example is given to demonstrate the limitation of using bulk integrated wave parameters to describe wave climate. The spatial distributions of the tracked WCSs are displayed and discussed in [section 4](#), followed by a summary in [section 5](#).

2. Data and methods

a. Wave hindcast data

Although measured wave spectra are available from the wave mode of spaceborne synthetic aperture radars, the data quality of these wave spectra is still not sufficient for studying wave climate ([Jiang et al. 2017](#)). Therefore, long-term global-scale spectral information is only available

from numerical wave models. The data used in this study are the spectral partition data obtained from the Integrated Ocean Waves for Geophysical and other Applications (IOWAGA) dataset from 1990 to 2015, a hindcast of WAVEWATCH-III (WAVEWATCH III Development Group 2016) using the physical parameterizations of Ardhuin et al. (2010) forced by the global 10-m wind data from the Climate Forecast System Reanalysis (CFSR). The spatial–temporal resolution of the model output used here is $0.5^\circ \times 0.5^\circ \times 3$ h. The directional spectrum computed in the model is spaced in 32 frequency bins increasing exponentially from 0.038 to 0.72 Hz, and 24 directional bins with 15° spacing. The spectrum at each grid point is partitioned into at most one wind-sea partition and up to five swell partitions using the method developed by Hanson and Phillips (2001). The partitioned H_s , T_m , and θ_m , which are the data used in this study, are computed for each spectral partition. In this WAVEWATCH-III setup, an explicit swell dissipation source term is included based on the space-borne observations of Ardhuin et al. (2009). The data show good agreement with observations from both buoys and remote sensing radar altimeters with respect to both overall H_s (Rascle and Ardhuin 2013; Stopa et al. 2016) and partitioned wave information, especially for ocean swells (e.g., Delpey et al. 2010; Stopa et al. 2016; Jiang et al. 2016a). The data are downloaded from the ftp server of IFREMER (ftp.ifremer.fr) where more detailed information about this dataset is also available (Rascle and Ardhuin 2013).

b. Probability density distributions of spectral partitions

The global partitioned H_s , T_m , and θ_m data of different spectral partitions from the IOWAGA dataset are organized in different seasons here such as: December–February (DJF), March–May (MAM), June–August (JJA), and September–November (SON). For each spectral partition at a given geographical location, the partitioned T_m and θ_m correspond to a specific position in the two-dimensional frequency–direction space. The frequency–direction positions of all wave partitions in the same location and season are collected. The frequency–direction space is again discretized into the aforementioned 32 frequency bins and 24 directional bins. The number of occurrences for spectral partitions in each frequency–direction bin is then counted without differentiating wind-sea and swell systems, and the result can be regarded as the probability density distributions of spectral partitions (PDSs) in frequency–direction space for different geographical locations and seasons. After that, the spectral partitioning algorithm is again applied on the smoothed PDSs to obtain the CWP. This method of using

PDSs to describe the spectral wave climate in a given geographical location is developed by Portilla et al. (2015) and has been proven to be an effective way to separate CWPs from different wind climate systems at any given location.

c. Integrated wave parameters for CWPs

For a given CWP labeled as CWP K at a certain geographical location, its mean H_s , T_m , and θ_m are calculated as follows:

$$\overline{H_s^K} = \sqrt{\sum_{i \in K} (H_{s,i}^K)^2 / N}, \quad (1)$$

$$\overline{T_m^K} = \left[\sum_{i \in K} T_{m,i}^K \times (H_{s,i}^K)^2 \right] / \left[\sum_{i \in K} (H_{s,i}^K)^2 \right], \quad (2)$$

$$\overline{\theta_m^K} = \left[\sum_{i \in K} \theta_{m,i}^K \times (H_{s,i}^K)^2 \right] / \left[\sum_{i \in K} (H_{s,i}^K)^2 \right], \quad (3)$$

where N is the length of 3-hourly time series in the corresponding season; $H_{s,i}^K$, $T_{p,i}^K$, and $\theta_{p,i}^K$ are the partitioned H_s , T_m , and θ_m of partition i (belonging to CWP K), respectively. Equation (3) is a weighted average of many directions that may range from 0° to 360° . Therefore, the values of partitioned θ_m in the same CWP, which are usually close to each other, are transformed (by subtracting or adding 360°) into a range where the difference between the maximum and minimum is less than 180° to solve the problem of averaging angles crossing $0^\circ/360^\circ$.

An alternative method for calculating mean H_s , T_m , and θ_m of different CWPs is to apply the spectral partitioning algorithm directly to the mean directional wave spectrum. However, the mean wave spectrum at a given location is smoother than the corresponding wave PDS, as pointed out by Portilla et al. (2015). Not all single CWPs are recognizable in the mean directional wave spectrum, especially those with relatively low energy. For example, Jiang and Mu (2019) show that the signature of the California low-level coastal jets (CLCJs) cannot be observed in the mean spectrum in the equatorial region but is well defined in the PDS. Therefore, the author recommends using the aforementioned method based on the partitioned PDS, instead of the mean directional wave spectrum, to calculate the mean H_s , T_m , and θ_m for each CWP.

d. Spatial tracking of WCSs

The spatial gradients of mean H_s , T_m , and θ_m are rather limited for the same WCS (CWPs originating from the same wind climate system) at two close geographical locations. This feature can be used to spatially

track the same WCS at different locations in the ocean. The spatial tracking algorithm used, presented by Jiang (2019), is originally designed for spatially tracking the wave event generated by the same meteorological event (e.g., a storm) from spectral partitions at different geographical locations. Meanwhile, this method is used in this study to spatially track the WCS generated by the same wind climate system from CWP at different geographical locations. The procedure of this algorithm is briefly described as follows:

1) Start from any CWP at any location. Search all its spatial neighbors one by one. For each neighboring point, find the CWP with the minimum spatial gradient of empirical spectral distance. The spatial gradient of empirical spectral distance between two CWPs at two different points, X and Y , is defined as

$$\delta = \frac{1}{d} \sqrt{\left(\frac{\overline{H_s^X} - \overline{H_s^Y}}{H_a}\right)^2 + \left(\frac{\overline{T_m^X} - \overline{T_m^Y}}{T_b}\right)^2 + \left(\frac{|\overline{\theta_m^X} - \overline{\theta_m^Y}| - \theta_{X,Y}}{\theta_c}\right)^2}, \quad (4)$$

where d is the spatial distance between two points; $\theta_{X,Y}$ is the change in nominal direction when a wave propagates from X to Y , an important correction at high latitudes; and H_a , T_b , and θ_c are empirical factors that need to be tuned. After tuning, it is found that $H_a = \min(\overline{H_s^X}, \overline{H_s^Y})$, $T_b = 1$ s, and $\theta_c = 10^\circ$ can give good results. Meanwhile, it is noted that small changes in these empirical factors have almost no impact on the result.

2) If the δ between two CWPs is less than the threshold δ_{th} , they can be regarded as the CWPs from the same WCS. The threshold is usually set as $\delta_{th} = 0.01 \text{ km}^{-1}$, which is determined from a tuning process, but this value is sometimes subjectively changed from case to case for a better tracking result.

3) When all neighboring points are searched, extend the above searching process to the second-order neighbor points (i.e., neighboring points of neighboring points) based on the wave parameters of first-order neighbors, and this process is operated recursively to higher-order neighbors (third, fourth, fifth, etc.) until no CWP can be included. After completing the tracking of the current WCS, start from step 1 again to track the next WCS until all systems are tracked.

In this procedure, the CWP is equivalent to the spectral partition in Jiang (2019) and the WCS is equivalent to the wave event in Jiang (2019). More details about this algorithm and its discussion can be found in Jiang (2019).

For each tracked WCS in the given season, its energy contribution to global wave climate, C , is computed as the ratio between the global spatial averaged wave energies of system K (E_K) and the total wave (E) as follows:

$$C = \frac{E_K}{E} = \frac{\sum [\overline{H_s^K}(\lambda, \varphi)^2 \cos \varphi]}{\sum \left[\sum_{t=1}^N H_{s,t}(\lambda, \varphi)^2 / N \right] \cos \varphi}, \quad (5)$$

where λ and φ are the longitude and latitude, respectively; N is the length of time series in the corresponding season; and $H_{s,t}(\lambda, \varphi)$ indicates the total H_s at time t . This energy contribution is similar to the variance contribution in EOF; it can be used to indicate the relative significance of each WCS.

3. Limitations of bulk wave parameters

An example is shown in Fig. 1 to illustrate the limitation of raw integrated wave parameters (H_s , T_m , θ_m , H_s^w , T_m^w , θ_m^w , H_s^s , T_m^s , θ_m^s). The left panel of Fig. 1 is an example of meridional profiles of H_s , T_m , and θ_m for the wind-sea, swells, and overall waves at 150°W in DJF. The seasonal maps for the global mean 10-m wind fields in DJF and JJA from CFSR that forces the model are shown in Fig. 2 as a reference. In Fig. 2, the color represents the mean wind speed and the directions of the arrows (not scaled with the background field) represent the seasonal mean wind directions (computed by averaging the zonal and meridional components of the wind separately). In Fig. 1, the magnitudes and directions of wind-seas are in good agreement with the wind field, and the parameters of the swells are close to those of the total waves because of the dominance of swells in the global wave field. The magnitudes and directions of the swells are consistent with those of the wind-seas to the south of $\sim 35^\circ\text{S}$ and to the north of $\sim 40^\circ\text{N}$, because the swells in this region are mainly young swells generated by the westerlies which have similar properties with the wind-seas in the same region. An abrupt change in swell/total mean θ_m , termed “swell front” (Young 1999), is observed at $\sim 20^\circ\text{S}$, which marks the dominance change of swells from the two hemispheres. However, a front should not really exist in the wave field because swell systems propagate almost independently. The result seems to be misleading to some extent. Moreover, the wind-seas also have abrupt changes in mean θ_m at $\sim 35^\circ\text{S}$ and $\sim 30^\circ\text{N}$, which corresponds to the dominance change of the westerlies and trade winds (Fig. 2a), and such “wind-sea fronts” might be also misleading sometimes.

The PDS along the same profile is plotted in the right panel of Fig. 1 at each 6° latitude. It is noted that an “oceanic convention” is used throughout the text that a direction of 0° corresponds to the wave propagating northward. At least three CWPs always coexist at any locations, and each of them corresponds to a remote or local wind climate system (Jiang and Mu 2019). The

FIG. 1. (left) The meridional profile of H_s , T_m (arrow color), and θ_m (arrow) for the wind-sea (orange line), swells (green line), and overall waves (blue line) at 150°W in DJF. The base map is used as the reference for the location of profile. It is noted that the wind-sea T_m and swell/total T_m share the same color bar but correspond to different values for the same color. (right) The PDS along the same profile at each 6° latitude; the latitudes are marked on the upper left of each subplot. A direction of 0° corresponds to the wave propagation toward the north in this paper.

signatures of swells generated by the westerlies in the North Pacific (NP) can be clearly observed in the South Pacific (SP) (i.e., the WCS at 160°–170°, 0.06–0.08 Hz in Figs. 1k–t). Similarly, the swells generated by the westerlies in the South Pacific (SP) can also be observed in the NP (i.e., the CWP at 20°–30°, 0.06–0.08 Hz in Figs. 1c–j). In the tropical region, four CWPs propagating almost perpendicularly with each other make the mean θ_m of total/wind-sea/swell loses the representativeness. Moreover, wind-sea and swell events can coexist in the same CWP. For instance, the mean θ_m of both wind-sea and swell at 57°S are eastward (the left panel), which means the eastward CWP at 57°S (Fig. 1t) consists of both wind-seas generated by the westerlies in

the SP and relatively young swells evolved from these wind-seas (swells propagating out of their sources or residual swells when wind diminishes). Because this CWP predominates in the westerlies, the mean θ_m of both wind-sea and swells are eastward at this location, and the information about the other two CWP is totally absent in the left panel of Fig. 1. This example shows that although traditional bulk wave parameters are useful for many purposes, sometimes they might not be sufficient for a detailed description of wave climates, even after the separation of wind-seas and swells.

The PDS profile in the right panel of Fig. 1 can present more information about the wave climate than the left panel. However, such an “array of wave spectra” is not a

FIG. 2. Seasonal averages of CFSR 10-m wind field for (a) DJF and (b) JJA. The arrows are not scaled with the background field.

good way of data analysis and visualization: First, the PDSs themselves do not contain the energy information for each CWP. Second, and more importantly, it is hard to visualize wave spectra on global scale with a high resolution. The right panel of Fig. 1 is only plotted every 6° of latitude, which is a rather coarse spatial resolution, and it is difficult to increase the resolution within a limited space. Such arrays of wave spectra are used to analyze global spectral wave climate with a $10^\circ \times 10^\circ$ resolution in Echevarria et al. (2019) where the propagation of swells from the westerlies of the two hemispheres is vaguely observable. The global distributions of PDS in DJF and JJA at $10^\circ \times 10^\circ$ resolution is also shown in Fig. S1 in the online supplemental material, where the propagation of some WCSs is also observable. Even with such a low spatial resolution, it is still hard to observe the features of global distributions of wave climate from these arrays. Meanwhile, global distributions of mean H_s , T_m , and θ_m for different WCSs can provide a better description of global spectral wave climate.

4. Spatial distributions of WCSs

The CWPs at all grid points are spatially tracked in different seasons to obtain global WCSs. Here, only the WCSs in DJF and JJA in the open ocean with a relatively large spatial coverage (larger than $600 1^\circ \times 1^\circ$ grid points) are shown in Figs. 3 and 4, respectively. A label is assigned to each WCS in each season, and each WCS reflects the wave climate generated by a wind climate system. To save space, several WCSs are plotted in the same map and share the same color scale if there is no overlap regarding their spatial extents. The figures with different WCSs plotted with different color scales can be found in the online supplemental material. Their energy contributions to the global (75°S – 75°N) wave climate are shown in Table 1. The WCSs shown in Figs. 3 and 4 can explain more than 90% of global wave energy.

The most interesting features in Figs. 3 and 4 are the patterns of generation and propagation of major WCSs, as the mean H_s , T_m , and θ_m of different WCSs are shown separately. The regions with highest H_s of different WCSs are in line with the region with high wind speed in Fig. 2. The spatial extent of a WCS can extend far away from their generation areas with the $H_s(T_m)$ decreasing (increasing) along the mean θ_m . The general propagation routes of these swells are also nicely shown by the mean θ_m (of the target WCS). For instance, how the waves in system JA travel from the SO to the coasts of California and Alaska along great circles on Earth's surface can be clearly observed according to the directions of arrows in Fig. 4, providing a good reference for tracking swell propagation using satellite or buoy array. It is already well

known that the swells generated by the winter storm in the Southern Hemisphere can sometimes propagate to the coast of California and Alaska (Munk et al. 1963; Snodgrass et al. 1966). The results obtained here can be regarded as the long-term averaged propagation paths of these swells, and the clear land blocking and island shadowing effects are along the direction of waves in this wave climate pattern can support this point of view. The mean T_m of any WCS generally increases with the propagation distance (along the mean θ_m), primarily because of two reasons: 1) nonlinear wave–wave interaction pumps the wave energy into lower frequencies during the evolution of waves, and 2) high-frequency swells dissipate faster than low-frequency ones (Ardhuin et al. 2009).

Some problems are also found in the results. Theoretically, the mean H_s in the boundaries of all WCSs should be close to zero, and the boundary outlines should be smooth in the open ocean. However, this is not the case for most WCSs. These problems are mainly due to the limitation of watershed-algorithm-based spectral partitioning method used twice in this study. The watershed-algorithm-based partitioning scheme (e.g., Hanson and Phillips 2001; Portilla et al. 2009) is a purely morphological method that does not consider wave dynamics. During spectrum partitioning, when the energy peak of a partition becomes smaller than the tail of a nearby partition, it might be overwhelmed by the nearby partition, and the watershed-algorithm scheme cannot identify it (Ailliot et al. 2013). In addition, it is noted that the WAVEWATCH-III retains at most six partitions at a given location, but sometimes more than six partitions might coexist in modeled wave spectra (although this case does not occur frequently) so that the partition with very low H_s might be also omitted in IOWAGA. Both of the two factors lead to the result that the partitioning with very low H_s might be missed in the dataset, and this affects the computation of PDS. Meanwhile, such a problem also exists during the “partitioning” of PDSs, leading to some discontinuous edges of WCSs. This problem of the spectral partitioning algorithm seems to be difficult to solve at this stage. In spite of this problem, this study presents a reasonable global view of wave climate separating the contributions of different WCSs.

a. Prevailing westerlies-generated WCSs

Systems DA, DB, and DD in Fig. 3 and systems JA, JD, and JL in Fig. 4 are the waves generated by the prevailing westerlies. They are the WCSs with the largest energy contributions to the global wave climate. In DJF, the maximum mean H_s ($\sim 5\text{ m}$) are found in the NP (system DA) and North Atlantic (NA; system DB) owing to the strong winter storm with averaged wind speed higher than 12 m s^{-1} (Fig. 2a). The waves

FIG. 3. Seasonal mean (left) H_s , (right) T_m , and θ_m (arrows) of the main tracked WCSs (labeled in purple letters) for DJF. The arrows are not scaled with the background fields. The systems without any spatial overlap are plotted on the same map, and the color scales are different for different subplots.

generated in these regions can gradually turn into swells and propagate to the coast of the Antarctic continent in the Southern Hemisphere (SH). In JJA, these two WCSs have less contribution to the global wave climate. With a maximum mean H_s of ~ 2 m, only a small amount of energy of systems JL and JD can propagate over the

equator due to the strong seasonality of the westerlies in the Northern Hemisphere (NH).

The WCS with the largest spatial extent in both DJF and JJA is generated by storm tracks in SH, because the Southern Ocean (SO) sectors of Pacific, Atlantic, and Indian Oceans are connected to each other. The highest

FIG. 4. As in Fig. 3, but for JJA.

mean H_s in systems JA and DD are both found in the SO Indian sector in all seasons. In JJA, the WCS generated by the westerlies in the SH (system JA) has a maximum mean H_s of more than 5 m and can travel across all the three oceans, and this WCS is overwhelming predominant in global wave climate with an energy contribution of almost 70%. Because the seasonality of the westerlies in the

SH is weaker than that in the NH, the WCS generated by the austral summer westerlies (system DD) also has a high mean H_s (~ 4 m) in the generation areas and can affect the wave climate in the entire Indian Ocean (IO) as well as a large proportion of the Pacific and Atlantic Oceans.

It is somewhat difficult to find the results of global distributions of spectral wave climate parameters in

TABLE 1. Energy contribution of WCSs.

DJF	System label	DA	DB	DC	DD	DE	DF	DG	DH	DI
	Contribution (%)	11	17	2	38	0.4	2	3	2	1
	System label	DJ	DK	DL	DM	DN	DO	DP	DQ	
	Contribution (%)	0.1	2	1	2	0.6	0.4	5	2	
JJA	System label	JA	JB	JC	JD	JE	JF	JG	JH	JI
	Contribution (%)	69	1	3	2	1	3	4	2	0.3
	System label	JJ	JK	JL	JM	JN				
	Contribution (%)	3	1	2	1	2				

previous studies that can be directly compared to the results obtained in this study. However, the results can be indirectly validated by the wave climate patterns derived from bulk wave parameters in previous studies (e.g., Young 1999; Alves 2006; Semedo et al. 2011; Jiang et al. 2017) and the wind climate patterns in Fig. 2. The patterns of these WCSs at their origins (westerlies) are in line with previous results obtained using bulk wave parameters (e.g., Semedo et al. 2011; Fan et al. 2014). However, because the mean H_s values of these WCSs decline rapidly out of the generation zone, previous studies on wave climate based on wave model hindcast or satellite data did not quantify the impacts of westerlies-generated WCSs on global wave climate in far fields, and even their existence is overwhelmed by the stronger waves generated in the other hemispheres. The results here show clearly that westerlies-generated swells can propagate to the equator in all seasons and can extend to the other end of the basin in hemispheric winters, although they may become less dominant after propagating over large distances. Besides, the mean T_m of the westerlies-generated WCSs can vary from less than 11 s in source regions to more than 15 s in far fields after propagating over more than 10 000 km due to the two reasons mentioned above. A problem of the result herein is that the mean H_s of the westerlies-generated WCSs is observed to increase in the far fields after propagating over 10 000 km, which is unreasonable. This is because the rotating extratropical storms can generate some strong local waves with similar T_m and θ_m to the waves originating from the other hemisphere, and these local and remote waves are identified as the same WCSs by the partitioning scheme, resulting in the “increase” of mean H_s in far fields. Because of these problems, only one significant digit is maintained in Table 1.

By keeping only the wind in selected oceanic areas active, Alves (2006) eliminates the impact of the waves outside the generation zone. The spatial patterns of mean H_s of systems DA and DB are similar to those of the annual mean H_s of waves originating from the extratropical NP and NA demonstrated in Alves (2006), respectively. According to the distributions of SO westerlies in Fig. 2, it should be more reasonable to consider the

SO westerlies-generated WCS as a single system, rather than as three systems that are separated subjectively in Alves (2006). However, the major propagating patterns of these waves are also similar in the two studies, showing the validity of the method and results in this study. The propagation patterns of these WCSs during their less dominant stages are also identified by Echevarria et al. (2019) in the first EOF mode of global wave spectra. But the EOF pattern of Echevarria et al. (2019) shows that the NH-generated swell can propagate into the SH farther than the more energetic SH swell, which propagates northward, which is in line neither with the results of the present research nor with Alves (2006). Checking the long-term PDS (available in Fig. S1), the signatures of the SH westerlies-generated WCSs can be observed to the north of 50°N in both DJF and JJA, whereas those of the NH westerlies-generated WCSs can be observed at 50°S only in DJF, also indicating that the SH-generated swell can propagate farther. It is worth noting that the first EOF mode in Echevarria et al. (2019) mainly represents the seasonality, instead of the strength of wave energy. The NH westerlies-generated swells are observed to propagate farther in Echevarria et al. (2019) simply because they have a stronger seasonality.

b. Trade winds-generated WCSs

Systems DC, DG, DI, DP, and DQ in Fig. 3 are the WCSs along the trade winds in DJF, and systems JF, JG, JH, JC, and JK in Fig. 4 are the corresponding WCSs in JJA. Although the trade winds are stronger in the winter hemisphere, all these five WCSs are clearly observed in all three oceans for all seasons. Because of the lower wind speed compared to the westerlies, both the mean H_s and the mean T_m of the WCSs generated by trade winds are lower than those generated by the westerlies. These trade winds-generated WCSs have the southwestward and northwestward components of mean θ_m in their southern and northern parts, respectively. The signatures of the regional winds effect along the five eastern boundary currents systems (the California and Canary Currents in JJA, and Humboldt, Benguela, and West Australia Currents in DJF; Carrasco et al. 2014; Semedo et al. 2018) can be observed as a part of these

trade winds–generated WCSs (the signals are clearer in the supplemental materials where different WCSs are shown with different color bars), which can be associated with the wind field illustrated in Fig. 2. For instance, the signature of the CLCJ is clearly shown as part of system JC generated by the trade winds in the NP. The range of influence of the waves generated by the CLCJ can reach the eastern tropical Pacific, which is consistent with the results of Jiang and Mu (2019), and can also be confirmed by the PDS array in the eastern tropical Pacific in Fig. S1b in the supplemental material. In addition, the WCS of trade winds in the SP in both DJF and JJA (i.e., systems DG and JG) has two centers of high mean H_s , which is in agreement with the structure of the trade wind zone in the SP. The regions in the “upstream” of the two WCSs have a lower T_m mean, showing that some of the westward waves originating from the trade wind zones to the west of the South America may receive energy again from the local wind in the western side of the basin.

Even in the center of the trade wind zones, the mean H_s of the trade winds–generated WCSs are generally lower than those of the westerlies-generated swells that have not propagated far away from their origins. Therefore, the impact of trade winds on the seasonal averages of total H_s fields is not well defined in either DJF or JJA [e.g., Figs. 2a and 3a of Semedo et al. (2011)] except for the one in the IO in JJA, which has a maximum mean H_s of more than 2.5 m (system JF). After separating the wind-sea and swell climates, these WCSs become observable along the trade wind zones in global distributions of the wind-sea H_s fields [e.g., Figs. 2c and 3c Semedo et al. (2011)]. However, it is difficult to display the trade wind–generated swell systems with lower H_s outside their origins by using bulk wave parameters. In Fig. S1 in the supplemental material, the signatures of the trade winds–generated swells are also well defined in both seasons (e.g., the northwestward and southwestward peaks along the equator). The signals of trade winds are also observed in Echevarria et al. (2019), but the spatial pattern is not that clear. Some of the patterns derived in this study can also be partially validated by the patterns derived by Alves (2006) where the tropical sections of the NA, South Atlantic (SA), eastern NP, eastern SP, and south IO are forced separately. However, the spatial extent of the patterns in Alves (2006) seems to be larger than in the present study. For example, in Alves (2006), the waves generated in the tropical eastern NP can propagate to Antarctic coasts and the waves generated in the tropical South IO can propagate to east coasts of South America. There are at least three reasons which can explain this difference: 1) The swell dissipation term is included in the IOWAGA hindcast whereas it is not taken into account in Alves (2006). Thus, the swell can

propagate farther in the model of Alves (2006). 2) Due to the seasonal shifting of the wind belts, the westerlies can sometimes reach south (north) of 30°N (30°S). Therefore, in Alves (2006), some signals of westerlies-generated waves are also displayed in the patterns of waves generated in the tropical oceans. This phenomenon is clearly shown in Fig. 13 of Alves (2006). 3) Even if some swells can propagate farther than the spatial extent shown in Figs. 3 and 4, the occurrence frequency of these swells decrease with the increase of the distance from the origins. The signals of these swells can sometimes be too low to be isolated in the long-term PDS, so that they are not identified in far fields.

c. Indian monsoon–generated WCSs

Systems DE and DJ in DJF and systems JB and JI in JJA are the WCSs generated by the Indian monsoon. Although their spatial extents and energy contributions to global wave climate are relatively small, the Indian monsoon has high wind speeds, especially in JJA, and their signals in the wave fields are well marked (e.g., Semedo et al. 2011; Ranjha et al. 2015). As the Indian monsoon is stronger in JJA, both systems JF and JI have much lower mean H_s and shorter T_m than systems DE and DJ. Within limited fetches, both the mean H_s and T_m of system JB are higher than those of system JI because the Somali coastal jet has higher wind speeds (Fig. 2). The directions of the Indian monsoon are mainly northeastward in JJA; thus, the ranges of influence the two WCSs are mainly the Arabian Sea and Bay of Bengal, respectively. However, it is observed that some of the waves in systems JB and JI can propagate eastward across the IO and reach the northwestern shore of Australia. This phenomenon can be also observed in Alves (2006) where the mean H_s field is forced by the wind in the northern IO.

d. Polar easterlies–generated WCSs

All the aforementioned WCSs correspond to regions with a relatively high local wind speed, which can be clearly observed in Fig. 2 (i.e., the westerlies, trade winds, and monsoon). However, it seems that the source regions of some WCSs are not very clearly characterized in the global distribution of wind speed. According to their spatial patterns, some of these WCSs should be generated by the polar easterlies. In DJF, systems DF, DH, and DK seem to be generated by the easterlies in the Bering Sea, the Norwegian Sea, and the coastal region of Antarctic, respectively, while systems JM and JN are polar easterlies–induced WCSs in JJA. In fact, the corresponding polar wind systems of these WCSs can be observed in the seasonal mean wind in Fig. 2, and the mean wind directions of these wind systems are also consistent with the mean θ_m of WCSs. However, because the polar easterlies–generated WCSs have a lower

wave energy than the westerlies-generated WCSs and their spatial impacting ranges have a large overlap, it seems that the polar easterlies-generated WCSs have often been overlooked in previous studies. Because Alves (2006) does not separate the polar regions and the extratropical regions, these patterns are not observed, either. Another WCS generated by the polar wind system is system DM, which corresponds to the south-eastward wind in the Labrador Sea (which is clear in Fig. 2a). This WCS can be partly observed in the distributions of mean H_s and wind-sea H_s at its origin, the Labrador Sea [Figs. 2a and 2c in Semedo et al. (2011)].

No polar easterlies-generated WCS is found in the NP in JJA, because the polar easterlies between 150°E and 120°W are mainly located in the Arctic Ocean in this period, as shown in Fig. 2b, and the Asian and North American continents block the polar easterlies-generated waves in the Arctic Ocean from propagating into the NP. System JN is also observed not to extend to the Pacific Ocean. A potential reason for this phenomenon is that the sea surface polar easterlies in the SO Pacific sector are mainly located in the Ross Sea to the south of 65°S (Fig. 2b), which is frozen in JJA. Therefore, these polar easterlies do not affect the waves in the SP in JJA.

e. Other WCSs

It seems to be relatively difficult to explain the rest of the tracked WCSs, including systems DL, DN, DO, JJ, and JE, because they do not correspond to any wind feature in Fig. 2. A potential reason for the formation of these WCSs is that they are the “remainder” WCSs generated by extratropical storms in the westerlies. Most of the waves generated in the westerlies are eastward, but the rotating winter storms can also generate westward waves, which might form some WCSs. Alves (2006) also shows that the wind in the extratropical regions can generate swell systems that propagate westward against the predominant wave directions. However, the westward wave patterns in Alves (2006) should be the superposition of these remainder WCSs and the polar easterlies-generated WCSs. In regions where systems DN and DO are located, the mean T_m and θ_m of these WCSs are different from those of the polar easterlies-generated WCSs so that they can be tracked by the long-term PDS. However, in some other regions (e.g., the region near 90°E in the SO), because the T_m and θ_m of so-called remainder WCSs are similar to those of the polar easterlies-generated WCSs, it is difficult to distinguish them from the long-term PDS.

5. Summary

Several WCSs often coexist at the same location, leading to the complex wave climate in the ocean. Using

integrated wave parameters (even if after wind-sea-swell separation) to characterize the global wave climate can only capture the feature of one or two predominant WCSs and often overlooks the WCSs with relatively low energy. In this study, an attempt is made to present a relatively comprehensive global view of wave climate by spatially tracking spectral WCSs in point-wise long-term PDSs. Tens of spatially consistent WCSs essentially independent of each other are tracked and each of them corresponds to a global or regional map of mean H_s , T_m , and θ_m of waves from the same origins, exhibiting more details of global wave climate. These WCSs reveal the process of how wind-seas gradually turn into swells, as well as the propagation route of global waves. Every region with relatively high wind speeds including the prevailing westerlies, the trade winds, polar easterlies, and Indian monsoon corresponds to an individual WCS. Among these WCSs, those generated by the westerlies, which can travel across the entire basin, play the most important role in the global wave climate. The patterns of these WCSs demonstrate the entire life cycle of ocean waves, from being dominant to become less dominant, from a climatic point of view. Such climatic patterns of global wave propagation obtained in this study can serve as a reference for designing satellite tracks and arranging buoy locations when one wants to track swell propagation using remote sensors (e.g., Jiang et al. 2016b) or buoy arrays (e.g., Snodgrass et al. 1966). Knowing the impacting range of different WCSs may help determine the modeling domain when running a regional wave model. Besides, WCSs from different sources may also have different properties when approaching the coasts, leading to different coastal processes and water-level responses. Therefore, some results of this study may also be useful for coastal research. This study mainly focuses on the seasonal variability of global wave climate while future exploration can be made to regional wave climate using data with higher spatial resolutions and to the interannual/long-term variabilities of each WCS using consistent data with longer time spans. Moreover, with the successful launching of CFOSAT (the China–France Oceanography Satellite) on 29 October 2018, observational global ocean wave spectra will be available from the Surface Waves Investigation and Monitoring (SWIM) sensor. With the accumulation of these remotely sensed wave spectra, the methodology employed in this study can also be used to present observational wave climate patterns in future works.

Acknowledgments. This work is jointly supported by the National Key Research and Development Program of China (2018YFC0309601), the National Natural Science Foundation of China (41806010), Laboratory for Regional Oceanography and Numerical Modeling,

Qingdao National Laboratory for Marine Science and Technology (2019A03), and the Guangdong Special Fund Program for Marine Economy Development (GDME-2018E001). The IOWAGA data are downloaded from IFREMER ftp (ftp.ifremer.fr). The author would like to thank Dr. Alexander Babanin, Dr. Alvaro Semedo, Dr. Qingxiang Liu, and three anonymous reviewers for their helpful comments and suggestions.

REFERENCES

- Aarnes, O. J., S. Abdalla, J. R. Bidlot, and Ø. Breivik, 2015: Marine wind and wave height trends at different ERA-Interim forecast ranges. *J. Climate*, **28**, 819–837, <https://doi.org/10.1175/JCLI-D-14-00470.1>.
- Ailliot, P., C. Maisondieu, and V. Monbet, 2013: Dynamical partitioning of directional ocean wave spectra. *Probab. Eng. Mech.*, **33**, 95–102, <https://doi.org/10.1016/j.pro bengmech.2013.03.002>.
- Alves, J. H. G. M., 2006: Numerical modeling of ocean swell contributions to the global wind-wave climate. *Ocean Modell.*, **11**, 98–122, <https://doi.org/10.1016/j.ocemod.2004.11.007>.
- Ardhuin, F., B. Chapron, and F. Collard, 2009: Observation of swell dissipation across oceans. *Geophys. Res. Lett.*, **36**, L06607, <https://doi.org/10.1029/2008GL037030>.
- , and Coauthors, 2010: Semi-empirical dissipation source functions for wind-wave models: Part I: Definition, calibration and validation. *J. Phys. Oceanogr.*, **40**, 1917–1941, <https://doi.org/10.1175/2010JPO4324.1>.
- Carrasco, A., A. Semedo, P. E. Isachsen, K. H. Christensen, and Ø. Saetra, 2014: Global surface wave drift climate from ERA-40: The contributions from wind-sea and swell. *Ocean Dyn.*, **64**, 1815–1829, <https://doi.org/10.1007/s10236-014-0783-9>.
- Chen, G., B. Chapron, R. Ezraty, and D. Vandemark, 2002: A global view of swell and wind sea climate in the ocean by satellite altimeter and scatterometer. *J. Atmos. Oceanic Technol.*, **19**, 1849–1859, [https://doi.org/10.1175/1520-0426\(2002\)019<1849:AGVOSA>2.0.CO;2](https://doi.org/10.1175/1520-0426(2002)019<1849:AGVOSA>2.0.CO;2).
- Cox, A. T., and V. R. Swail, 2001: A global wave hindcast over the period 1958–1997: Validation and climate assessment. *J. Geophys. Res.*, **106**, 2313–2329, <https://doi.org/10.1029/2001JC000301>.
- Delpey, M. T., F. Ardhuin, F. Collard, and B. Chapron, 2010: Space-time structure of long ocean swell fields. *J. Geophys. Res.*, **115**, C12037, <https://doi.org/10.1029/2009JC005885>.
- Echevarria, E. R., M. A. Hemer, and N. J. Holbrook, 2019: Seasonal variability of the global spectral wind-wave climate. *J. Geophys. Res. Oceans*, **124**, 2924–2939, <https://doi.org/10.1029/2018JC014620>.
- Fan, Y., S. Lin, I. M. Held, Z. Yu, and H. L. Tolman, 2012: Global ocean surface wave simulation using a coupled atmosphere–wave model. *J. Climate*, **25**, 6233–6252, <https://doi.org/10.1175/JCLI-D-11-00621.1>.
- , I. M. Held, S. Lin, and X. Wang, 2013: Ocean warming effect on surface gravity wave climate change for the end of the twenty-first century. *J. Climate*, **26**, 6046–6066, <https://doi.org/10.1175/JCLI-D-12-00410.1>.
- , S. Lin, S. M. Griffies, and M. A. Hemer, 2014: Simulated global swell and wind-sea climate and their responses to anthropogenic climate change at the end of the twenty-first century. *J. Climate*, **27**, 3516–3536, <https://doi.org/10.1175/JCLI-D-13-00198.1>.
- Gulev, S. K., and V. Grigorieva, 2004: Last century changes in ocean wind wave height from global visual wave data. *Geophys. Res. Lett.*, **31**, L24302, <https://doi.org/10.1029/2004GL021040>.
- , —, A. Sterl, and D. Woolf, 2003: Assessment of the reliability of wave observations from voluntary observing ships: Insights from the validation of a global wind wave climatology based on voluntary observing ship data. *J. Geophys. Res.*, **108**, 3236, <https://doi.org/10.1029/2002JC001437>.
- Hanson, J. L., and O. M. Phillips, 2001: Automated analysis of ocean surface directional wave spectra. *J. Atmos. Oceanic Technol.*, **18**, 277–293, [https://doi.org/10.1175/1520-0426\(2001\)018<0277:AAOOSD>2.0.CO;2](https://doi.org/10.1175/1520-0426(2001)018<0277:AAOOSD>2.0.CO;2).
- Hemer, M. A., J. A. Church, and J. R. Hunter, 2010: Variability and trends in the directional wave climate of the Southern Hemisphere. *Int. J. Climatol.*, **30**, 475–491, <https://doi.org/10.1002/joc.1900>.
- , Y. Fan, N. Mori, A. Semedo, and X. Wang, 2013: Projected changes in wave climate from a multi-model ensemble. *Nat. Climate Change*, **3**, 471–476, <https://doi.org/10.1038/nclimate1791>.
- Jiang, H., 2019: Spatially tracking wave events in partitioned numerical wave model outputs. *J. Atmos. Oceanic Technol.*, **36**, 1933–1944, <https://doi.org/10.1175/JTECH-D-18-0228.1>.
- , and L. Mu, 2019: Wave climate from spectra and its connections with local and remote wind climate. *J. Phys. Oceanogr.*, **49**, 543–559, <https://doi.org/10.1175/JPO-D-18-0149.1>.
- , A. V. Babanin, and G. Chen, 2016a: Event-based validation of swell arrival time. *J. Phys. Oceanogr.*, **46**, 3563–3569, <https://doi.org/10.1175/JPO-D-16-0208.1>.
- , J. E. Stopa, H. Wang, R. Husson, A. Mouche, B. Chapron, and G. Chen, 2016b: Tracking the attenuation and nonbreaking dissipation of swells using altimeters. *J. Geophys. Res. Oceans*, **121**, 1446–1458, <https://doi.org/10.1002/2015JC011536>.
- , A. Mouche, H. Wang, A. Babanin, B. Chapron, and G. Chen, 2017: Limitation of SAR quasi-linear inversion data on swell climate: An example of global crossing swells. *Remote Sens.*, **9**, 107, <https://doi.org/10.3390/rs9020107>.
- Munk, W. H., G. R. Miller, F. E. Snodgrass, and N. F. Barber, 1963: Directional recording of swell from distant storms. *Philos. Trans. Roy. Soc. London*, **A255**, 505–584, <https://doi.org/10.1098/rsta.1963.0011>.
- Portilla, J., 2018: The global signature of ocean wave spectra. *Geophys. Res. Lett.*, **45**, 267–276, <https://doi.org/10.1002/2017GL076431>.
- , F. J. Ocampo-Torres, and J. Monbaliu, 2009: Spectral partitioning and identification of wind sea and swell. *J. Atmos. Oceanic Technol.*, **26**, 107–122, <https://doi.org/10.1175/2008JTECHO609.1>.
- , L. Cavaleri, and G. P. V. Vledder, 2015: Wave spectra partitioning and long term statistical distribution. *Ocean Modell.*, **96**, 148–160, <https://doi.org/10.1016/j.ocemod.2015.06.008>.
- Ranjha, R., M. Tjernström, A. Semedo, G. Svensson, and R. M. Cardoso, 2015: Structure and variability of the Oman coastal low-level jet. *Tellus*, **67A**, 25285, <https://doi.org/10.3402/tellusa.v67.25285>.
- Rascle, N., and F. Ardhuin, 2013: A global wave parameter database for geophysical applications. Part 2: Model validation with improved source term parameterization. *Ocean Modell.*, **70**, 174–188, <https://doi.org/10.1016/j.ocemod.2012.12.001>.
- Semedo, A., K. Sušelj, A. Rutgersson, and A. Sterl, 2011: A global view on the wind sea and swell climate and variability from ERA-40. *J. Climate*, **24**, 1461–1479, <https://doi.org/10.1175/2010JCLI3718.1>.
- , R. Weiss, A. Behrens, A. Sterl, L. Bengtsson, and H. Günther, 2013: Projection of global wave climate change towards the end of the 21st century. *J. Climate*, **26**, 8269–8288, <https://doi.org/10.1175/JCLI-D-12-00658.1>.

- , P. M. M. Soares, and G. Lemos, 2018: Waves along eastern boundary currents—The regional winds effect. *Ocean Modell.*, **129**, 39–57, <https://doi.org/10.1016/j.ocemod.2018.07.005>.
- Snodgrass, F. E., G. W. Groves, K. Hasselmann, G. R. Miller, W. H. Munk, and W. H. Powers, 1966: Propagation of ocean swell across the Pacific. *Philos. Trans. Roy. Soc. London*, **A249**, 431–497, <https://doi.org/10.1098/RSTA.1966.0022>.
- Stopa, J. E., F. Ardhuin, A. V. Babanin, and S. Zieger, 2016: Comparison and validation of physical wave parameterizations in spectral wave models. *Ocean Modell.*, **103**, 2–17, <https://doi.org/10.1016/j.ocemod.2015.09.003>.
- WAVEWATCH III Development Group, 2016: User manual and system documentation of WAVEWATCH III version 5.16. Tech. Note 329, NOAA/NWS/NCEP/MMAB, 326 pp., <http://polar.ncep.noaa.gov/waves/wavewatch/manual.v5.16.pdf>.
- Young, I. R., 1999: Seasonal variability of the global ocean wind and wave climate. *Int. J. Climatol.*, **19**, 931–950, [https://doi.org/10.1002/\(SICI\)1097-0088\(199907\)19:9<931::AID-JOC412>3.0.CO;2-O](https://doi.org/10.1002/(SICI)1097-0088(199907)19:9<931::AID-JOC412>3.0.CO;2-O).
- , S. Zieger, and A. V. Babanin, 2011: Global trends in wind speed and wave height. *Science*, **332**, 451–455, <https://doi.org/10.1126/science.1197219>.
- , J. Vinoth, S. Zieger, and A. V. Babanin, 2012: Investigation of trends in extreme value wave height and wind speed. *J. Geophys. Res.*, **117**, C00J06, <https://doi.org/10.1029/2011JC007753>.