Evaluating the Eastward Propagation of the MJO in CMIP5 and CMIP6 Models Based on a Variety of Diagnostics

YUE LI,^a JIYE WU,^a JING-JIA LUO,^a AND YOUNG MIN YANG^a

^a *Institute of Climate Application Research (ICAR)/CICFEM/KLME/ILCEC, Nanjing University of Information Science and Technology, Nanjing, China*

(Manuscript received 18 May 2021, in final form 24 November 2021)

ABSTRACT: Given the climatic importance of the Madden–Julian oscillation (MJO), this study evaluates the capability of CMIP6 models in simulating MJO eastward propagation in comparison with their CMIP5 counterparts. To understand the representation of MJO simulation in models, a set of diagnostics with respect to MJO-associated dynamic and thermodynamic structures is applied, including large-scale zonal circulation, vertical structures of diabatic heating and equivalent potential temperature, moisture convergence at the planetary boundary layer (PBL), and the east–west asymmetry of moisture tendency relative to the MJO convection. The simulated propagation of the MJO in CMIP6 models shows an overall improvement in realistic speed and longer distance, which displays a robust linear regression relationship against the above-mentioned dynamic and thermodynamic structures. The improved MJO propagation in CMIP6 largely benefits from better representation of premoistening processes that is primarily contributed by improved PBL moisture convergence. In addition, the convergence of moisture and meridional advection of moisture prior to the MJO convection are enhanced in CMIP6, while the zonal advection of moisture is as weak as that in CMIP5. The increased convergence of moisture is a result of enhanced lower-tropospheric moisture and divergence, and the enhanced meridional advection of moisture can be caused by sharpened meridional gradient of mean lower-tropospheric moisture in the western Pacific. Further examination of the lower-tropospheric moisture budget reveals that the enhanced zonal asymmetry of the moisture tendency in CMIP6 is driven by the drying process to the west of the MJO convection, which is attributed to the negative vertical and zonal advective of moisture.

KEYWORDS: Tropics; Madden–Julian Oscillation; Model comparison; Model evaluation/performance; Intraseasonal variability

1. Introduction

The Madden–Julian oscillation (MJO), first documented by [Madden and Julian \(1971, 1972\)](#), is a dominant mode of tropical intraseasonal fluctuation (~30–90 days) with an eastward-moving (~5 m s^{−1} on average speed) large-scale envelope of convective cloud clusters. The MJO not only has a profound impact on tropical weather and climate phenomena, but also influences a wide region outside the tropics via atmospheric teleconnections (e.g., [Alvarez et al. 2016](#); [Arcodia et al. 2020](#); [Garfinkel et al. 2014](#); [Kim et al. 2020](#); [Klotzbach 2010, 2014](#); [Wheeler et al. 2009](#); [S. Zhou et al. 2012](#); [Zhou et al. 2020](#)). It has also been considered one of important sources of predictability at the intraseasonal-to-seasonal time scale (e.g., [Vitart et al. 2017](#); [Waliser 2011](#)). However, state-of-art general circulation models (GCMs) still show insufficient capability in simulating and predicting MJO, in part due to lack of a poor representation of the core processes that generate the atmospheric instability and slow eastward propagation (e.g., [Ahn et al. 2017](#); [Hung et al. 2013](#); [Jiang et al. 2015](#); [Lin et al. 2006](#); [Ling et al. 2017, 2019](#); [Fu et al. 2013](#)).

Numerous previous observational studies have revealed that the typical MJO structure is a deep active convection with enhanced column moisture and strong upward motion, accompanied by prevailing overturning circulation near the convective center (e.g., [Jiang et al. 2020a](#); [Kiladis et al. 2005](#); [Zhang 2005](#)). The observed MJO envelope is characterized by an east–west asymmetry relative to the deep convective center in dynamic and thermodynamic fields. To the east of the MJO convective center, warm SST ([Hendon and Glick 1997](#)) and enhanced lower-tropospheric moisture anomalies ([Tian et al. 2010](#)) provide a favorable condition for the building up of shallow cumuli and congestus that transition to deep convection ([Chen and Del Genio 2009](#)). To the west of the MJO convective center, the upper-tropospheric stratiform precipitation leads to dry and cold air in the lower troposphere ([Kiladis et al. 2005](#)). Such an asymmetry involves a series of multiscale interactions and feedbacks among moisture, convection, cloud radiation, and circulation.

A variety of theories have been proposed to explain the atmospheric instability preceding the center of active convection and how it supports the growth of MJO structure and powers the eastward propagation ([Jiang et al. 2020a](#); [Zhang et al. 2020](#)), ranging from some earlier theories, such as the wave-conditional instability of the second kind (CISK; [Lau and Peng 1987](#)), the wind-induced surface heat exchange mechanism (WISHE; [Emanuel 1987](#); [Neelin et al. 1987](#)), and the modified wave-CISK with frictional PBL convergence feedback (frictional CISK; [Wang 1988](#); [Wang and Rui 1990](#)),

Supplemental information related to this paper is available at the Journals Online website: <https://doi.org/10.1175/JCLI-D-21-0378.s1>.

Corresponding author: Jing-Jia Luo, jingjia_luo@hotmail.com and jjluo@nuist.edu.cn

DOI: 10.1175/JCLI-D-21-0378.1

© 2022 American Meteorological Society. For information regarding reuse of this content and general copyright information, consult the [AMS Copyright Policy \(www.ametsoc.org/PUBSReuseLicenses\)](#).

to some modern theories, such as the skeleton theory that explains the two-way interaction of planetary-scale envelope of MJO and sub-planetary-scale wave activity (Majda and Stechmann 2009), the gravity wave theory (Yang and Ingersoll 2011, 2013), the moisture mode theory under weak temperature gradient balance (WTG-moisture mode; Adames and Kim 2016; Sobel and Maloney 2012, 2013) or incorporating WISHE as a propagation mechanism (WISHE-moisture mode; Fuchs and Raymond 2005, 2007, 2017), and the trio-interaction theory (Wang et al. 2016; Wang and Chen 2017). Most modern theories emphasize the important role of the moist thermodynamics in the MJO.

In the last decades, the WTG-based moisture mode theory has been widely used to diagnose the MJO in observations and models (e.g., Sobel and Maloney 2013; Adames and Kim 2016; Hsu et al. 2014; Hu et al. 2020); this considers the inclusion of prognostic moisture to be important to the MJO-related atmospheric disturbance. In this moisture-mode framework, the wind associated with the MJO instantaneous response to the equatorial heat source to form a Matsuno–Gill structure and the intraseasonal heating anomalies are in WTG balance (Jiang et al. 2020a). The eastward propagation of the MJO is maintained by large-scale horizontal and vertical moisture advection, which contributes to a zonal asymmetry of the column-integrated moist static energy (MSE) tendency with positive (negative) anomalies to the east (west) of the MJO convection. This theory also emphasizes the role of longwave cloud–radiation feedback in the free atmosphere in generating and sustaining intraseasonal precipitation anomalies.

Different from the moisture mode under WTG balance, which mainly focuses on the moisture evolution, the trio-interaction theory depicts a three-way interaction among convective heating, moisture, and coupled Kelvin–Rossby wave related large-scale dynamics (Wang et al. 2016). The trio-interaction theory is a complex construct that is integrated from several major existing theoretical models, including the frictionally coupled Kelvin–Rossby wave, the moisture mode, the frictionally coupled dynamic moisture mode, the MJO skeleton, and the gravity wave theory, emphasizing the role of moisture convergence occurring at the planetary boundary layer (PBL). The structure of MJO active convection resembles a Matsuno–Gill response to a moving near-equatorial heat source (Matsuno 1966; Gill 1980), which displays a Kelvin wave response to the east of the heating and a Rossby wave response to the west. The PBL moisture convergence plays an important role in coupling the equatorial Rossby and Kelvin waves with convective heating and generating the zonal planetary-scale instability that thereby leads to the eastward propagation (Wang et al. 2016). Hu et al. (2020) found that the zonal asymmetry of moisture in the PBL can be contributed by the PBL moisture convergence, promoting the propagation at the absence of positive MSE tendency to the east of the convection.

Although the MJO simulation has largely been improved due to the progresses in theoretical understanding of MJO dynamics, the ability of GCMs in reproducing MJO propagation is still far from satisfactory (e.g., Hung et al. 2013; Lin

et al. 2006). The poor capabilities of MJO simulation in GCMs can be due to the biases in mean state. For example, Ray et al. (2011) found that the deficiencies in MJO simulations can be attributed to errors in the atmospheric mean states. Gonzalez and Jiang (2017) found that the spatial pattern of the mean lower-tropospheric moisture over the Indo-Pacific is highly correlated with the MJO propagating skill in model. Through a multimodel intercomparison, Jiang (2017) showed that the deficiencies in simulating the MJO circulation and its related seasonal mean moisture entropy are responsible for the poor representations of MJO eastward propagation across the tropical Indian Ocean in some GCMs. In addition, systematic biases in simulated mean moisture can lead to the poor MJO simulation in GCMs (Jiang et al. 2020b).

Additionally, some insufficient parameterized processes that are associated with MJO dynamics can lead to poor simulated MJO (e.g., Lin et al. 2006, 2007, 2008; Thayer-Calder and Randall 2009; Li et al. 2009; Deng and Wu 2010; Subramanian et al. 2011; Kim and Kang 2012; L. Zhou et al. 2012). It is suggested that appropriate changes in parameterizations, especially of cumulus convection, can improve the MJO simulation (e.g., Lin et al. 2006, 2008; Chikira and Sugiyama 2010). For example, Zhang and Song (2009) emphasized the importance of correct representation of the interactions between shallow and deep convections in GCMs. They found that the premoistening process at the lower troposphere triggered by shallow convection can facilitate the development of deep convection, thereby leading to a realistic MJO simulation. Seo and Wang (2010) showed the important role of the stratiform heating in properly simulating the MJO via influencing the vertical heating structure and moisture–stratiform instability process. The convective–stratiform cloud processes can also interact with moisture and large-scale circulation to favor the low-level moisture convergence to the east of MJO convection, which has been demonstrated to be important to MJO propagation (Seo and Wang 2010).

To better evaluate GCMs' representation of the MJO, a suite of diagnostics has been developed by the U.S. Climate Variability and Predictability (CLIVAR) MJO Working Group (Waliser et al. 2009). In addition, the process-oriented (Kim et al. 2014) and dynamics-oriented (Wang et al. 2018) diagnostics are often used as supplementary analyses for intermodel comparison of the simulated MJO in GCMs. The process-oriented diagnostics aim to identify processes in model that are crucial to the MJO simulation, while the dynamics-oriented diagnostics, derived from the trio-interaction theory, examine whether a model can represent the three-dimensional dynamic and thermodynamic structures associated with the MJO. These diagnostics have been widely used in model intercomparison studies (e.g., Ahn et al. 2017, 2020; Gonzalez and Jiang 2017; Jiang 2017; Jiang et al. 2015; Kim et al. 2009, 2014; Wang et al. 2018). Jiang et al. (2015) showed that about 25% of participating models can simulate a reasonable MJO, which is a great improvement regarding to MJO simulation fidelity in multimodel group. The Coupled Model Intercomparison Project (CMIP), as one of the important multimodel groups, has been evaluated in terms of the

characteristics of the MJO since phase 3 (CMIP3; [Hung et al. 2013](#)). [Hung et al. \(2013\)](#) evaluated the simulation of MJO in 20 CMIP5 models with comparison to CMIP3 and found that only one of 20 models can simulate realistic eastward propagation of MJO. [Ahn et al. \(2017\)](#) evaluated the performance of CMIP5 models in simulating MJO using process-oriented diagnosis and suggested that improvements on the processes associated with moisture and convection and the gross moist stability can boost the MJO simulation. Recently, [Ahn et al. \(2020\)](#) evaluated 34 models in CMIP6 on their abilities in reproducing the MJO propagation across the Maritime Continent in comparison with 25 models of CMIP5. They found that the CMIP6 models show realistically enhanced propagation across the Maritime Continent due to enhanced horizontal moisture advection. However, the comparison conducted by [Ahn et al. \(2020\)](#) was only based on the moisture budget analyses, which lacks depiction of large-scale dynamical and thermodynamical structure associated with the MJO.

In this study, we attempt to evaluate the eastward-propagating characteristics of the simulated MJO in CMIP5 and CMIP6 models in a comprehensive way by mainly applying dynamics-oriented diagnostics, along with several basic process-oriented diagnostics. These dynamics-oriented diagnostics include large-scale zonal circulation, vertical profiles of diabatic heating and equivalent potential temperature, and PBL moisture convergence, which can provide insight into whether the MJO reproduced by these models occurs for the right reasons. Instead of evaluating CMIP models for the purpose of classification, this study seeks to answer the questions whether CMIP6 models as a whole show a better representation of MJO propagation than CMIP5 models, and which MJO-related dynamic or thermodynamic structures or processes in CMIP6 models should be credited for the improvement, if any.

2. Data and method

a. Model and validation data

In this study, we analyze one ensemble member of historical runs, namely r1i1p1 for CMIP5 and r1i1p1f1 for CMIP6. This ensemble member is the first member of preindustrial control (piControl) experiment from the condition of 1 January 1850. For a fair comparison, we select 25 CMIP5 ([Taylor et al. 2012](#)) and 25 CMIP6 ([Eyring et al. 2016](#)) models for a common period of 1979–2005 (27 years in total). Model details are listed in [Table 1](#). The daily outputs of outgoing longwave radiation (OLR), precipitation, sea surface temperature (SST), specific humidity (q), horizontal winds, temperature, and vertical velocity in pressure coordinates (ω) are adopted.

The observational OLR daily data is derived from National Oceanic and Atmospheric Administration (NOAA) interpolated outgoing longwave radiation datasets for the period of 1979–2005 ([Liebmann and Smith 1996](#)). The observed daily precipitation and SST data used to evaluate models' seasonal means are from the Tropical Rainfall Measuring Mission (TRMM; [Huffman et al. 2007](#)) for the period of 1998–2005

and the Optimum Interpolation Sea Surface Temperature version 2 (OISSTv2; [Reynolds et al. 2007](#)) for the period of 1981–2005, respectively. Other atmospheric variables are obtained from European Centre for Medium-Range Weather Forecasts (ECMWF) ERA-Interim daily-averaged reanalysis data ([Dee et al. 2011](#)). The horizontal resolution of all models and validation data is interpolated into $2.5^\circ \times 2.5^\circ$. This study only examines the MJO occurring in the extended boreal winter from November to April.

b. Modified regression method

In this study, we employ the linear least squares regression method to isolate the MJO convection-related signal. Prior to regression, intraseasonal anomalies of each variable are obtained by applying a 20–100-day Lanczos bandpass filter. The MJO-related metrics are then produced by regressing the intraseasonal anomalies of interested variables onto the intraseasonal OLR anomalies averaged over the reference box in the equatorial eastern Indian Ocean (EIO; 10°S – 10°N , 70° – 100°E). To avoid the excessive impacts from contribution of maximum or minimum values near the reference box, we apply a min-max normalization method to the regression in a larger area (15°S – 15°N , 65° – 105°E) than the EIO reference box. That is, the values in that box are regressed onto normalized OLR intraseasonal anomalies, while those outside of the box are regressed onto nonnormalized ones. The min-max normalization method is shown below:

$$x' = \frac{x - \min(x)}{\max(x) - \min(x)},$$

where x' denotes the normalized values, and $\min(x)$ and $\max(x)$ represent the minimum and maximum values. After normalization, the OLR intraseasonal anomalies are rescaled between 0 and 1.

Given that the regression coefficients only reflect the negative or positive relationship of regressed fields to OLR in EIO, we multiply -1 standard deviation (STD) of observed OLR intraseasonal anomalies in the reference box to all regression coefficients for a better interpretation. As the -1 STD of OLR is always negative, it can represent the active convection over the reference box. Accordingly, we can simply interpret the regressed fields as the case when active convection is located at the EIO.

3. Results

a. Mean state

Previous studies have shown that biases in atmospheric mean states are highly likely to cause poor MJO statistics in simulations (e.g., [Gonzalez and Jiang 2017](#); [Jiang et al. 2020b](#); [Kim et al. 2011](#); [Ray et al. 2011](#)). We first evaluate the fidelity of the simulated mean states of OLR (as the proxy of convective precipitation), 850-hPa zonal wind (U850), and lower tropospheric moisture in extended boreal winter season from November to April ([Fig. 1](#)). As some recent studies argued that the air–sea interactions can be important to MJO

TABLE 1. List of CMIP5 and CMIP6 models used in this study.

No.	Model	Institute	Atmospheric horizontal resolution (lon $\times$ lat)	Cumulus scheme and modification
CMIP5 models				
1	ACCESS1.0	Centre for Australian Weather and Climate Research (CAWCR), Australia, Australia	$1.875^\circ \times 1.250^\circ$	Gregory and Rowntree (1990) Gregory and Rowntree (1990), Hewitt et al. (2011)
2	ACCESS1.3			
3	CanESM2	Canadian Centre for Climate Modeling and Analysis (CCCma), Canada	$2.813^\circ \times 2.813^\circ$	Zhang and McFarlane (1995)
4	CMCC-CESM	Centro Euro-Mediterraneo sui Cambiamenti Climatici (CMCC), Italy	$3.750^\circ \times 3.750^\circ$	Tiedtke (1989), Nordeng (1994)
5	CMCC-CM		$0.750^\circ \times 0.750^\circ$	
6	CMCC-CMS	Commonwealth Scientific and Industrial Research Organization (CSIRO), Australia	$1.875^\circ \times 1.875^\circ$	Gregory and Rowntree (1990), Gregory (1995)
7	CSIRO-Mk3.6.0		$1.875^\circ \times 1.875^\circ$	
8	FGOALS-g2	Institute of Atmospheric Physics, Chinese Academy of Sciences, China	$2.813^\circ \times 3.000^\circ$	Zhang and McFarlane (1995), Zhang and Mu (2005) Tiedtke (1989), Nordeng (1994)
9	FGOALS-s2		$2.813^\circ \times 1.667^\circ$	
10	GFDL CM3	Geophysical Fluid Dynamics Laboratory (GRDL), National Oceanic and Atmospheric Administration (NOAA), United States	$2.500^\circ \times 2.000^\circ$	Donner (1993), Donner et al. (2001), Wilcox and Donner (2007) Moorthi and Suarez (1992)
11	GFDL-ESM2G			
12	GFDL-ESM2M	Institut Pierre-Simon Laplace (IPSL), France	$3.750^\circ \times 1.875^\circ$	Emanuel (1991)
13	IPSL-CM5A-LR		$2.500^\circ \times 1.268^\circ$	
14	IPSL-CM5A-MR		$3.750^\circ \times 1.875^\circ$	
15	IPSL-CM5B-LR	Atmosphere and Ocean Research Institute/ National Institute for Environmental Studies/JAMSTEC, Japan	$0.563^\circ \times 0.563^\circ$	Grandpeix and Lafore (2010), Grandpeix et al. (2010) Arakawa and Schubert (1974), Pan and Randall (1998), Emori et al. (2001) Chikira and Sugiyama (2010) Pan and Randall (1998), Emori et al. (2001) Tiedtke (1989), Nordeng (1994)
16	MIROC4h			
17	MIROC5			
18	MIROC-ESM-CHEM			
19	MIROC-ESM			
20	MPI-ESM-LR	Max Planck Institute for Meteorology (MPI), Germany	$1.875^\circ \times 1.875^\circ$	Tiedtke (1989), Nordeng (1994)
21	MPI-ESM-MR			
22	MPI-ESM-P			
23	MRI-CGCM3	Meteorological Research Institute (MRI), Japan	$1.125^\circ \times 1.125^\circ$	Arakawa and Schubert (1974), Tiedtke (1989), Yoshimura et al. (2015) Arakawa and Schubert (1974), Yoshimura et al. (2015), Kain and Fritsch (1990, 1993)
24	MRI-ESM1			
25	NorESM1-M	Norwegian Climate Centre (NCC), Norwegian	$2.500^\circ \times 1.895^\circ$	Zhang and McFarlane (1995), Neale et al. (2008), Richter and Rasch (2008)
CMIP6 models				
1	ACCESS-CM2	Centre for Australian Weather and Climate Research (CAWCR), Australia	$1.875^\circ \times 1.250^\circ$	Gregory and Rowntree (1990)
2	ACCESS-ESM1-5			
3	AWI-ESM-1-1-LR	Alfred Wegener Institute (AWI), Helmholtz Centre for Polar and Marine Research, Germany	$1.875^\circ \times 1.875^\circ$	Tiedtke (1989), Nordeng (1994)
4	BCC-ESM1	Beijing Climate Center (BCC), China	$2.813^\circ \times 2.813^\circ$	Wu (2012)
5	CanESM5	Canadian Centre for Climate Modeling and Analysis (CCCma), Canada	$2.813^\circ \times 2.813^\circ$	Zhang and McFarlane (1995)
6	CESM2	NSF-DOE-NCAR, United States	$1.250^\circ \times 0.938^\circ$	Zhang and McFarlane (1995)
7	CESM2-FV2		$2.500^\circ \times 1.875^\circ$	
8	CESM2-WACCM		$1.250^\circ \times 0.938^\circ$	

TABLE 1. (Continued)

No.	Model	Institute	Atmospheric horizontal resolution (lon $\times$ lat)	Cumulus scheme and modification
9	CESM2-WACCM-FV2		$2.500^\circ \times 1.875^\circ$	
10	CMCC-CM2-HR4	Centro Euro-Mediterraneo sui Cambiamenti Climatici (CMCC), Italy	$1.250^\circ \times 0.938^\circ$	Zhang and McFarlane (1995)
11	EC-Earth3-AerChem	Rosby Centre, Swedish Meteorological and Hydrological Institute, European	$0.703^\circ \times 0.703^\circ$	Tiedtke (1989)
12	EC-Earth3-Veg	EC-Earth consortium, Europe	$1.125^\circ \times 1.125^\circ$	
13	EC-Earth3-Veg-LR	EC-Earth consortium, Europe	$1.125^\circ \times 1.125^\circ$	
14	FGOALS-g3	Institute of Atmospheric Physics, Chinese Academy of Sciences, China	$2.000^\circ \times 2.250^\circ$	Zhang and McFarlane (1995), Zhang and Mu (2005), Wu et al. (2007)
15	GFDL-CM4	Geophysical Fluid Dynamics Laboratory (GRDL), National Oceanic and Atmospheric Administration (NOAA), United States	$1.000^\circ \times 1.000^\circ$	Arakawa and Schubert (1974), Bretherton et al. (2004)
16	INM-CM4-8	Institute of Numerical Mathematics (INM), Russia	$2.000^\circ \times 1.500^\circ$	Betts (1986)
17	INM-CM5-0			
18	IPSL-CM6A-LR	Institut Pierre-Simon Laplace (IPSL), France	$2.500^\circ \times 1.250^\circ$	Emanuel (1991)
19	MIROC6	Atmosphere and Ocean Research Institute/ National Institute for Environmental Studies/JAMSTEC, Japan	$1.406^\circ \times 1.406^\circ$	Chikira and Sugiyama (2010)
20	MPI-ESM-1-2-HAM	Max Planck Institute for Meteorology (MPI), Germany	$1.875^\circ \times 1.875^\circ$	Tiedtke (1989), Nordeng (1994), Stevens et al. (2013)
21	MPI-ESM1-2-HR		$0.938^\circ \times 0.938^\circ$	
22	MPI-ESM1-2-LR		$1.875^\circ \times 1.875^\circ$	
23	MRI-ESM2-0	Meteorological Research Institute (MRI), Japan	$1.125^\circ \times 1.125^\circ$	Arakawa and Schubert (1974), Yoshimura et al. (2015), Kain and Fritsch (1990, 1993)
24	NESM3	Nanjing University of Information Science and Technology (NUIST), China	$1.875^\circ \times 1.875^\circ$	Tiedtke (1989), Nordeng (1994), Stevens et al. (2013)
25	SAM0-UNICON	Seoul National University (SNU), South Korea	$1.250^\circ \times 0.938^\circ$	Zhang and McFarlane (1995), Neale et al. (2008), Park and Bretherton (2009)

propagation (e.g., DeMott et al. 2015, 2016; Fu et al. 2017; Zhu et al. 2017), the simulated mean state of SST is also examined in Fig. 1.

In the observations, the relatively low OLR (Fig. 1a-1) and intensive lower-tropospheric moisture (Fig. 1c-1) are expected to respond to high SST (Fig. 1d-1) over the Indo-western Pacific sector, indicative of active convections over these regions during boreal winter season. Observed mean U850 shows relatively weak westerlies (up to 4 m s^{-1}) over the tropical Indian Ocean, Maritime Continent region, and western Pacific, while predominant easterlies are found over other regions (Fig. 1b-1). Compared to observation, the OLR in CMIP6 is generally overestimated except for small areas in the western-central tropical Indian Ocean and eastern Maritime Continent (Fig. 1a-2), suggesting an overall underestimated convective rainfall over tropical Indo-Pacific, particularly over the Pacific cold tongue region. Over the eastern tropical Indian Ocean, the Maritime Continent, and around 15°S off the equatorial tropical Pacific, the OLR is significantly lower in CMIP6 than that in CMIP5 (Fig. 1a-3), indicating relatively strong mean convection over these areas in CMIP6. The CMIP6 models show an overall easterly bias of U850 except for a westerly bias found over the southern

area of the Maritime Continent (Fig. 1b-2). Such westerly biases are much stronger than those in CMIP5 models (Fig. 1b-3). The mean state of lower-tropospheric moisture shows dry bias over the eastern Indian Ocean and the Pacific cold tongue in CMIP6 models (Fig. 1c-2), which can be a result of cold bias of SST over these areas (Fig. 1d-2). It is evident that, compared to CMIP5, the seasonal mean SST is warmer (Fig. 1d-3) and the lower atmosphere is wetter (Fig. 1c-3) in CMIP6, which can provide more moisture to favor the propagation of the MJO (Arnold et al. 2015).

To assess the biases of mean states, Fig. 2 shows the scatterplots of normalized root-mean-square error (NRMSE) against the pattern correlation coefficient (PCC) for each individual model and their multimodel means. The multimodel means (MMMs) for CMIP5 and CMIP6 models are denoted as CMIP5-MMM and CMIP6-MMM hereafter. The values of the PCC and NRMSE are calculated based on the domain of 20°S – 20°N , 30° – 210°E as shown in Fig. 1 and the NRMSE is obtained by dividing the RMSE with one standard deviation of the observed value to facilitate the comparison between models with different scales. As expected, the model with a higher PCC tends to have a lower NRMSE. The performances of simulating mean states of OLR (Fig. 2a) and

FIG. 1. (a-1)–(d-1) The mean states of extended boreal winter seasonal (November–April) for observational and reanalysis, (a-2)–(d-2) mean-state biases in winter for CMIP6 models, and (a-3)–(d-3) Difference of mean-state in winter season between CMIP6 and CMIP5 models in terms of OLR, 850-hPa zonal wind (U850), 850–700-hPa averaged specific humidity (q), and SST. The statistically significant biases and differences at 95% confidence level are denoted by black dots. The Pacific equatorial cold tongue region is denoted on plots in the center column.

U850 (Fig. 2b) are comparable among CMIP5 and CMIP6 models given similar spreads of PCCs and NRMSE. However, CMIP6 models show improved mean states of lower atmospheric moisture (Fig. 2c) and SST (Fig. 2d) due to models with higher PCCs and lower NRMSEs clustered.

b. Wavenumber–frequency power spectra

It is known that the tropical convective rainfall can be organized to move eastward or westward as a form of convectively coupled equatorial waves (CCEWs) with a wide range of spatial and temporal scales (Kiladis et al. 2009). Through a space–time spectral analysis of the OLR, the CCEWs can be decomposed into some symmetric and antisymmetric components (Fig. 3). In the symmetric components of the observed OLR spectrum, strong eastward waves can be identified as the forms of the Kelvin waves (~ 2 – 10 zonal wavenumbers, ~ 3 – 16 days) and the MJO-related waves (~ 1 – 3 zonal wavenumbers, ~ 30 – 90 days), and relatively weaker westward $n = 1$ equatorial Rossby waves are shown as a spatial–temporal variation of ~ 2 – 5 zonal wavenumbers and period > 20 days (Fig. 3a). In models, the symmetric waves are overall underestimated. However, the CMIP6-MMM does show some improvement on the simulations of eastward Kelvin wave (Fig. 3c) compared to its CMIP5-MMM counterpart (Fig. 3b), despite faster propagation speeds than observation. In the antisymmetric components, strong mixed Rossby–gravity waves and $n = 0$ eastward inertia–gravity waves are also identified in the observation (Fig. 3d), which can be found in CMIP6-MMM, albeit with much weaker power (Fig. 3f), but they are absent in CMIP5-MMM (Fig. 3e).

To isolate the spatial and temporal signals that are related to the MJO, we apply wavenumber–frequency power spectra on the OLR and U850 anomalies in winter (Figs. 4a–c). The power spectra are operated on each year and then averaged over all years of data. The dominant eastward powers for both OLR and U850 in observation are shown in 1–3 zonal wavenumbers and 0.01–0.03 cycles per day (i.e., ~ 30 – 100 days). The patterns of spectra for CMIP5- and CMIP6-MMM are similar; that is, the eastward power of OLR is prone to concentrate at 1–4 zonal wavenumbers and low-frequency zone (period > 100 days), whereas that of U850 is dominant at 1–2 zonal wavenumbers and ~ 30 – 100 days (Figs. 4b,c). However, the magnitudes of the eastward powers are higher in CMIP6-MMM for both OLR and U850. This may indicate stronger eastward signals of the simulated MJO in CMIP6-MMM.

To quantify the capability of eastward propagation, we calculate the ratios of eastward power over westward power (i.e., E/W herein), the ratios of simulated eastward power over observed eastward power (i.e., E/O), and the coherence (Coh^2) between OLR and U850. The E/W ratio suggests the relative strength of eastward power to its westward counterpart in observation or models, whereas the E/O ratio indicates how strong the eastward power in model compared to that in the observation. The Coh^2 obtained by a cross spectrum shows the coupling relationship between OLR and U850. As the values of Coh^2 are between 0 and 1, we normalize the E/W and E/O ratios using the min-max normalization method to fit all metrics in one heat map shown in Fig. 4d. The warmer color in Fig. 4d indicates stronger eastward propagation power or tighter coupling relationship between OLR and

FIG. 2. Scatterplot of pattern correlation coefficient (PCC) and normalized RMSE (NRMSE) for the mean state in winter of (a) OLR, (b) 850-hPa zonal wind, (c) 1000–700-hPa averaged specific humidity, and (d) SST based on CMIP5 (blue) and CMIP6 (red) simulations. In each plot, large hollow circles represent the multimodel means.

U850. Through this heat map, the models with strong or weak eastward powers can be easily identified. For example, it is obvious that the eastward power and the OLR–U850 coupling relationship are relatively strong in observation, and the CMIP6-MMM shows a general improvement from CMIP5-MMM in these aspects. We can also find the poor simulations, such as MIROC-ESM-CHEM, MIROC-ESM, and CanESM2 in CMIP5 and INM-CM4-8, INM-CM5-0, and IPSL-CM6A-LR in CMIP6.

c. MJO eastward propagation skill

The slow eastward propagation of the active convective center at a speed of $\sim 5 \text{ m s}^{-1}$ is one of the most prominent features of the MJO, which is used to reflect how well the model can reproduce realistic MJO (e.g., Jiang et al. 2015; Wang and Lee 2017; Wang et al. 2018). Conventionally, the lag–longitude diagrams of 10°N – 10°S averaged intraseasonal OLR anomalies regressed onto the EIO intraseasonal OLR

anomalies are a good way to depict the propagation of the MJO whose convective center is in the EIO (Figs. 5a–c). In the observations (Fig. 5a), the convection starts approximately 10 days preceding the maximum convection in the reference box and maintains a continuous eastward propagation from 60°E to the vicinity of 180° with a speed of $\sim 5 \text{ m s}^{-1}$ (i.e., propagating almost 120° in 40 days). The convection in CMIP5-MMM initiated near 60°E shows 5 days delay compared to the observation, which takes ~ 25 days to propagate eastward from 60° to 130°E with an average speed of $\sim 2.55 \text{ m s}^{-1}$ (Fig. 5b). In contrast, the propagation speed of the convection in CMIP6-MMM is comparable to observation, although its convective center fails to propagate as far as that in the observation (Fig. 5c). The intermodel comparison based on individual lag–longitude diagrams suggests that the simulated propagation speed has been generally improved from CMIP5 to CMIP6 models (see Figs. S1 and S2 in the online supplemental material). Although the Maritime Continent barrier exists in

FIG. 3. (a)–(c) Symmetric and (d)–(f) antisymmetric space–time OLR spectra for observation (NOAA), CMIP5-MMM, and CMIP6-MMM, respectively. The spectra are operated on the daily OLR over the entire tropical channel within the latitudes of 15°S – 15°N . The symmetric and antisymmetric components are displayed as the ratios relative to the background spectra. The x axis represents the zonal wavenumber with positive (negative) values indicating eastward (westward) propagation. The y axis represents the frequency in units of cycle per day (cpd).

both the CMIP5 and CMIP6 models, it is nevertheless weaker in CMIP6 models. This result is in agreement with the findings from Ahn et al. (2020).

To evaluate how well each model can reproduce the characteristics of the MJO eastward propagation, we calculate the PCCs between the simulated lag–longitude diagrams and that of the observation (Fig. 5d). It is argued that the conventional PCCs based on the regression coefficients can be dominant by the weights near the reference box, possibly leading to unreliable high values of PCCs. Such that, we also calculate the weights-removed PCCs by normalizing the regression coefficients in reference box before the pattern correlation applied. The results show that the modified PCCs that represent models' propagation skills are generally lower than those without normalization (not shown). However, the normalization shows limit influence on the ranks of models, namely, the good (poor) models retaining relatively high (low) PCC values. The correlation coefficients between these two sets of PCCs are significantly high (i.e., 0.91 for CMIP5 models and 0.97 for CMIP6 models). Thus, we still adopt the convective calculation of PCCs throughout this study to be consistent with previous studies on model intercomparison of MJO. It

is shown that there are 48% of the CMIP6 models having their PCCs exceeding 0.90% versus 28% of the CMIP5 models. In addition, the PCC of the multimodel mean of CMIP6 is higher (0.95) than that of CMIP5-MMM (0.92). In last section, the capability of simulated eastward power of OLR has been evaluated in the form of E/W ratio. Here, we calculate the correlation coefficients between PCCs and E/W ratios of OLR for CMIP5 and CMIP6 models (Fig. 5e). It shows that these two metrics are strongly positively correlated.

Given its simplicity, the PCC based on the OLR lag–longitude diagram is defined as the MJO eastward propagation skill of a model and used to explore the linear regression relationships against several MJO-associated processes in the following sections.

d. Zonal wind spatial structure in the lower troposphere

Observationally, the three-dimensional structure of the MJO reveals a tightly coupled relationship between deep convection and zonal wind (e.g., Chen and Wang 2018, 2020; Zhang 2005). The observed intraseasonal anomalies of U850 display a zonal asymmetry with easterlies to the east of the convective center and westerlies to the west (Fig. 6a).

FIG. 4. (a)–(c) The wavenumber–frequency power spectra of 10°S – 10°N averaged OLR (shaded) and U850 (contour) anomalies in winter season for observational/reanalysis data, CMIP5-MMM, and CMIP6-MMM. (d) The wavenumber–frequency spectra metrics based on the “MJO band” at 1–3 wavenumber and 30–60 days. A detailed description of these metrics is provided in section 3b.

Furthermore, the associated pair of cyclonic gyres straddling the equator can be found to the west of convective center in observation (e.g., [Adames and Kim 2016](#); [Wang et al. 2016](#); [Zhang 2005](#)). Such a large-scale U850 spatial pattern resembles a coupled Kelvin–Rossby wave structure, which has been considered essential to the MJO dynamics ([Wang et al. 2018](#)). The CMIP5- and CMIP6-MMM can reproduce such a U850 spatial pattern to some extent. Although the strength of easterly anomalies in CMIP6-MMM is slightly weaker than in ERA-Interim ([Fig. 6c](#)), it is stronger than that in CMIP5-MMM ([Fig. 6b](#)). We use the PCC of U850 wind speed in the domain of 15°S – 15°N , 30° – 180°E between the models and observation to represent the skill in simulating U850 spatial structure. It is shown that most CMIP models can reproduce reasonable U850 spatial structure (Figs. S3 and S4). The U850 spatial structure in the CMIP6-MMM shows a higher PCC (0.94; [Fig. 6c](#)) than that in CMIP5-MMM (0.89; [Fig. 6b](#)), suggesting an improved representation of the U850 spatial structure in CMIP6-MMM. To further examine the contribution of the U850 spatial structure to the MJO eastward propagation, we employ a linear regression of MJO propagation skill against U850 spatial structure ([Fig. 6d](#)). The CMIP6-MMM shows a higher regression coefficient (0.71) than that in CMIP5-MMM (0.55) with high positive correlation coefficient of 0.73, indicating a coherent and robust linear relationship between the

U850 spatial structure and the MJO eastward propagation in CMIP6 models.

Previous observational studies showed that the zonal asymmetrical structure of low-level zonal winds can be important to the propagation of the MJO. The zonal asymmetrical structure is characterized by the stronger easterlies to the east relative to westerlies to the west of the convective center at 850 hPa. In [Wang et al. \(2018\)](#), the zonal asymmetry of the low atmospheric zonal wind was examined using the ratio of maximum easterly speed versus maximum westerly speed averaged over the equatorial Indo-western Pacific region (5°S – 5°N , 30°E – 180°). Following [Wang et al. \(2018\)](#), we calculate this ratio to quantify the zonal asymmetry of U850. As the spatial structure of U850 resembles the Kelvin–equatorial Rossby wave structure responding to a diabatic heat source ([Gill 1980](#); [Matsuno 1966](#)), we refer to the easterly/westerly ratio as the K/R ratio to distinguish it from the E/W power ratio in section 3b. The K/R ratio in ERA-Interim is ~ 1.19 , indicative of relatively stronger easterlies than westerlies in U850. In contrast, the K/R ratios in both multimodel means are lower than 1.0 (i.e., 0.91 for CMIP5-MMM and 0.99 for CMIP6-MMM), suggesting weakened U850 zonal asymmetrical structure, and small K/R ratios (i.e., < 1.0) can be found across more than 60% of CMIP models ([Tables 2 and 3](#)). In

FIG. 5. (a)–(c) Lagged time–longitude diagrams of regressed intraseasonal OLR anomalies averaged over 10°S–10°N for observation, CMIP5-MMM, and CMIP6-MMM. Red dashed line indicates the 5 m s⁻¹ propagation speed. (d) The PCC between each model and observation is calculated based on their lag–longitude diagrams, which is defined as MJO eastward propagation skill. (e) Scatterplot of the MJO eastward propagation skill against E/W ratio. The correlations are also indicated in the plot.

addition, the low regression coefficients of the MJO propagation skill onto the simulated K/R ratio suggest that the K/R ratio alone might not be an effective contributor to the propagation of MJO in CMIP models.

e. MJO dynamics-related vertical structures

The MJO convective complex consists of multiscale dynamic and thermodynamic processes that couples convection with moisture, heating, and waves associated with large-

FIG. 6. The regressed intraseasonal anomalies of U850 wind speed (shaded) and 850-hPa winds (vector) during the extended winter season (November–April) for (a) ERA-Interim, (b) CMIP5-MMM, and (c) CMIP6-MMM. (d) Scatterplot of the MJO eastward propagation skill against the PCCs of the spatial pattern of U850 wind speed (defined as U850 spatial structure). Blue and red circles represent CMIP5 and CMIP6 models, respectively, with the large hollow ones representing the multimodel means. The correlations between models and observation are also indicated in the plot.

TABLE 2. Pattern correlations between CMIP5 model and observation with respect to a range of metrics, K/R ratios, and western Pacific convective instability indices (CIIs) for CMIP5 models and observation. The models are ranked in a descending order according to their MJO skill. The outliers are marked by an asterisk (*).

	CMIP5 models	MJO skill	U850 structure	Q1 structure	EPT structure	Moisture structure	Divergence structure	Zonal wind structure	Omega structure	MFC skill	∂_q skill	K/R ratio	Western Pacific CII
1	NorESM1-M	0.9534	0.9045	0.9029	0.8427	0.7890	0.7248	0.9226	0.9055	0.7270	0.7074	1.2900	0.1122
2	CMCC-CM	0.9225	0.8446	0.9226	0.8716	0.9076	0.6880	0.8485	0.9303	0.7673	0.9093	0.7183	0.2638
3	IPSL-CM5A-MR	0.9190	0.7260	0.8078	0.8160	0.7458	0.5172	0.7088	0.8268	0.5285	0.6521	1.5765	0.0147
4	IPSL-CM5B-LR	0.9147	0.8480	0.8992	0.8641	0.8939	0.6672	0.9096	0.9163	0.7387	0.9124	1.4811	0.2828
5	MRI-CGCM3	0.9138	0.9579	0.7447	0.7135	0.8300	0.6888	0.8749	0.7637	0.8107	0.8464	1.0857	0.1193
6	MRI-ESM1	0.9110	0.8116	0.7580	0.7082	0.8161	0.6652	0.8595	0.7746	0.8273	0.8189	0.9491	0.1514
7	GFDL-CM3	0.9094	0.9183	—	0.8988	0.8883	0.7347	0.8967	—	0.7975	0.7381	0.6903	0.1687
8	ACCESS1.0	0.8895	0.9007	0.9045	0.7371	0.8599	0.6733	0.8703	0.9106	0.7081	0.7646	0.8966	0.1524
9	MIROC5	0.8817	0.9125	0.8858	0.8173	0.7742	0.7497	0.9046	0.9000	0.7535	0.9005	0.7554	0.2373
10	ACCESS1.3	0.8802	0.9279	0.9419	0.8478	0.8779	0.7380	0.8946	0.9375	0.6545	0.6489	0.7365	0.1602
11	CSIRO-Mk3.6.0	0.8642	0.8468	0.7980	0.7760	0.8406	0.6285	0.9012	0.8154	0.6275	0.7677	0.9236	0.1162
12	FGOALS-s2	0.8562	0.8283	0.8871	0.7590	0.7701	0.6830	0.7958	0.8757	0.6059	0.6569	1.1019	0.1696
13	CMCC-CMS	0.8550	0.8442	0.8656	0.8819	0.8987	0.7305	0.8551	0.8692	0.7603	0.8757	0.7576	0.3210
14	IPSL-CM5A-LR	0.8524	0.6266	0.7493	0.7681	0.6824	0.4499	0.6825	0.7588	0.4816	0.5610	1.7327	0.0297
15	MPI-ESM-MR	0.8418	0.7458	0.8623	0.8168	0.8288	0.7380	0.7739	0.8651	0.6893	0.7889	0.7201	0.2349
16	GFDL-ESM2M	0.8356	0.8616	0.8607	0.8298	0.8589	0.7003	0.8738	0.8767	0.6809	0.7083	1.0336	0.2181
17	GFDL-ESM2G	0.8287	0.7670	0.7600	0.7902	0.7827	0.6355	0.8254	0.8122	0.6091	0.7040	0.9325	0.2290
18	MPI-ESM-LR	0.8209	0.8108	0.8725	0.8461	0.8757	0.7319	0.8264	0.8731	0.6531	0.7776	0.6489	0.1924
19	MPI-ESM-P	0.8165	0.8116	0.8602	0.8257	0.8459	0.7420	0.8138	0.8656	0.6467	0.7532	0.5715	0.2151
20	MIROC4h	0.8101	0.7726	0.6437*	0.7873	0.7580	0.6708	0.8544	0.6824	0.4761	0.7848	2.7308	-0.1063
21	MIROC-ESM	0.7795	0.7274	0.4446*	0.7597	0.8633	0.6700	0.8116	0.7000	0.5298	0.5891	0.8381	0.0322
22	MIROC-ESM-CHEM	0.7540	0.6438	0.8419	0.7474	0.6239	0.6390	0.7402	0.8329	0.5205	0.4618	0.8969	0.2799
23	CMCC-CESM	0.7311	0.6573	0.7975	0.7636	0.8546	0.5966	0.7851	0.7862	0.6089	0.7590	0.5819	0.2689
24	FGOALS-g2	0.7211	0.7352	0.7811	0.8262	0.7697	0.6397	0.7865	0.7667	0.5151	0.6569	0.6795	0.0529
25	CanESM2	0.6285	0.6667	0.7668	0.7416	0.7296	0.4823	0.5973	0.7588	0.4238	0.5057	1.1082	0.0497
	CMIP5-MMM	0.9209	0.8894	0.9314	0.8935	0.9059	0.7628	0.9138	0.9265	0.7911	0.8528	0.9054	0.1692
	Observation	—	—	—	—	—	—	—	—	—	—	1.1938	0.3899
	Regression coefficient	—	0.5483	0.5173	0.4846	0.4200	0.4000	0.6486	0.5618	0.4758	0.3719	0.0163	0.1100
	Correlated to MJO skill	—	0.7175	0.4181	0.3502	0.3974	0.4238	0.6745	0.5338	0.7282	0.5948	0.0993	0.1473

TABLE 3. As in Table 2, but for CMIP6 models.

	CMIP6 models	MJO skill	U850 structure	Q1 structure	EPT structure	Moisture structure	Divergence structure	Zonal wind structure	Omega structure	MFC skill	$\partial q'$ skill	K/R ratio	Western Pacific CII
1	EC-Earth3-Veg	0.9451	0.8911	0.9446	0.8915	0.9160	0.7760	0.8902	0.9412	0.8189	0.8477	0.6264	0.1848
2	FGOALS-g3	0.9425	0.9230	0.8730	0.8829	0.8889	0.8067	0.8868	0.8752	0.7751	0.8176	0.9931	0.0646
3	CESM2-FV2	0.9420	0.9578	0.8549	0.8259	0.7487	0.8101	0.9252	0.9450	0.8297	0.8338	0.1472	0.1857
4	ACCESS-CM2	0.9356	0.9301	0.9374	0.8115	0.7209	0.7209	0.8616	0.9372	0.7920	0.8406	0.8644	0.0244
5	CESM2-WACCM-FV2	0.9346	0.9396	0.8673	0.8043	0.7243	0.8099	0.9170	0.9151	0.8026	0.8946	1.4774	0.0727
6	EC-Earth3-Veg-LR	0.9343	0.9204	0.9361	0.8863	0.9150	0.7631	0.9102	0.9272	0.8099	0.8222	0.7215	0.1100
7	ACCESS-ESM1-5	0.9331	0.9129	0.9402	0.8276	0.8705	0.7286	0.8805	0.9290	0.7541	0.6938	0.8345	0.0415
8	EC-Earth3-AerChem	0.9323	0.8299	0.9208	0.8597	0.8801	0.7171	0.7946	0.9137	0.7930	0.8020	0.9952	0.0607
9	CESM2-WACCM	0.9309	0.9393	0.9132	0.8662	0.8102	0.7787	0.9532	0.9333	0.8176	0.9112	1.4065	0.1138
10	SAM0-UNICON	0.9184	0.9484	0.8900	0.7954	0.7508	0.7486	0.8934	0.9332	0.8253	0.8942	1.1493	0.1058
11	GFDL-CM4	0.9122	0.9508	0.9162	0.8671	0.8880	0.8380	0.9249	0.9441	0.8340	0.8651	0.5980	0.2087
12	CESM2	0.9089	0.9246	0.9028	0.8326	0.7445	0.7772	0.9367	0.9236	0.7611	0.9023	1.5297	0.1924
13	MIROC6	0.8970	0.9188	0.9293	0.8278	0.8377	0.7442	0.9185	0.9322	0.7773	0.9353	0.7995	0.1411
14	NESM3	0.8847	0.8879	0.9164	0.9019	0.9252	0.8116	0.8715	0.9233	0.7239	0.8576	0.6875	0.2487
15	IPSL-CM6A-LR	0.8776	0.6952	0.9009	0.8088	0.8327	0.6649	0.8511	0.8463	0.5535	0.7034	1.3874	0.1547
16	BCC-ESM1	0.8772	0.7908	0.8508	0.8733	0.8690	0.7299	0.8096	0.8628	0.6673	0.7935	2.1155	0.7797
17	MRI-ESM2-0	0.8662	0.9463	0.9028	0.8175	0.8599	0.6613	0.8943	0.8840	0.7045	0.8119	1.2182	0.2108
18	MPI-ESM1-2-HR	0.8595	0.9038	0.9233	0.8416	0.8262	0.7116	0.8180	0.9309	0.7407	0.7656	0.7302	0.3148
19	CanESM5	0.8399	0.8164	0.8639	0.8270	0.8047	0.7003	0.8067	0.8574	0.5198	0.4359	0.8826	0.1286
20	AWI-ESM-1-1-LR	0.8241	0.8442	0.8745	0.8984	0.9073	0.7395	0.9036	0.8900	0.5399	0.7786	0.7914	0.2109
21	CMCC-CM2-HR4	0.8228	0.8721	0.8349	0.8399	0.7919	0.7547	0.8815	0.8518	0.4576	0.5831	1.4163	0.0158
22	MPI-ESM1-2-LR	0.7986	0.7882	0.8620	0.8422	0.8761	0.6561	0.8300	0.8710	0.5802	0.6942	0.7358	0.2687
23	MPI-ESM-1-2-HAM	0.7692	0.7642	0.8137	0.8103	0.7961	0.6664	0.8065	0.8100	0.5127	0.7130	0.6103	0.2003
24	INM-CM5-0	0.7341	0.7951	0.7600	0.7725	0.7770	0.6186	0.7487	0.7479	0.4579	0.5963	0.7456	0.0468
25	INM-CM4-8	0.6471	0.7056	0.7303	0.6190	0.7246	0.3742	0.7948	0.6838	0.4370	0.3742	0.4561	-0.0236
	CMIP6-MMM	0.9454	0.9405	0.9435	0.9090	0.9172	0.9361	0.9383	0.9486	0.8175	0.9322	0.9896	0.1859
	Observation	—	—	—	—	—	—	—	—	—	—	1.1938	0.3899
	Regression coefficient	—	0.7071	1.1432	0.8759	0.4250	0.6791	0.9172	1.0613	0.4850	0.4108	0.0613	0.0311
	Correlated to MJO skill	—	0.7341	0.8371	0.6670	0.3620	0.8410	0.6642	0.9057	0.8874	0.7776	0.3114	0.0631

FIG. 7. Longitude–pressure diagrams of the regressed intraseasonal anomalies of (a-1)–(a-3) diabatic heating ($Q1$; shaded) and the zonal circulation (m s^{-1} for zonal wind and 0.01 Pa s^{-1} for vertical velocity; vector), (b-1)–(b-3) divergence with positive value indicating divergence, and (c-1)–(c-3) equivalent potential temperature (EPT; shaded) and specific humidity (contour). (left) ERA-Interim, (center) CMIP5-MMM, and (right) CMIP6-MMM. All plots are averaged over 10°S – 10°N along the longitude from 30°E to 180° during the winter season (November–April). The PCCs of CMIP5-MMM and CMIP6-MMM with ERA-Interim are indicated in corresponding plots.

scale circulation (Wang et al. 2018). The vertical structure of the MJO envelope is characterized by a prominent east–west asymmetry with shallow cumuli to the east of the convective center and a westward spread of stratiform at upper atmosphere behind the convection (Jiang et al. 2020a). In the following sections, we diagnose and evaluate the performance of CMIP models in simulating several dynamic and thermodynamics fields that are essential to MJO dynamics in a form of a longitude–pressure diagram. The regressed pattern of each field in the longitude–pressure diagram is defined as its vertical structure.

1) DIABATIC HEATING AND ZONAL CIRCULATION

We first diagnose the vertical structures of diabatic heating ($Q1$) and zonal circulation associated with the MJO convection (Figs. 7a-1, 7a-2, and 7a-3). The $Q1$ is often estimated as a residual of the heat budget and is calculated following Yanai et al. (1973), shown as follows:

$$Q1 = c_p \frac{\partial T}{\partial t} - c_p (\omega s - \mathbf{V} \cdot \nabla T),$$

FIG. 8. Scatterplots of the MJO eastward propagation skill against PCCs of (a) diabatic heating, (b) zonal wind, (c) vertical velocity, (d) equivalent potential temperature, (e) specific humidity, and (f) divergence. For the scatterplots, blue and red circles represent CMIP5 and CMIP6 models, with large hollow ones representing their multimodel means, respectively. The regression coefficients (rc) and correlation coefficients (r) between models and ERA-Interim are indicated in each plot.

where c_p , T , t , and ω denote the specific heat at constant pressure, the temperature, the time, and the vertical velocity in pressure coordinates, respectively; $s = -(T/\theta)(\partial T/\partial t)$ is the static stability with θ being the potential temperature; $\mathbf{V}$ and ∇ represent the horizontal velocity vector and the horizontal gradient operator, respectively.

In Fig. 7a-1, ERA-Interim shows a strong diabatic heating over the MJO convective center and a relatively weak cooling ahead of the convection, which favors a large-scale zonal circulation preceding the MJO convection. To the west of the deep convection, the Q1 shows heating in the upper troposphere and cooling in the lower troposphere. Both CMIP5- and CMIP6-MMM show similar patterns of Q1 to the ERA-Interim, with strong heating in the convective center and moderate cooling to the east of convection, although the heating in the upper troposphere to the west of convection is absent in models (Figs. 7a-2 and 7a-3). The convective heating in the deep convection is prone to expand zonally in CMIP6-MMM, leading to excessive upward flow in the convective center and shortened zonal circulation behind the convection. Compared to CMIP6-MMM, the maximum cooling is in the vicinity of 130°E in CMIP5-MMM, which causes shortened zonal circulation ahead of convection (Fig. 7a-2). As the zonal circulation largely relies on the zonal atmospheric pressure

gradient, the weakened zonal gradient of atmospheric heating to the east of convection leads to weaker easterlies in the lower troposphere, which can explain the smaller K/R ratio of U850 in CMIP models.

To investigate the importance of Q1 and zonal circulation components to the MJO propagation skill, we calculate the PCCs of the vertical structures of Q1, zonal wind, and vertical velocity between models and the reanalysis. There are three models in CMIP5 removed from the regression due either to missing data (GFDL CM3 and MIROC4h) or to obvious model errors in the upper troposphere (MIROC-ESM). Compared to CMIP5 models, their CMIP6 counterparts show higher regression coefficients and strong positive correlation for all three fields (Figs. 8a-c), particularly the Q1 and vertical velocity. This indicates that strong linear relationships between the MJO eastward propagation skill and vertical structures of Q1 and zonal circulation are more coherent and robust in CMIP6 models.

2) DIVERGENCE

The large-scale zonally distribution of divergence can help to understand the coupling between convection and zonal circulation. In ERA-Interim, the maximum convergence occurs

at 700–500 hPa while strongest divergence is located at 200 hPa over the MJO convective center (Fig. 7b-1). A divergence in the lower troposphere paired with an upper-level convergence are found behind the MJO convective center. Ahead of the convective center, divergence is predominant except for a convergence over western Pacific occurring below 850 hPa. Compared to the reanalysis, CMIP models simulate lower positions of divergence in the upper troposphere (Figs. 7b-2 and 7b-3), which are probably attributed to the sparse pressure levels recorded in CMIP models. The CMIP5-MMM shows a weaker pair of divergence and convergence over the convective center than that in the CMIP6-MMM. In addition, the regression relationship between the MJO propagation skill and vertical structure of divergence is more robust in CMIP6 models (Fig. 8f).

3) EQUIVALENT POTENTIAL TEMPERATURE AND MOISTURE

In addition to the forcing from large-scale circulation, previous studies showed that an unstable atmosphere ahead of the MJO convection can also promote the eastward propagation of the MJO (e.g., Zhang 2005; Jiang 2017). The equivalent potential temperature (EPT), a measurement to explain the moist-adiabatic process, is related to the stability of a column of air in the atmosphere. Thus, we use the pressure–longitude diagram of the intraseasonal EPT anomalies to depict atmospheric stability associated with the MJO convection (Figs. 7c-1, 7c-2, and 7c-3). In Fig. 7c-1, the reanalysis shows a concave-eastward shape in the EPT vertical profile with relatively higher EPT at the lower troposphere extending eastward (i.e., the propagation direction). This vertical gradient of the EPT can trigger a destabilizing process to the east of the MJO convection that is driven by a buoyancy force arising from hydrostatic instability. The moisture at lower troposphere also shows an eastward extension (Fig. 7c-1), which suggests a premoistening process ahead of the MJO convection. These predestabilizing and premoistening processes can facilitate the eastward propagation of the MJO. However, there are no such processes found in the CMIP5-MMM (Fig. 7c-2). In contrast, CMIP6-MMM can reproduce these processes albeit with a weak amplitude (Fig. 7c-3). In addition, both CMIP5- and CMIP6-MMM can reproduce the dry and cold anomalies behind the convection.

Using the PCCs between the observed and simulated EPT and moisture profiles to represent the EPT and moisture vertical structures, we examine the contributions of the EPT and moisture vertical structures to the MJO propagation skill (Figs. 8d,e). The CMIP5 and CMIP6 models show similar linear regression coefficients for moisture vertical structure (Fig. 8e), while the linear regression relationship is more robust and coherent in the CMIP6 models than their CMIP5 counterparts with respect to EPT vertical structure (Fig. 8d).

To quantify the atmospheric instability ahead of the MJO convection, we calculate the convective instability index (CII) following Wang et al. (2018). In Wang et al. (2018), the CII was defined as the EPT difference between 850 and 400 hPa. In this study, the CII is calculated as the difference between

the 1000–700-hPa averaged EPT and 500-hPa EPT, due to limited pressure levels available and large amount of missing data found below 700 hPa in some CMIP models. Positive CII values suggest cooler and denser air aloft, which can be used to interpret instable atmosphere. As the instability prior to MJO convection can promote the eastward propagation, we calculate the CII averaged over the western Pacific (120°–150°E) that can be found in Tables 2 and 3. The western Pacific CII for reanalysis, CMIP5-MMM, and CMIP6-MMM are ~ 0.39 , ~ 0.17 , and ~ 0.19 , respectively. This suggests that the atmospheric instability ahead of the MJO convection is underestimated in CMIP models. However, there is no obvious linear relationship between the MJO propagation skill and western Pacific CII (Tables 2 and 3) for either CMIP5 or CMIP6 models.

In summary, the moistening and destabilizing processes prior to the active deep convection are important to the MJO propagation. The destabilizing forces ahead of the MJO convection are similar in CMIP5 and CMIP6 models but weaker than the observation, while the premoistening process is substantially improved in the CMIP6 models as increased moisture at the lower troposphere is found over the western Pacific (Fig. 7c-3).

f. Lagged time–longitude regression analyses

In addition to the vertical structures, it is also important to diagnose the propagation patterns of the variables associated with the MJO. In this section, we use the lagged time–longitude diagram of regressed anomalies to evaluate the simulated propagation of total precipitation, moisture convergence and moisture tendency in the lower troposphere (Fig. 9).

The lagged time–longitude patterns of precipitation (Figs. 9a-1, 9a-2, and 9a-3) are consistent with those of OLR (Figs. 5a-c), and CMIP6-MMM shows a general improvement in simulating precipitation eastward propagation from CMIP5-MMM. The lagged time series of precipitation averaged over 85°–95°E (where maximum precipitation occurs around the lag = 0) provide insight into the timing of precipitation at this site (Fig. 9a-4). The negative (positive) lag can be interpreted as the timing before (after) the MJO convective center enters the site of 85°–95°E. It is obvious that models underestimated maximum precipitation near lag = 0, especially CMIP5-MMM. The peak precipitation of observation occurs ~ 2 days lagged, while CMIP5- and CMIP6-MMM show their maximum precipitation at lag = 0. At this site, precipitation anomalies of observation become positive ~ 9 days earlier than the MJO convective center, and CMIP5-MMM (CMIP6-MMM) shows ~ 11 days (~ 10 days) earlier. In addition, the duration of positive anomalies of precipitation for CMIP5-MMM at this site is longer than observation and CMIP6-MMM.

Previous studies suggested that the moisture convergence occurring at the PBL is crucial to the moistening process ahead of MJO convection (e.g., Maloney and Hartmann 1998; Benedict and Randall 2007; Wang et al. 2016, 2018; Hu et al. 2020). Due to limited pressure levels and missing data distributed over the Maritime Continent region in some models, we consider the vertical averaged moisture flux convergence

FIG. 9. (a-1)–(a-4) The lagged time–longitude diagrams of the regressed anomalies of total precipitation for observation/reanalysis, CMIP5-MMM and CMIP6-MMM, and the lead-lag time series of regressed anomalies averaged over 85° – 95° E with observation in black, CMIP5 in blue and CMIP6-MMM in red. (b-1)–(b-4), (c-1)–(c-4) As in (a-1)–(a-4), but showing the regressed anomalies of 1000–700-hPa vertically averaged moisture flux convergence and moisture tendency, respectively. All lag–longitude diagrams are averaged over 10° S– 10° N and displayed along the longitude of 30° E– 180° . Red dashed lines represent the propagation speed of 5 m s^{-1} . For moisture convergence plots, positive (negative) values indicate convergence (divergence). The PCCs between the multimodel means and observation/reanalysis are indicated.

from 1000 to 700 hPa as the PBL moisture convergence. In the observation, the PBL moisture convergence shows a similar propagation pattern to that of precipitation (Fig. 9b-1). In addition, a strong and persistent convergence can be found over the western Pacific when the MJO convection occurs in the EIO (Fig. 9b-1). Both CMIP5-MMM (PCC = 0.79) and CMIP6-MMM (PCC = 0.82) can simulate the eastward propagation of the PBL moisture convergence (Figs. 9b-2 and 9b-3). The moisture convergence over the western Pacific ahead of MJO convection can also be found in CMIP6-MMM, albeit

with weaker amplitude (Fig. 9b-3), while it is absent in CMIP5-MMM (Fig. 9b-2). We calculate the linear regression of the MJO propagation skill against the PCCs of PBL moisture convergence (shown in Tables 2 and 3), and the regression coefficient is 0.48 for both CMIP5 and CMIP6 models, suggesting similar contributions to premoistening process from the PBL moisture convergence. Furthermore, the high correlation coefficients (i.e., $r = 0.73$ for CMIP5 and $r = 0.89$ for CMIP6) indicate that this result is robust in models. An examination of lagged time series of PBL moisture

FIG. 10. The longitudinal variation of 10°S–10°N averaged regressed intraseasonal anomalies of (a) total precipitation, (b) SST, (c) latent heat flux, (d) specific humidity, (e) divergence, and (f) moisture tendency. The specific humidity, divergence, and moisture tendency are vertically averaged from 1000 to 700 hPa. In each panel, thick black line represents observation or reanalysis, and thin blue and red lines represent CMIP5-MMM and CMIP6-MMM, respectively. Blue and yellow strips mark the locations of EIO reference box used for regression and western Pacific region.

convergence shows that the maximum moisture convergence in CMIP5- and CMIP6-MMM is earlier than that in observation, and the moisture convergence in CMIP5-MMM occurs earlier and is stronger than CMIP6-MMM (Fig. 9b–4). We also note that the PBL moisture convergence occurs approximately 1 day earlier than precipitation no matter in observation or multimodel means.

According to the moist-mode theory, a positive moisture tendency at the low-level atmosphere prior to the MJO convection can promote the propagation of the MJO (e.g., Sobel and Maloney 2013; Adames and Wallace 2015; Adames and Kim 2016; Raymond and Fuchs 2009; Hsu et al. 2014; Hu et al. 2020). The moisture tendency is a temporal derivative of moisture, which indicates the rate of moistening or drying. Here, we use the 1000–700-hPa averaged intraseasonal regressed moisture tendency to depict the moistening process associated with the MJO. In Fig. 9c–1, the observation shows a positive moisture tendency occurring 10 days in advance of the MJO convection. When the MJO convective center locates in the eastern Indian Ocean, there exists a positive tendency over the western Pacific and a negative tendency to the west of convective center. The CMIP6-MMM can simulate a better positive tendency to the east of MJO convective center and east–west asymmetry of the moisture tendency (Fig. 9c–3) than the CMIP5-MMM (Fig. 9c–2). We also calculate the PCC of moisture tendency lagged time–longitude

diagram between the observation and models (Tables 2 and 3) and examine its contribution to the MJO propagation skill by using regression analysis. The results show a similar regression coefficient of 0.37 for CMIP5 ($r = 0.59$) and 0.41 for CMIP6 ($r = 0.78$) models, which suggests that the contributions from moisture tendency at lower troposphere are similar in CMIP5 and CMIP6 models. The lagged time series show that the peak of positive moisture tendency occurring at 10 days earlier and the moisture tendency reverse to negative ~1–2 days before the MJO convection in reanalysis and CMIP6-MMM (Fig. 9c–4), whereas the moisture tendency in CMIP5-MMM shows a few days of delay in peak and reversal timing.

g. East–west zonal asymmetry

One of the prominent characteristics of the MJO structure is the east–west zonal asymmetry. In this section, we examine the zonal distribution of several interested variables when the MJO convective center is at the EIO in observation and CMIP5- and CMIP6-MMM (Fig. 10). In observation, to the west of the MJO convection, increased positive anomalies of precipitation are found to extend from 50°E to the convective center, which is probably contributed by the upper-tropospheric stratiform precipitation (Fig. 10a). To the east of the MJO convection, the observed precipitation retains positive anomalies over the western Pacific, which can be related to

the precipitation from shallow cumuli. Compared to observation, CMIP5- and CMIP6-MMM show positive precipitation anomalies to the west of convection, albeit with shorter distance and weaker amplitude (Fig. 10a). However, to the east of convection, only CMIP6-MMM reproduces the comparable positive anomalies to the observation.

To the west of MJO convection, the observed SST (Fig. 10a) shows cold anomalies as a result of increased anomalies of precipitation (Fig. 10b) and upward latent heat flux (Fig. 10c). In contrast, the warm anomalies of SST are observed over the western Pacific, due to decreased amplitude of positive precipitation anomalies and increased downward latent heat flux. The SST and latent heat flux in CMIP5- and CMIP6-MMM shows similar longitudinal variation to observation. Compared to CMIP5-MMM, the SST anomalies are colder to the west of convection and warmer to the east of convection in CMIP6-MMM (Fig. 10b).

In the previous section, a strong observed PBL moisture flux convergence can be found to the east of the MJO convection. The CMIP6-MMM can reproduce such moisture flux convergences, while the CMIP5-MMM cannot. It can be partly explained by increased positive anomalies of moisture (Fig. 10d) along with enhanced convergence (Fig. 10e) in the lower troposphere over the western Pacific in CMIP6-MMM compared to CMIP5-MMM. As the moisture flux convergence is contributed by the horizontal advection of moisture and convergence of moisture, enhanced convergence with more moisture can increase the convergence of moisture. In the MJO convective center, the simulated convergence of moisture in CMIP6-MMM is even stronger than the observation.

In Fig. 10f, the moisture tendency at the lower troposphere shows an east–west asymmetry in observation with increased tendency to the east of the MJO convection and decreased tendency to the west of the MJO convection. The CMIP5- and CMIP6-MMM can capture a similar zonal asymmetry of moisture tendency to the observation, although the moistening (drying) process to the east (west) of the convective center is underestimated. Compared to CMIP5-MMM, the enhanced east–west asymmetry in CMIP6-MMM is mainly contributed by the increased amplitude of negative moisture tendency to the west of the MJO convection; that is, the drying processes behind the MJO convective center is improved in CMIP6-MMM.

To further examine the specific processes that cause the east–west asymmetry of moisture tendency, a diagnosis on moisture budget was applied. Following Yanai et al. (1973), the moisture equation on the intraseasonal time scale can be written as follows:

$$\partial_t q' = -(\omega \partial_p q)' - (u \partial_x q)' - (v \partial_y q)' - \frac{Q'_2}{L},$$

where q , ω , u , and v denote the specific humidity, zonal velocity, meridional velocity, and vertical pressure velocity, respectively; Q_2 represents the apparent moisture sink, and L ($= 2.5 \times 10^6 \text{ J kg}^{-1}$) is the latent heat due to condensation or evaporation processes. The prime indicates the

FIG. 11. The 1000–700-hPa vertically averaged moisture budget components for the (a) east (10°S – 10°N , 135° – 150°E) and (b) west (10°S – 10°N , 45° – 60°E), and (c) their differences. Models' standard deviation error bars are indicated in each panel.

intraseasonal variability. The terms on the right-hand side of the equation represent the changes of moisture due to vertical advection, zonal advection, meridional advection, and the moistening due to condensational heating and evaporative cooling, respectively.

Each moisture budget term in the area with the maximum amplitude of increased (decreased) moisture tendency (shown in Fig. 10f) is averaged to represent the value in the east (west) pole. It is shown that models display quite large dispersions from their multimodel means in almost all budget terms (Fig. 11). The positive tendency to the east of convective center is mainly contributed by the vertical moisture advection in observation and CMIP6-MMM (Fig. 11a). For CMIP5-MMM, the positive tendency in the east pole results from the meridional moisture advection and it is surprising that vertical moisture advection is found to be negative in CMIP5-MMM. The meridional moisture advection in east pole is enhanced in CMIP6-MMM, probably due to the sharpened gradient of mean moisture at lower troposphere. It is worth noting that

FIG. 12. Schematic diagram of three-dimensional MJO structure when active convective center is in the EIO. Possible processes associated with the MJO eastward propagation skills are marked by orange circles. The coefficients from the linear least squares regression of the MJO eastward propagation skill onto each process in the CMIP5 and CMIP6 models are indicated. This diagram is modified based on Fig. 1 of Wang et al. (2016).

the moisture advection via zonal wind in the east area is substantially underestimated in both CMIP5 and CMIP6 models.

In the west pole, the negative moisture tendency is contributed by moisture advection from all directions in observation, while for CMIP5- and CMIP6-MMM, the vertical advection plays a primary role. The meridional moisture advection for CMIP6-MMM is largely reduced. The amplified east–west asymmetry of moisture tendency in CMIP6-MMM is mainly contributed by enhanced drying processes occurred at the west pole.

4. Conclusions and discussion

This study evaluated the eastward propagation skill of the MJO in CMIP5 and CMIP6 models and examined several possible processes that can cause atmospheric instability and propagation of the MJO in the CMIP models. It is evident that the CMIP6 models show an overall improvement in simulating the eastward propagation skill compared to their CMIP5 counterpart, with more realistic propagation speed and longer propagation distance (Fig. 5). Such an improvement can be partly explained by the better simulated low-level mean moisture in CMIP6 with enhanced moisture distributed over the southern Maritime Continent that leads to reinforced horizontal mean moisture gradient (Gonzalez and Jiang 2017). The improved propagation in CMIP6 also benefits from the warmer mean SST and low-level equatorial westerlies rather than easterlies over the tropical Indo-Pacific sector (DeMott et al. 2015). We also analyzed the linear regression relationship of the eastward propagation of the MJO against some dynamic or thermodynamic structures

or processes at intraseasonal scale to ensure that such an improvement in CMIP6 models is from right reasons.

The first process is associated with the large-scale zonal circulation, in which the east–west asymmetry of wave-related zonal winds is considered important to the propagation of MJO. The observation suggests stronger easterlies anomalies with a K/R ratio of ~ 1.19 . Most CMIP5 and CMIP6 models show lower K/R ratios than observation, and no linear regression relationship is found between the K/R ratio and the simulated MJO propagation skill. For instance, the K/R ratios in some models are greater than 1.0 (i.e., relatively stronger easterlies to westerlies) or comparable to the observation, but their propagation skills are not well simulated (Figs. S1 and S2), such as the CanESM2 in CMIP5 and the BCC-ESM1 in CMIP6. However, these models show poor U850 spatial structures in common. It is found that most models with relatively low pattern correlations with the observed U850 spatial structure (Tables 2 and 3) tend to poorly simulate the eastward propagation of the MJO. The spatial structure of U850 can reveal how well it represents the low pressure associated with the Kelvin wave to the east of the convection and a pair of twin low pressure systems associated with the Rossby wave straddling the equator to the west (Wang et al. 2018), thus being proved to be more important than the K/R ratio itself in our study.

Wang et al. (2016) suggested that the enhanced Kelvin easterlies in model can facilitate the PBL convergence and formation of shallow cumuli to the east of the MJO convection, which boosts moisture buildup via frictional CISK mechanism, therefore promoting the eastward propagation of MJO. The CMIP6 shows strengthened Kelvin-wave easterlies at 850

FIG. 13. Scatterplots of (a) standard deviation against the mean of all PCC metrics and (b) the linear regression of the MJO eastward propagation against the mean of all PCC metrics.

hPa than CMIP5, which is also supported by stronger signal of Kelvin wave found in the space–time U850 spectra in CMIP6. Such improvements may benefit better simulations on diabatic heating vertical structure and PBL moisture convergence in CMIP6.

In CMIP6, the diabatic heating over the deep convective center shows a slightly zonal expansion, and moderate cooling occurs over the western Pacific, leading to a large-scale zonal circulation. In contrast, the maximum cooling in CMIP5 is in the vicinity of 130°E, which causes the shortened circulation to the east of deep convective center. In addition, the regression coefficient between diabatic heating and MJO propagation is substantially higher in CMIP6 models, which indicates that the increased fraction of variance across CMIP6 models can be explained by this regression. In other words, the CMIP6 models with proper simulated diabatic heating are likely to have better representation of MJO eastward propagation. Previous studies have shown that the vertical structure of diabatic heating in numerical models are predominated by cumulus parameterization (Seo and Wang 2010; Seo and Son 2012). The better representation of diabatic heating may be a result from improved cumulus parameterization in CMIP6.

Improved PBL moisture convergence can also contribute to the better representation of zonal circulation. For CMIP6, the PBL moisture convergence is enhanced over the western Pacific, as are the lower-tropospheric averaged moisture and divergence compared to CMIP5. The reinforced moisture convergence in CMIP6 can facilitate the coupling of Rossby and Kelvin wave in a Matsuno–Gill response and generating a zonal planetary-scale instability, which may reinforce the easterlies in the zonal circulation. Enhanced Kelvin wave easterlies to the east of MJO deep convection may contribute to the eastward propagation through inducing the premoistening effect by the vertical and horizontal moisture advections (Wang et al. 2017).

Although convective instability to the east of the convection is generally weaker in models than that in the observation, it is slightly improved in CMIP6 models. The CMIP6 models also show better representation of the premoistening processes occurring at the PBL. This may explain why CMIP6 models can better capture the characteristics of eastward propagation of the MJO than the CMIP5 models. Our results attribute the better premoistening in the CMIP6 models to the better representations of the PBL moisture convergence. The moisture convergence can be partitioned to the convergence of moisture and the meridional and zonal advections of moisture. Over the western Pacific, the convergence of moisture is increased due to enhanced lower-tropospheric moisture and divergence, and the meridional advection of moisture is enhanced, which may result from the sharpened meridional gradient of mean low-tropospheric moisture in the western Pacific. In contrast, the zonal advection of moisture is as weak as that in CMIP5. Moreover, improved shallow cumulus parameterization in CMIP6 can also lead to a better transition from local shallow moistening by shallow and congestus convection to deep convection. Hirota et al. (2018) showed that the shallow convection in MIROC6 can better transport PBL moisture upward to moisten the lower free troposphere at MJO initial stage, which is improved from MIROC5.

To the west of the MJO convective center, the CMIP6 shows slightly increased stratiform precipitation, drier at the lower troposphere and colder SST compared to CMIP5. This improvement is probably due to a better simulation of stratiform heating that can be attributed to an improvement of clouds/microphysics parameterization schemes in CMIP6. Wang et al. (2017) found that the presence of a stratiform heating behind the MJO convection plays an important role in strengthening lower-tropospheric poleward flows to the east of MJO convection. As the diabatic heating is the combined effects of radiative fluxes, phase changes of water, and

sensible heat flux from Earth's surface, this study has not evaluated which of the above-mentioned processes are responsible for the better representation of diabatic heating or stratiform heating at the rear in CMIP6 models, which deserves a further study in the future.

In addition, the enhanced east–west asymmetry of moisture tendency in CMIP6 is determined by the excessive lower-tropospheric drying process to the west of the MJO convection, which is mainly contributed by the negative vertical and zonal advections. The enhanced asymmetry may also explain the better simulation of propagation of the MJO in CMIP6.

In conclusion, the CMIP6 models show an overall improvement in simulating the MJO propagation compared to the CMIP5 models, which boosts our confidence when predicting MJO using CMIP6 models. The schematic diagram in Fig. 12 summarizes the quantified contribution of each process to the MJO eastward propagation skill in the CMIP5 and CMIP6 models. In the CMIP6 models, the vertical structures of diabatic heating (Q1), zonal circulation (consisting of zonal wind and vertical velocity), and equivalent potential temperature (EPT) are found to be the most important contributors to the MJO propagation skill, which are substantially improved in CMIP6 models. In addition, the linear regression relationships between these metrics and MJO eastward propagation skill tend to be more coherent and robust in CMIP6 compared to the CMIP5 models. These metrics are not isolated. The means of all PCC-derived metrics and their standard deviations (Fig. 13a) are found to be clustered in the area with high PCC and low standard deviation for CMIP6 models. The scatterplots of the linear regression of MJO propagation skill against the mean PCC of all metrics (Fig. 13b) showed that a good model from CMIP6 can generally simulate all metrics well.

Given better representation of moisture in the CMIP6 models, questions then arise what processes should be accredited to this improvement. As the CMIP6 models show a warmer SST and increased lower-tropospheric moisture in the winter mean state, the moisture transfer associated with air–sea interaction might be a potential reason, which warrants a further investigation.

Acknowledgments. This work is supported by National Natural Science Foundation of China (Grants 42088101 and 42030605) and National Key R&D Program of China (Grant 2020YFA0608004). We acknowledge the World Climate Research Programme's Working Group on Coupled Modeling, which is responsible for CMIP5 and CMIP6, and the climate modeling groups for producing and making available their model output. We thank the three anonymous reviewers for their valuable comments that helped to improve the manuscript.

Data availability statement. The observed daily outgoing longwave radiation data are available from National Oceanic and Atmospheric Administration (NOAA) at https://psl.noaa.gov/data/gridded/data.interp_OLR.html, as cited in

Liebmann and Smith (1996). The daily precipitation data are available from the Tropical Rainfall Measuring Mission at <https://gpm.nasa.gov/data>, as cited in Huffman et al. (2007). The sea surface temperature is available from the NOAA Optimum Interpolation Sea Surface Temperature at <https://www.ncei.noaa.gov/data/sea-surface-temperature-optimum-interpolation/v2/access/avhrr-only>, as cited in Reynolds et al. (2007). All ERA-Interim daily-averaged reanalysis data are available from the European Centre for Medium-Range Weather Forecasts (ECMWF) at <https://apps.ecmwf.int/datasets/data/interim-full-daily/levtype=sfc/>, as cited in Dee et al. (2011). The outputs from CMIP5 and CMIP6 models are available at <https://esgf-node.llnl.gov/>, as cited in Taylor et al. (2012) and Eyring et al. (2016) respectively.

REFERENCES

- Adames, Á. F., and J. M. Wallace, 2015: Three-dimensional structure and evolution of the moisture field in the MJO. *J. Atmos. Sci.*, **72**, 3733–3754, <https://doi.org/10.1175/JAS-D-15-0003.1>.
- , and D. Kim, 2016: The MJO as a dispersive, convectively coupled moisture wave: Theory and observations. *J. Atmos. Sci.*, **73**, 913–941, <https://doi.org/10.1175/JAS-D-15-0170.1>.
- Ahn, M.-S., D. Kim, K. R. Sperber, I.-S. Kang, E. Maloney, D. Waliser, and H. Hendon, 2017: MJO simulation in CMIP5 climate models: MJO skill metrics and process-oriented diagnosis. *Climate Dyn.*, **49**, 4023–4045, <https://doi.org/10.1007/s00382-017-3558-4>.
- , and Coauthors, 2020: MJO propagation across the Maritime Continent: Are CMIP6 models better than CMIP5 models? *Geophys. Res. Lett.*, **47**, e2020GL087250, <https://doi.org/10.1029/2020GL087250>.
- Alvarez, M. S., C. S. Vera, G. N. Kiladis, and B. Liebmann, 2016: Influence of the Madden Julian Oscillation on precipitation and surface air temperature in South America. *Climate Dyn.*, **46**, 245–262, <https://doi.org/10.1007/s00382-015-2581-6>.
- Arakawa, A., and W. H. Schubert, 1974: Interaction of a cumulus cloud ensemble with the large-scale environment. Part I. *J. Atmos. Sci.*, **31**, 674–701, [https://doi.org/10.1175/1520-0469\(1974\)031<0674:IOACCE>2.0.CO;2](https://doi.org/10.1175/1520-0469(1974)031<0674:IOACCE>2.0.CO;2).
- Arcodia, M. C., B. P. Kirtman, and L. S. P. Siqueira, 2020: How MJO teleconnections and ENSO interference impacts U.S. precipitation. *J. Climate*, **33**, 4621–4640, <https://doi.org/10.1175/JCLI-D-19-0448.1>.
- Arnold, N. P., M. Branson, Z. Kuang, D. A. Randall, and E. Tziperman, 2015: MJO intensification with warming in the superparameterized CESM. *J. Climate*, **28**, 2706–2724, <https://doi.org/10.1175/JCLI-D-14-00494.1>.
- Benedict, J. J., and D. Randall, 2007: Observed characteristics of the MJO relative to maximum rainfall. *J. Atmos. Sci.*, **64**, 2332–2354, <https://doi.org/10.1175/JAS3968.1>.
- Betts, A. K., 1986: A new convective adjustment scheme. Part I: Observational and theoretical basis. *Quart. J. Roy. Meteor. Soc.*, **112**, 677–691, <https://doi.org/10.1002/qj.49711247307>.
- Bretherton, C. S., J. R. McCaa, and H. Grenier, 2004: A new parameterization for shallow cumulus convection and its application to marine subtropical cloud-topped boundary layers. Part I: Description and 1D results. *Mon. Wea. Rev.*, **132**, 864–882, [https://doi.org/10.1175/1520-0493\(2004\)132<0864:ANPFS>2.0.CO;2](https://doi.org/10.1175/1520-0493(2004)132<0864:ANPFS>2.0.CO;2).

- Chen, G., and B. Wang, 2018: Effects of enhanced front Walker cell on the eastward propagation of the MJO. *J. Climate*, **31**, 7719–7738, <https://doi.org/10.1175/JCLI-D-17-0383.1>.
- , and —, 2020: Circulation factors determining the propagation speed of the Madden–Julian oscillation. *J. Climate*, **33**, 3367–3380, <https://doi.org/10.1175/JCLI-D-19-0661.1>.
- Chen, Y., and A. D. Del Genio, 2009: Evaluation of tropical cloud regimes in observations and a general circulation model. *Climate Dyn.*, **32**, 355–369, <https://doi.org/10.1007/s00382-008-0386-6>.
- Chikira, M., and M. Sugiyama, 2010: A cumulus parameterization with state-dependent entrainment rate. Part I: Description and sensitivity to temperature and humidity profiles. *J. Atmos. Sci.*, **67**, 2171–2193, <https://doi.org/10.1175/2010JAS3316.1>.
- Dee, D. P., and Coauthors, 2011: The ERA-Interim reanalysis: Configuration and performance of the data assimilation system. *Quart. J. Roy. Meteor. Soc.*, **137**, 553–597, <https://doi.org/10.1002/qj.828>.
- DeMott, C. A., N. P. Klingaman, and S. J. Woolnough, 2015: Atmosphere–ocean coupled processes in the Madden–Julian Oscillation. *Rev. Geophys.*, **53**, 1099–1154, <https://doi.org/10.1002/2014RG000478>.
- , J. J. Benedict, N. P. Klingaman, S. J. Woolnough, and D. A. Randall, 2016: Diagnosing ocean feedbacks to the MJO: SST-modulated surface fluxes and the moist static energy budget. *J. Geophys. Res. Atmos.*, **121**, 8350–8373, <https://doi.org/10.1002/2016JD025098>.
- Deng, L., and X. Wu, 2010: Effects of convective processes on GCM simulations of the Madden–Julian oscillation. *J. Climate*, **23**, 352–377, <https://doi.org/10.1175/2009JCLI3114.1>.
- Donner, L. J., 1993: A cumulus parameterization including mass fluxes, vertical momentum dynamics, and mesoscale effects. *J. Atmos. Sci.*, **50**, 889–906, [https://doi.org/10.1175/1520-0469\(1993\)050<0889:ACPIMF>2.0.CO;2](https://doi.org/10.1175/1520-0469(1993)050<0889:ACPIMF>2.0.CO;2).
- , C. J. Seman, R. S. Hemler, and S. Fan, 2001: A cumulus parameterization including mass fluxes, convective vertical velocities, and mesoscale effects: Thermodynamic and hydrological aspects in a general circulation model. *J. Climate*, **14**, 3444–3463, [https://doi.org/10.1175/1520-0442\(2001\)014<3444:ACPIMF>2.0.CO;2](https://doi.org/10.1175/1520-0442(2001)014<3444:ACPIMF>2.0.CO;2).
- Emanuel, K. A., 1987: An air–sea interaction model of intraseasonal oscillations in the tropics. *J. Atmos. Sci.*, **44**, 2324–2340, [https://doi.org/10.1175/1520-0469\(1987\)044<2324:AASIMO>2.0.CO;2](https://doi.org/10.1175/1520-0469(1987)044<2324:AASIMO>2.0.CO;2).
- , 1991: A scheme for representing cumulus convection in large-scale models. *J. Atmos. Sci.*, **48**, 2313–2335, [https://doi.org/10.1175/1520-0469\(1991\)048<2313:ASFRCC>2.0.CO;2](https://doi.org/10.1175/1520-0469(1991)048<2313:ASFRCC>2.0.CO;2).
- Emori, S., T. Nozawa, A. Numaguti, and I. Uno, 2001: Importance of cumulus parameterization for precipitation simulation over East Asia in June. *J. Meteor. Soc. Japan*, **79**, 939–947, <https://doi.org/10.2151/jmsj.79.939>.
- Eyring, V., S. Bony, G. A. Meehl, C. A. Senior, B. Stevens, R. J. Stouffer, and K. E. Taylor, 2016: Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization. *Geosci. Model Dev.*, **9**, 1937–1958, <https://doi.org/10.5194/gmd-9-1937-2016>.
- Fu, X., J.-Y. Lee, P.-C. Hsu, H. Taniguchi, B. Wang, W. Wang, and S. Weaver, 2013: Multi-model MJO forecasting during DYNAMO/CINDY period. *Climate Dyn.*, **41**, 1067–1081, <https://doi.org/10.1007/s00382-013-1859-9>.
- , W. Wang, T. Shinoda, H.-L. Ren, and X. Jia, 2017: Toward understanding the diverse impacts of air–sea interactions on MJO simulations. *J. Geophys. Res. Oceans*, **122**, 8855–8875, <https://doi.org/10.1002/2017JC013187>.
- Fuchs, Z., and D. J. Raymond, 2005: Large-scale modes in a rotating atmosphere with radiative-convective instability and WISHE. *J. Atmos. Sci.*, **62**, 4084–4094, <https://doi.org/10.1175/JAS3582.1>.
- , and —, 2007: A simple, vertically resolved model of tropical disturbances with a humidity closure. *Tellus*, **59A**, 344–354, <https://doi.org/10.1111/j.1600-0870.2007.00230.x>.
- , and —, 2017: A simple model of intraseasonal oscillations. *J. Adv. Model. Earth Syst.*, **9**, 1195–1211, <https://doi.org/10.1002/2017MS000963>.
- Garfinkel, C., J. J. Benedict, and E. D. Maloney, 2014: Impact of the MJO on the boreal winter extratropical circulation. *Geophys. Res. Lett.*, **41**, 6055–6062, <https://doi.org/10.1002/2014GL061094>.
- Gill, A. E., 1980: Some simple solutions for heat-induced tropical circulation. *Quart. J. Roy. Meteor. Soc.*, **106**, 447–462, <https://doi.org/10.1002/qj.49710644905>.
- Gonzalez, A. O., and X. Jiang, 2017: Winter mean lower tropospheric moisture over the Maritime Continent as a climate model diagnostic metric for the propagation of the Madden–Julian Oscillation. *Geophys. Res. Lett.*, **44**, 2588–2596, <https://doi.org/10.1002/2016GL072430>.
- Grandpeix, J., and J. Lafore, 2010: A density current parameterization coupled with Emanuel’s convection scheme. Part I: The models. *J. Atmos. Sci.*, **67**, 881–897, <https://doi.org/10.1175/2009JAS3044.1>.
- , —, and F. Cheruy, 2010: A density current parameterization coupled with Emanuel’s convection scheme. Part II: 1D simulations. *J. Atmos. Sci.*, **67**, 898–922, <https://doi.org/10.1175/2009JAS3045.1>.
- Gregory, D., 1995: The representation of moist convection in atmospheric models. *Proc. Seminar on Parametrization of Sub-grid Scale Physical Processes*, Reading, United Kingdom, ECMWF, 77–113, <https://www.ecmwf.int/sites/default/files/elibrary/1994/9641-representation-moist-convection-atmospheric-models.pdf>.
- , and P. R. Rowntree, 1990: A mass flux convection scheme with representation of cloud ensemble characteristics and stability-dependent closure. *Mon. Wea. Rev.*, **118**, 1483–1506, [https://doi.org/10.1175/1520-0493\(1990\)118<1483:AMFCSW>2.0.CO;2](https://doi.org/10.1175/1520-0493(1990)118<1483:AMFCSW>2.0.CO;2).
- Hendon, H. H., and J. Glick, 1997: Intraseasonal air–sea interaction in the tropical Indian and Pacific Oceans. *J. Climate*, **10**, 647–661, [https://doi.org/10.1175/1520-0442\(1997\)010<0647:IASIIT>2.0.CO;2](https://doi.org/10.1175/1520-0442(1997)010<0647:IASIIT>2.0.CO;2).
- Hewitt, H. T., D. Copsey, I. D. Culverwell, C. M. Harris, R. S. R. Hill, A. B. Keen, A. J. McLaren, and E. C. Hunke, 2011: Design and implementation of the infrastructure of HadGEM3: The next-generation Met Office climate modelling system. *Geosci. Model Dev.*, **4**, 223–253, <https://doi.org/10.5194/gmd-4-223-2011>.
- Hirota, N., T. Ogura, H. Tatebe, H. Shiogama, M. Kimoto, and H. Watanabe, 2018: Roles of shallow convective moistening in the eastward propagation of the MJO in MIROC6. *J. Climate*, **31**, 3033–3037, <https://doi.org/10.1175/JCLI-D-17-0246.1>.
- Hsu, P.-C., T. Li, and H. Murakami, 2014: Moisture asymmetry and MJO eastward propagation in an aquaplanet general circulation model. *J. Climate*, **27**, 8747–8760, <https://doi.org/10.1175/JCLI-D-14-00148.1>.
- Hu, F., T. Li, J. Gao, and L. Hao, 2020: Reexamining the moisture mode theories of the Madden–Julian oscillation based on

- observational analyses. *J. Climate*, **34**, 839–853, <https://doi.org/10.1175/JCLI-D-20-0441.1>.
- Huffman, G. J., and Coauthors, 2007: The TRMM Multisatellite Precipitation Analysis (TMPA): Quasi-global, multiyear, combined-sensor precipitation estimates at fine scales. *J. Hydrometeorol.*, **8**, 38–55, <https://doi.org/10.1175/JHM560.1>.
- Hung, M.-P., J.-L. Lin, W. Wang, D. Kim, T. Shinoda, and S. J. Weaver, 2013: MJO and convectively coupled equatorial waves simulated by CMIP5 climate models. *J. Climate*, **26**, 6185–6214, <https://doi.org/10.1175/JCLI-D-12-00541.1>.
- Jiang, X., 2017: Key processes for the eastward propagation of the Madden–Julian Oscillation based on multimodel simulations. *J. Geophys. Res. Atmos.*, **122**, 755–770, <https://doi.org/10.1002/2016JD025955>.
- , and Coauthors, 2015: Vertical structure and physical processes of the Madden–Julian Oscillation: Exploring key model physics in climate simulations. *J. Geophys. Res. Atmos.*, **120**, 4718–4748, <https://doi.org/10.1002/2014JD022375>.
- , and Coauthors, 2020a: Fifty years of research on the Madden–Julian Oscillation: Recent progress, challenges, and perspectives. *J. Geophys. Res. Atmos.*, **125**, e2019JD030911, <https://doi.org/10.1029/2019JD030911>.
- , E. Maloney, and H. Su, 2020b: Large-scale controls of propagation of the Madden–Julian Oscillation. *npj Climate Atmos. Sci.*, **3**, 29, <https://doi.org/10.1038/s41612-020-00134-x>.
- Kain, J. S., and J. M. Fritsch, 1990: A one-dimensional entraining/detraining plume model and its application in convective parameterization. *J. Atmos. Sci.*, **47**, 2784–2802, [https://doi.org/10.1175/1520-0469\(1990\)047<2784:AODEPM>2.0.CO;2](https://doi.org/10.1175/1520-0469(1990)047<2784:AODEPM>2.0.CO;2).
- , and —, 1993: Convective parameterization for mesoscale models: The Kain–Fritsch scheme. *The Representation of Cumulus Convection in Numerical Models*, Meteor. Monogr., No. 46, Amer. Meteor. Soc., 165–170.
- Kiladis, G. N., K. H. Straub, and P. T. Haertel, 2005: Zonal and vertical structure of the Madden–Julian oscillation. *J. Atmos. Sci.*, **62**, 2790–2809, <https://doi.org/10.1175/JAS3520.1>.
- , M. C. Wheeler, P. T. Haertel, K. H. Straub, and P. E. Roundy, 2009: Convectively coupled equatorial waves. *Rev. Geophys.*, **47**, RG2003, <https://doi.org/10.1029/2008RG000266>.
- Kim, D., and I.-S. Kang, 2012: A bulk mass flux convection scheme for climate model: Description and moisture sensitivity. *Climate Dyn.*, **38**, 411–429, <https://doi.org/10.1007/s00382-010-0972-2>.
- , and Coauthors, 2009: Application of MJO simulation diagnostics to climate models. *J. Climate*, **22**, 6413–6436, <https://doi.org/10.1175/2009JCLI3063.1>.
- , A. H. Sobel, E. D. Maloney, D. M. W. Frierson, and I.-S. Kang, 2011: A systematic relationship between intraseasonal variability and mean state bias in AGCM simulations. *J. Climate*, **24**, 5506–5520, <https://doi.org/10.1175/2011JCLI4177.1>.
- , and Coauthors, 2014: Process-oriented MJO simulation diagnostic: Moisture sensitivity of simulated convection. *J. Climate*, **27**, 5379–5395, <https://doi.org/10.1175/JCLI-D-13-00497.1>.
- Kim, S., J.-S. Kug, and K.-H. Seo, 2020: Impacts of MJO on the intraseasonal temperature variation in East Asia. *J. Climate*, **33**, 8903–8916, <https://doi.org/10.1175/JCLI-D-20-0302.1>.
- Klotzbach, P. J., 2010: On the Madden–Julian oscillation–Atlantic hurricane relationship. *J. Climate*, **23**, 282–293, <https://doi.org/10.1175/2009JCLI2978.1>.
- , 2014: The Madden–Julian oscillation’s impacts on worldwide tropical cyclone activity. *J. Climate*, **27**, 2317–2330, <https://doi.org/10.1175/JCLI-D-13-00483.1>.
- Lau, K.-M., and L. Peng, 1987: Origin of low-frequency (intraseasonal) oscillations in the tropical atmosphere. Part I: Basic theory. *J. Atmos. Sci.*, **44**, 950–972, [https://doi.org/10.1175/1520-0469\(1987\)044<0950:OOLFOI>2.0.CO;2](https://doi.org/10.1175/1520-0469(1987)044<0950:OOLFOI>2.0.CO;2).
- Li, C., X. Jia, J. Ling, W. Zhou, and C. Zhang, 2009: Sensitivity of MJO simulations to diabatic heating profiles. *Climate Dyn.*, **32**, 167–187, <https://doi.org/10.1007/s00382-008-0455-x>.
- Liebmann, B., and C. A. Smith, 1996: Description of a complete (interpolated) outgoing longwave radiation dataset. *Bull. Amer. Meteor. Soc.*, **77**, 1275–1277, <https://doi.org/10.1175/1520-0477-77.6.1274>.
- Lin, J.-L., and Coauthors, 2006: Tropical intraseasonal variability in 14 IPCC AR4 climate models. Part I: Convective signals. *J. Climate*, **19**, 2665–2690, <https://doi.org/10.1175/JCLI3735.1>.
- , D. Kim, M. Lee, and I.-S. Kang, 2007: Effects of cloud-radiative heating on atmospheric general circulation model (AGCM) simulations of convectively coupled equatorial waves. *J. Geophys. Res.*, **112**, D24107, <https://doi.org/10.1029/2006JD008291>.
- , M.-I. Lee, D. Kim, I.-S. Kang, and D. M. W. Frierson, 2008: The impacts of convective parameterization and moisture triggering on AGCM-simulated convectively coupled equatorial waves. *J. Climate*, **21**, 883–909, <https://doi.org/10.1175/2007JCLI1790.1>.
- Ling, J., C. Zhang, S. Wang, and C. Li, 2017: A new interpretation of the ability of global models to simulate the MJO. *Geophys. Res. Lett.*, **44**, 5798–5806, <https://doi.org/10.1002/2017GL073891>.
- , Y. Zhao, and G. Chen, 2019: Barrier effect on MJO propagation by the Maritime Continent in the MJO task force/GEWEX atmospheric system study models. *J. Climate*, **32**, 5529–5547, <https://doi.org/10.1175/JCLI-D-18-0870.1>.
- Madden, R. A., and P. R. Julian, 1971: Detection of a 40–50 day oscillation in the zonal wind in the tropical Pacific. *J. Atmos. Sci.*, **28**, 702–708, [https://doi.org/10.1175/1520-0469\(1971\)028<0702:DOADOI>2.0.CO;2](https://doi.org/10.1175/1520-0469(1971)028<0702:DOADOI>2.0.CO;2).
- , and —, 1972: Description of global-scale circulation cells in the tropics with a 40–50 day period. *J. Atmos. Sci.*, **29**, 1109–1123, [https://doi.org/10.1175/1520-0469\(1972\)029<1109:DOGSCC>2.0.CO;2](https://doi.org/10.1175/1520-0469(1972)029<1109:DOGSCC>2.0.CO;2).
- Majda, A. J., and S. N. Stechmann, 2009: The skeleton of tropical intraseasonal oscillations. *Proc. Natl. Acad. Sci. USA*, **106**, 8417–8422, <https://doi.org/10.1073/pnas.0903367106>.
- Maloney, E. D., and D. L. Hartmann, 1998: Frictional moisture convergence in a composite life cycle of the Madden–Julian oscillation. *J. Climate*, **11**, 2387–2403, [https://doi.org/10.1175/1520-0442\(1998\)011<2387:FMCIAC>2.0.CO;2](https://doi.org/10.1175/1520-0442(1998)011<2387:FMCIAC>2.0.CO;2).
- Matsuno, T., 1966: Quasi-geostrophic motions in the equatorial area. *J. Meteor. Soc. Japan*, **44**, 25–43, https://doi.org/10.2151/jmsj1965.44.1_25.
- Moorthi, S., and M. J. Suarez, 1992: Relaxed Arakawa–Schubert: A parameterization of moist convection for general circulation models. *Mon. Wea. Rev.*, **120**, 978–1002, [https://doi.org/10.1175/1520-0493\(1992\)120<0978:RASAP0>2.0.CO;2](https://doi.org/10.1175/1520-0493(1992)120<0978:RASAP0>2.0.CO;2).
- Neale, R. B., J. H. Richter, and M. Jochum, 2008: The impact of convection on ENSO: From a delayed oscillator to a series of events. *J. Climate*, **21**, 5904–5924, <https://doi.org/10.1175/2008JCLI2244.1>.
- Neelin, J. D., I. M. Held, and K. H. Cook, 1987: Evaporation–wind feedback and low-frequency variability in the tropical atmosphere. *J. Atmos. Sci.*, **44**, 2341–2348, [https://doi.org/10.1175/1520-0469\(1987\)044<2341:EWFALF>2.0.CO;2](https://doi.org/10.1175/1520-0469(1987)044<2341:EWFALF>2.0.CO;2).
- Nordeng, T. E., 1994: Extended versions of the convective parameterization scheme at ECMWF and their impact on the mean

- and transient activity of the model in the tropics. ECMWF Tech. Memo. 206, 41 pp., <https://doi.org/10.21957/e34xwhysw>.
- Pan, D. M., and D. A. Randall, 1998: A cumulus parameterization with a prognostic closure. *Quart. J. Roy. Meteor. Soc.*, **124**, 949–981, <https://doi.org/10.1002/qj.49712454714>.
- Park, S., and C. S. Bretherton, 2009: The University of Washington shallow convection and moist turbulence schemes and their impact on climate simulations with the Community Atmosphere Model. *J. Climate*, **22**, 3449–3469, <https://doi.org/10.1175/2008JCLI2557.1>.
- Ray, P., C. Zhang, M. W. Moncrieff, J. Dudhia, J. M. Caron, L. R. Leung, and C. Bruyere, 2011: Role of the atmospheric mean state on the initiation of the Madden–Julian Oscillation in a tropical channel model. *Climate Dyn.*, **36**, 161–184, <https://doi.org/10.1007/s00382-010-0859-2>.
- Raymond, D. J., and Ž. Fuchs, 2009: Moisture modes and the Madden–Julian oscillation. *J. Climate*, **22**, 3031–3046, <https://doi.org/10.1175/2008JCLI2739.1>.
- Reynolds, R. W., T. M. Smith, C. Liu, D. B. Chelton, K. S. Casey, and M. G. Schlax, 2007: Daily high-resolution-blended analyses for sea surface temperature. *J. Climate*, **20**, 5473–5496, <https://doi.org/10.1175/2007JCLI1824.1>.
- Richter, J. H., and P. J. Rasch, 2008: Effects of convective momentum transport on the atmospheric circulation in the Community Atmosphere Model, version 3. *J. Climate*, **21**, 1487–1499, <https://doi.org/10.1175/2007JCLI1789.1>.
- Seo, K.-H., and W. Wang, 2010: The Madden–Julian oscillation simulated in the NCEP Climate Forecast System Model: The importance of stratiform heating. *J. Climate*, **23**, 4770–4793, <https://doi.org/10.1175/2010JCLI2983.1>.
- , and S.-W. Son, 2012: The global atmospheric circulation response to tropical diabatic heating associated with the Madden–Julian oscillation during northern winter. *J. Atmos. Sci.*, **69**, 79–96, <https://doi.org/10.1175/2011JAS3686.1>.
- Sobel, A. H., and E. D. Maloney, 2012: An idealized semi-empirical framework for modeling the Madden–Julian oscillation. *J. Atmos. Sci.*, **69**, 1691–1705, <https://doi.org/10.1175/JAS-D-11-0118.1>.
- , and —, 2013: Moisture modes and the eastward propagation of the MJO. *J. Atmos. Sci.*, **70**, 187–192, <https://doi.org/10.1175/JAS-D-12-0189.1>.
- Stevens, B., and Coauthors, 2013: Atmospheric component of the MPI-M Earth System Model: ECHAM6. *J. Adv. Model. Earth Syst.*, **5**, 146–172, <https://doi.org/10.1002/jame.20015>.
- Subramanian, A. C., M. Jochum, A. J. Miller, R. Murtugudde, R. B. Neale, and D. E. Waliser, 2011: The Madden–Julian oscillation in CCSM4. *J. Climate*, **24**, 6261–6282, <https://doi.org/10.1175/JCLI-D-11-00031.1>.
- Taylor, K. E., R. J. Stouffer, and G. A. Meehl, 2012: An overview of CMIP5 and the experiment design. *Bull. Amer. Meteor. Soc.*, **93**, 485–498, <https://doi.org/10.1175/BAMS-D-11-00094.1>.
- Thayer-Calder, K., and D. A. Randall, 2009: The role of convective moistening in the Madden–Julian oscillation. *J. Atmos. Sci.*, **66**, 3297–3312, <https://doi.org/10.1175/2009JAS3081.1>.
- Tian, B. J., D. E. Waliser, E. J. Fetzer, and Y. L. Yung, 2010: Vertical moist thermodynamic structure of the Madden–Julian oscillation in atmospheric infrared sounder retrievals: An update and a comparison to ECMWF Interim Re-Analysis. *Mon. Wea. Rev.*, **138**, 4576–4582, <https://doi.org/10.1175/2010MWR3486.1>.
- Tiedtke, M., 1989: A comprehensive mass flux scheme for cumulus parameterization in large-scale models. *Mon. Wea. Rev.*, **117**, 1779–1800, [https://doi.org/10.1175/1520-0493\(1989\)117<1779:ACMFSF>2.0.CO;2](https://doi.org/10.1175/1520-0493(1989)117<1779:ACMFSF>2.0.CO;2).
- Vitart, F., and Coauthors, 2017: The Subseasonal to Seasonal Prediction (S2S) Project database. *Bull. Amer. Meteor. Soc.*, **98**, 163–173, <https://doi.org/10.1175/BAMS-D-16-0017.1>.
- Waliser, D. E., 2011: Predictability and forecasting. *Intraseasonal Variability of the Atmosphere–Ocean Climate System*, 2nd ed. W. K. M. Lau and D. E. Waliser, Eds., Springer, 433–468.
- , and Coauthors, 2009: MJO simulation diagnostics. *J. Climate*, **22**, 3006–3030, <https://doi.org/10.1175/2008JCLI2731.1>.
- Wang, B., 1988: Dynamics of tropical low-frequency waves: An analysis of the moist Kelvin wave. *J. Atmos. Sci.*, **45**, 2051–2065, [https://doi.org/10.1175/1520-0469\(1988\)045<2051:DOTLFW>2.0.CO;2](https://doi.org/10.1175/1520-0469(1988)045<2051:DOTLFW>2.0.CO;2).
- , and H. Rui, 1990: Dynamics of the coupled moist Kelvin–Rossby wave on an equatorial β -plane. *J. Atmos. Sci.*, **47**, 397–413, [https://doi.org/10.1175/1520-0469\(1990\)047<0397:DOTCMK>2.0.CO;2](https://doi.org/10.1175/1520-0469(1990)047<0397:DOTCMK>2.0.CO;2).
- , and G. Chen, 2017: A general theoretical framework for understanding essential dynamics of Madden–Julian Oscillation. *Climate Dyn.*, **49**, 2309–2328, <https://doi.org/10.1007/s00382-016-3448-1>.
- , and S.-S. Lee, 2017: MJO propagation shaped by zonal asymmetric structures: Results from 24 GCM simulations. *J. Climate*, **30**, 7933–7952, <https://doi.org/10.1175/JCLI-D-16-0873.1>.
- , F. Liu, and G. Chen, 2016: A trio-interaction theory for Madden–Julian Oscillation. *Geosci. Lett.*, **3**, 34, <https://doi.org/10.1186/s40562-016-0066-z>.
- , and Coauthors, 2018: Dynamics-oriented diagnostics for the Madden–Julian oscillation. *J. Climate*, **31**, 3117–3135, <https://doi.org/10.1175/JCLI-D-17-0332.1>.
- Wang, L., T. Li, E. Maloney, and B. Wang, 2017: Fundamental causes of propagating and nonpropagating MJOs in MJOTF/GASS models. *J. Climate*, **30**, 3743–3769, <https://doi.org/10.1175/JCLI-D-16-0765.1>.
- Wheeler, M. C., H. H. Harry, S. Cleland, H. Meinke, and A. Donald, 2009: Impacts of the Madden–Julian oscillation on Australian rainfall and circulation. *J. Climate*, **22**, 1482–1498, <https://doi.org/10.1175/2008JCLI2595.1>.
- Wilcox, E. M., and J. J. Donner, 2007: The frequency of extreme rain events in satellite rain-rate estimates and an atmospheric general circulation model. *J. Climate*, **20**, 53–69, <https://doi.org/10.1175/JCLI3987.1>.
- Wu, T., 2012: A mass-flux cumulus parameterization scheme for large-scale models: Description and test with observations. *Climate Dyn.*, **38**, 725–744, <https://doi.org/10.1007/s00382-011-0995-3>.
- Wu, X. Q., L. P. Deng, X. L. Song, and G. J. Zhang, 2007: Coupling of convective momentum transport with convective heating in global climate simulations. *J. Atmos. Sci.*, **64**, 1334–1349, <https://doi.org/10.1175/JAS3894.1>.
- Yanai, M., S. Esbensen, and J.-H. Chu, 1973: Determination of bulk properties of tropical cloud clusters from large-scale heat and moisture budgets. *J. Atmos. Sci.*, **30**, 611–627, [https://doi.org/10.1175/1520-0469\(1973\)030<0611:DOBPOT>2.0.CO;2](https://doi.org/10.1175/1520-0469(1973)030<0611:DOBPOT>2.0.CO;2).
- Yang, D., and A. P. Ingersoll, 2011: Testing the hypothesis that the MJO is a mixed Rossby–gravity wave packet. *J. Atmos. Sci.*, **68**, 226–239, <https://doi.org/10.1175/2010JAS3563.1>.
- , and —, 2013: Triggered convection, gravity waves, and the MJO: A shallow-water model. *J. Atmos. Sci.*, **70**, 2476–2486, <https://doi.org/10.1175/JAS-D-12-0255.1>.

- Yoshimura, H., R. Mizuta, and H. Murakami, 2015: A spectral cumulus parameterization scheme interpolating between two convective updrafts with semi-Lagrangian calculation of transport by compensatory subsidence. *Mon. Wea. Rev.*, **143**, 597–621, <https://doi.org/10.1175/MWR-D-14-00068.1>.
- Zhang, C., 2005: Madden–Julian Oscillation. *Rev. Geophys.*, **43**, RG2003, <https://doi.org/10.1029/2004RG000158>.
- , A. F. Adames, B. Khouider, B. Wang, and D. Yang, 2020: Four theories of the Madden–Julian Oscillation. *Rev. Geophys.*, **58**, e2019RG000685, <https://doi.org/10.1029/2019RG000685>.
- Zhang, G. J., and N. A. McFarlane, 1995: Sensitivity of climate simulations to the parameterization of cumulus convection in the Canadian Climate Center general circulation model. *Atmos.–Ocean*, **33**, 407–446, <https://doi.org/10.1080/07055900.1995.9649539>.
- , and M. Mu, 2005: Effects of modifications to the Zhang–McFarlane convection parameterization on the simulation of the tropical precipitation in the National Center for Atmospheric Research Community Climate Model, version 3. *J. Geophys. Res.*, **110**, D09109, <https://doi.org/10.1029/2004JD005617>.
- , and X. Song, 2009: Interaction of deep and shallow convection is key to Madden–Julian Oscillation simulation. *Geophys. Res. Lett.*, **36**, L09708, <https://doi.org/10.1029/2009GL037340>.
- Zhou, L., R. B. Neale, M. Jochum, and R. Murtugudde, 2012: Improved Madden–Julian oscillations with improved physics: The impact of modified convection parameterizations. *J. Climate*, **25**, 1116–1136, <https://doi.org/10.1175/2011JCLI4059.1>.
- Zhou, S., M. L’Heureux, S. Weaver, and A. Kumar, 2012: A composite study of the MJO influence on the surface air temperature and precipitation over the continental United States. *Climate Dyn.*, **38**, 1459–1471, <https://doi.org/10.1007/s00382-011-1001-9>.
- Zhou, W., D. Yang, S.-P. Xie, and J. Ma, 2020: Amplified Madden–Julian oscillation impacts in the Pacific–North America region. *Nat. Climate Change*, **10**, 654–660, <https://doi.org/10.1038/s41558-020-0814-0>.
- Zhu, J., W. Wang, and A. Kumar, 2017: Simulations of MJO propagation across the Maritime Continent: Impacts of SST feedback. *J. Climate*, **30**, 1689–1704, <https://doi.org/10.1175/JCLI-D-16-0367.1>.