

# Trends and Periodicities of Rainfall Over India

P. JAGANNATHAN and B. PARTHASARATHY—*Indian Institute of Tropical Meteorology, Poona 5, India*

**ABSTRACT**—Trends and periodicities in the annual rainfall of India were studied using data for 48 stations having record lengths of over 70 yr. Increasing or decreasing rainfall tendencies were found over large continuous areas in India. These trends, however, are not significant over all the stations in the areas but only at a few places distributed at random. Wherever a trend is significant, it has persistence or a periodicity of more than 40 yr. Quasi-

biennial oscillation (QBO) is exhibited at several stations in the areas of increasing or decreasing trend. Similarly, the 11-yr cycle (solar cycle) is also exhibited in both areas. The QBO and the solar cycle are both present at only three stations, however. This casts a serious doubt on whether or not there is any physical relation between the QBO and the solar cycle, at least for rainfall.

## 1. INTRODUCTION

Much work has been done in various countries on trends and fluctuations of rainfall and temperature, notably by Reynolds (1953), Kraus (1960), Willett (1950), and Callendar (1961). Angell et al. (1966) observed quasi-biennial oscillations (QBO) in the tropospheric zonal winds of the Tropics and subtropics, and Landsberg et al. (1963) found evidence of QBO in the surface temperature in many parts of the world. Pramanik and Jagannathan (1953) found systematic variations in the annual rainfall over certain parts of India. Koteswaram and Alvi (1969) and Bhargava and Bansal (1969) reported a QBO in the southwest monsoon/annual rainfall over some stations on the west coast of south India. Jagannathan and Bhalme (1973) showed that rainfall in India during the monsoon season (June to September) has oscillations corresponding to the sunspot cycle. Further, they have found that the QBO is significantly present in some of the parameters. Their study, however, was restricted to the monsoon season alone and that too was based on pentad rainfall. Here, we propose to examine the oscillations, if any, in the annual rainfall and to study their nature. For this purpose a network of 48 stations having long-period homogeneous rainfall data has been selected (table 1). The annual rainfall,  $A_0$ , is taken for computational convenience as equal to  $\Sigma R_i/12$ , where  $R_i$  is the monthly rainfall.

## 2. PERSISTENCE

The various alternatives to randomness have the common property of low-frequency variation, which introduces positive serial correlation at small lags. To calculate the lag-one serial correlation, we used the following formula:

$$r_1 = \frac{(N-1) \sum_{i=1}^{N-1} x_i x_{i+1} - \left( \sum_{i=1}^{N-1} x_i \right) \left( \sum_{i=2}^N x_i \right)}{\sqrt{\left[ (N-1) \sum_{i=1}^{N-1} x_i^2 - \left( \sum_{i=1}^{N-1} x_i \right)^2 \right] \left[ (N-1) \sum_{i=2}^N x_i^2 - \left( \sum_{i=2}^N x_i \right)^2 \right]}} \quad (1)$$

The significance of the lag-one correlation,  $r_1$ , is tested using the one-tail 95-percent significance point of the Gaussian distribution (WMO 1966). The test value  $(r_1)_t$  is computed from

$$(r_1)_t = \frac{-1 + t_g \sqrt{N-2}}{N-1} \quad (2)$$

where  $t_g$  is the value of the standard deviate in the Gaussian distribution corresponding to the desired level of significance. The  $r_1$  values for all the stations are given in table 2.

Gilman et al. (1963) have given the method of finding the persistence of the first-order, linear Markov process, which is a dominant form of trend. Accordingly, the serial correlations at lag two and lag three were compared with  $r_2^3$  and  $r_1^3$ , respectively. When  $r_1$  was negative, it was tested against the two-tailed value and interpreted as indicative of marked high-frequency oscillations. The significant values are indicated in table 2. They show that the sample values of  $r_1$  are positive and significantly greater than the test value (at 95 percent) at Trivandrum, Fort Cochin, Akola, Veraval, Sagar, and Shillong with the values of  $r_2$  and  $r_3$  equal to or greater than  $r_2^3$  and  $r_1^3$ , respectively. This indicates Markov linear-type persistence at these stations. The  $r_1$  is also significant at Daltonganj, Allahabad, and Darjeeling, but the values of  $r_2$  and  $r_3$  are less than  $r_2^3$  and  $r_1^3$ , respectively, indicating that the persistence is not linear. At Bangalore

and Darbhanga,  $r_1$  is significantly negative, indicating the presence of high-frequency oscillations.

### 3. TREND

The Mann-Kendall rank statistic has been suggested as a powerful test (Kendall and Stuart 1961) when the most likely alternative to randomness is linear or non-linear trend. The statistic  $\tau$  is computed from

$$\tau = \frac{4\sum n_i}{N(N-1)} - 1 \quad (3)$$

where  $n_i$  is the number of values larger than the  $i$ th value in the series subsequent to its position in the time series. The test statistic ( $\tau$ ), is

$$(\tau)_i = \pm t_g \sqrt{\frac{4N+10}{9N(N-1)}} \quad (4)$$

where  $t_g$  is the value of  $t$  at the probability point in the Gaussian distribution appropriate to the two-tailed test. Table 3 gives the Mann-Kendall rank statistic values significant at 95 and 99 percent. A positive value indicates that the trend is one of increasing tendency while a negative value indicates a decreasing tendency. This test has been applied to all the series irrespective of whether or not the first test indicated a trend. We found a significant increasing trend at Trivandrum, Fort Cochin, Bellary, Belgaum, Masulipatnam, Bombay, Veraval, Indore, and Leh, and significant decreasing trend at Darjeeling and Simla. The rest of the stations do not exhibit significant trends.

### 4. LOW-PASS FILTER

To understand the nature of this trend, we subjected the series to a "low-pass filter" (WMO 1966), thus suppressing the high-frequency oscillations. The weights used were the nine ordinates of the Gaussian probability curve (0.01, 0.05, 0.12, 0.20, 0.24, 0.20, 0.12, 0.05, and 0.01). The response curve of the Gaussian low-pass filter has a response function that is equal to unity at infinite wavelengths; it then tails off asymptotically to zero with decreasing wavelength. The response is approximately

$$R(f) = \exp[-2\pi^2\sigma_g^2 f^2] \quad (5)$$

where  $\sigma_g$  is the appropriate standard deviation (i.e.,  $6\sigma_g = 10$  yr). The trend is not linear but oscillatory, consisting of periods of more than 10 yr in duration. A few filtered series along with the unfiltered series are shown as examples in figure 1.

### 5. POWER SPECTRUM ANALYSIS

Having found these oscillations in the annual rainfall, we now wish to see if there are any cycles. To do this, we conducted a power spectrum analysis of the time series of the mean annual rainfall for all stations. Tukey's (1950)

TABLE 1.—Rainfall data used in this study

| Station symbol | Name of station | Period of data | Station symbol | Name of station | Period of data |
|----------------|-----------------|----------------|----------------|-----------------|----------------|
| TRV*           | Trivandrum      | 1890-1967      | AHM            | Ahmadabad       | 1869-1967      |
| PBN            | Pamban          | 1893-1967      | SGR            | Sagar           | 1871-1967      |
| F.CHN          | Fort Cochin     | 1864-1967      | DTG            | Daltanganj      | 1893-1967      |
| KDK            | Kodaikanal      | 1901-1967      | DMK            | Dumka           | 1883-1967      |
| MNG            | Mangalore       | 1864-1967      | SLC            | Silchar         | 1869-1967      |
| BNG            | Bangalore       | 1837-1967      | KTA            | Kotah           | 1869-1967      |
| MDS            | Madras          | 1813-1967      | ALB            | Allahabad       | 1845-1967      |
| BLY            | Bellary         | 1869-1967      | SHL            | Shillong        | 1867-1967      |
| BLG            | Belgaum         | 1866-1967      | PTN            | Patna           | 1868-1967      |
| MPT            | Masulipatnam    | 1863-1967      | DBN            | Darbhanga       | 1875-1967      |
| HYD            | Hyderabad       | 1893-1967      | GHT            | Gauhati         | 1849-1967      |
| VSK            | Visakhapatnam   | 1866-1967      | JDP            | Jodhpur         | 1897-1967      |
| PNA            | Poona           | 1856-1967      | JSM            | Jaisalmer       | 1883-1967      |
| BMB            | Bombay          | 1847-1967      | JPR            | Jaipur          | 1875-1967      |
| JGD            | Jagdapur        | 1910-1967      | DJG            | Darjeeling      | 1868-1967      |
| AKL            | Akola           | 1870-1967      | AGR            | Agra            | 1862-1967      |
| CTK            | Cuttack         | 1867-1967      | DBH            | Dibrugarh       | 1902-1967      |
| VVL            | Veraval         | 1893-1967      | BKR            | Bikaner         | 1878-1967      |
| NGP            | Nagpur          | 1855-1967      | DLH            | New Delhi       | 1875-1967      |
| SBP            | Sambalpur       | 1867-1967      | MKS            | Mukteswar       | 1898-1967      |
| SGR            | Sagar Island    | 1869-1967      | LDN            | Ludhiana        | 1868-1967      |
| DWK            | Dwarka          | 1901-1967      | SML            | Simla           | 1862-1967      |
| CAL            | Calcutta        | 1829-1967      | SRN            | Srinagar        | 1893-1967      |
| IND            | Indore          | 1877-1967      | LEH            | Leh             | 1876-1967      |

\*Stations as listed run from southern India to northern India in figure 2.

TABLE 2.—Lag-one serial correlation,  $r_1$

| Name of station | $r_1$   | Name of station | $r_1$  |
|-----------------|---------|-----------------|--------|
| Trivandrum      | +0.208* | Ahmadabad       | +0.092 |
| Pamban          | -.122   | Sagar           | +.264* |
| Fort Cochin     | +.170*  | Daltanganj      | +.188† |
| Kodaikanal      | -.087   | Dumka           | -.072  |
| Mangalore       | -.018   | Silchar         | +.029  |
| Bangalore       | -.231†  | Kotah           | +.125  |
| Madras          | +.084   | Allahabad       | +.303† |
| Bellary         | +.054   | Patna           | +.101  |
| Belgaum         | +.110   | Shillong        | +.209* |
| Masulipatnam    | +.146   | Darbhanga       | -.173† |
| Hyderabad       | +.068   | Gauhati         | -.003  |
| Visakhapatnam   | -.018   | Jodhpur         | +.119  |
| Poona           | +.103   | Jaisalmer       | -.005  |
| Bombay          | +.008   | Jaipur          | -.007  |
| Jagdapur        | +.098   | Agra            | -.047  |
| Akola           | +.177*  | Darjeeling      | +.255† |
| Cuttack         | -.016   | Dibrugarh       | +.062  |
| Veraval         | +.232*  | Bikaner         | +.029  |
| Nagpur          | +.098   | New Delhi       | +.114  |
| Sambalpur       | +.100   | Mukteswar       | -.072  |
| Sagar Island    | +.079   | Ludhiana        | -.114  |
| Dwarka          | -.028   | Simla           | +.144  |
| Calcutta        | -.004   | Srinagar        | -.083  |
| Indore          | +.101   | Leh             | +.109  |

\*Significant Markov linear persistence at 95 percent.

†Significant correlation coefficient at 95 percent.

procedure for computing power spectra has been followed. To achieve satisfactory resolution in the spectrum, we chose a maximum lag,  $m$ , as large<sup>est</sup> as possible but not exceeding one-third of the total number of years of the record involved in the analysis. To reduce the chances of picking up a high power, we conducted the analysis sepa-

TABLE 3.—Mann-Kendall rank statistic,  $\tau$

| Name of station | $\tau$  | Name of station | $\tau$ |
|-----------------|---------|-----------------|--------|
| Trivandrum      | +0.211* | Ahmadabad       | +0.037 |
| Pamban          | -.034   | Sagar           | +.080  |
| Fort Cochin     | +.185*  | Daltonganj      | +.051  |
| Kodaikanal      | -.020   | Dumka           | -.001  |
| Mangalore       | +.078   | Silchar         | +.081  |
| Bangalore       | +.023   | Kotah           | +.079  |
| Madras          | +.033   | Allahabad       | +.027  |
| Bellary         | +.181*  | Patna           | -.050  |
| Belgaum         | +.232*  | Shillong        | +.071  |
| Masulipatnam    | +.143†  | Darbhanga       | -.014  |
| Hyderabad       | +.044   | Gauhati         | -.022  |
| Visakhapatnam   | -.104   | Jodhpur         | +.145  |
| Poona           | +.040   | Jaisalmer       | -.048  |
| Bombay          | +.149†  | Jaipur          | +.012  |
| Jagdalpur       | -.102   | Agra            | +.127  |
| Akola           | +.074   | Darjeeling      | -.197* |
| Cuttack         | -.002   | Dibrugrah       | -.117  |
| Veraval         | +.304*  | Bikaner         | -.010  |
| Nagpur          | +.056   | New Delhi       | +.077  |
| Sambalpur       | +.069   | Mukteswar       | +.001  |
| Sagar Island    | +.057   | Ludhiana        | +.067  |
| Dwarka          | +.113   | Simla           | -.225* |
| Calcutta        | -.088   | Srinagar        | +.020  |
| Indore          | +.159†  | Leh             | +.204* |

\*Significant value at 99-percent level  
 †Significant value at 95-percent level

rately with five or six different maximum lags that might have arisen due to "aliasing effect" consistent with the above restrictions. The null hypothesis for this purpose was considered in accordance with whether or not the series revealed any persistence. If the persistence was of the Markov linear-type, the appropriate red noise spectrum and the associated 99-, 95-, and 90-percent limits were calculated, and the individual peaks were tested with reference to these limits. If the lag-one correlation was significantly greater in magnitude than zero but higher lag correlation did not taper off exponentially, the spectral estimates in the first half were tested with reference to the red noise spectrum and the rest against white noise. In the absence of any persistence, the spectral estimates were tested against the white noise spectrum.

Significant periods (at 99-, 95-, 90-percent levels) revealed by the spectral analysis are given in table 4. This table shows low-frequency oscillations (more than 40 yr) at Belgaum, Bombay, Akola, Veraval, Nagpur, Indore, Sagar, Dumka, Allahabad, Agra, Simla, and Leh. Cycles of nearly 11 yr are seen at Madras, Jagdalpur, Silchar, Kotah, Allahabad, Jodhpur, Darjeeling, and New Delhi. Figure 2 shows these stations and also those that have exhibited trends.

A quasi-biennial oscillation is observed at Bangalore, Madras, Bombay, Jagdalpur, Dumka, Silchar, Darbhanga, Jaipur, Agra, Bikaner, Ludhiana, and Srinagar. It is important to note that of all the stations having nearly 11-yr cycles, only Madras, Jagdalpur, and Silchar have ex-




FIGURE 1.—Examples of filtered (10-yr weighted moving average, thick curve) and unfiltered (thin curve) values of rainfall showing increasing trends (A, B, C, D, E, F, G, H), decreasing trends (J), and no significant trend (I).

hibited QBO (fig. 2). QBO, it may be pointed out, is often considered as part of the solar cycle (nearly 11 yr) which in turn is intimately related to the ultraviolet emission (Shapiro and Ward 1962, Staley 1963).

## 6. CONCLUSION AND REMARKS

This study shows that the areas having increasing and decreasing trend in the annual rainfall are practically contiguous. However, this trend is not significant over all the stations in the area but only at a few places distributed at random. Wherever this trend is significant, it has persistence or a periodicity of more than 40 yr. QBO is exhibited at several stations in the areas of increasing and

TABLE 4.—Power spectrum results

| Name of station | Class interval of significant periods (yr) | Name of station | Class interval of significant periods (yr) |
|-----------------|--|-----------------|--|
| Trivandrum      | - -  | Indore          | ∞- 88.0†                                   |
| Pamban          | 6.7- 5.8                                   | Ahmadabad       | 17.6- 12.5†                                |
| Fort Cochin     | 5.8- 5.1†                                  | Sagar           | 18.8- 12.2                                 |
|                 | 4.6- 4.1                                   |                 | 4.8- 4.5                                   |
|                 | 3.5- 3.2                                   |                 | 44.0- 26.4                                 |
| Kodakanal       | 4.1- 3.5*                                  | Daltonganj      | 14.6- 12.0†                                |
| Mangalore       | - -  | Dumka           | 29.3- 12.5†                                |
| Bangalore       | 5.6- 5.3                                   |                 | 44.0- 26.4                                 |
|                 | 3.2- 3.0                                   |                 | 2.9- 2.6*                                  |
|                 | 2.2- 2.0                                   |                 | 12.0- 8.8**                                |
| Madras          | 35.2- 25.1                                 |                 | 2.8- 2.2**                                 |
|                 | 13.5- 11.7                                 |                 | 12.5- 8.0*                                 |
|                 | 2.4- 2.3**                                 |                 | 8.0- 6.7                                   |
|                 | 2.3- 2.2                                   |                 | 44.0- 26.4                                 |
| Bellary         | - -  |                 | 26.4- 18.8*                                |
| Belgaum         | ∞-132.0*                                   |                 | 12.0- 7.7†                                 |
|                 | 132.0- 44.0                                |                 | 4.1- 3.8                                   |
| Masulpatnam     | 8.0- 6.7                                   |                 | - -  |
| Hyderabad       | - -  |                 | 4.0- 3.5*                                  |
| Visakhapatnam   | 4.0- 3.7                                   |                 | 2.3- 2.1†                                  |
| Poona           | 4.8- 4.2*                                  |                 | 3.3- 3.0†                                  |
| Bombay          | ∞- 44.0†                                   |                 | 12.5- 8.0†                                 |
|                 | 4.8- 4.5†                                  |                 | 3.2- 2.8                                   |
|                 | 2.4- 2.2†                                  |                 | 14.6- 12.0                                 |
| Jagdapur        | 12.5- 9.7                                  |                 | 2.4- 2.2                                   |
|                 | 9.7- 8.0*                                  |                 | ∞- 44.0                                    |
|                 | 2.5- 2.3                                   |                 | 3.2- 2.6*                                  |
| Akola           | 44.0- 26.4†                                |                 | 14.6- 10.1†                                |
|                 | 26.4- 18.8                                 |                 | 29.3- 17.6                                 |
| Cuttack         | 8.8- 7.7                                   |                 | 14.6- 12.0                                 |
|                 | 3.7- 3.3                                   |                 | 8.8- 7.7                                   |
| Varaval         | ∞- 88.0                                    |                 | 2.4- 2.3                                   |
| Nagpur          | 132.0- 26.4                                |                 | 12.0- 10.1†                                |
| Sambalpur       | 10.5- 7.7                                  |                 | 7.7- 6.9                                   |
|                 | 3.0- 2.9                                   |                 | - -  |
| Sagar Island    | 4.8- 4.5†                                  |                 | 8.8- 6.9†                                  |
|                 | 3.3- 3.0                                   |                 | 2.9- 2.4†                                  |
| Dwarka          | - -  |                 | ∞-132.0†                                   |
| Calcutta        | 4.2- 4.0                                   |                 | 132.0- 44.0                                |
|                 | 4.0- 3.9*                                  |                 | 8.0- 6.7                                   |
|                 | 3.3- 3.2                                   |                 | 2.3- 2.1                                   |
|                 |  |                 | 44.0- 26.4                                 |

\*Significant at 99-percent level  
 †Significant at 95-percent level

decreasing trend. Similarly, the 11-yr cycle (solar cycle) is also exhibited in both these areas. However, both the QBO and 11-yr cycle are present at three stations only. This is an important feature that has to be seriously considered when we seek a physical explanation for the QBO in the rainfall. Presently, QBO in any meteorological parameter is generally believed to be intimately connected with the solar cycle. If the solar phenomena are responsible for the QBO, we should reasonably expect the solar cycle also at the same place. Since this association is absent at certain stations we have studied, it is rather difficult for us to visualize at present how the solar cycle and QBO are related.

**ACKNOWLEDGMENT**

Authors express their sincere thanks to K. Raghavan for going through the manuscript critically and offering very valuable suggestions.




FIGURE 2.—Observing stations from which rainfall data were used. Here, (●) indicates increasing trend, (○) indicates decreasing trend, Q indicates QBO, and underline indicates 11-yr cycle. (Note that all stations indicate either an increasing or decreasing trend; those with significant trends are named in section 3.)

**REFERENCES**

Angell, J. K., Korshover, J., and Carpenter, T. H., "Note Concerning the Period of the Quasi-biennial Oscillation," *Monthly Weather Review*, Vol. 95, No. 3, Mar. 1966, pp. 319-323.

Bhargava, B. N., and Bansal, R. K., "A Quasi-biennial Oscillation in Precipitation at Some Indian Stations," *Indian Journal of Meteorology and Geophysics*, Vol. 20, No. 2, New Delhi, Jan. 1969, pp. 127-128.

Callendar, G. S., "Temperature Fluctuations and Trends Over the Earth," *Quarterly Journal of the Royal Meteorological Society*, Vol. 87, No. 371, London, England, Jan. 1961, pp. 1-12.

Gilman, D. L., Fuglister, F. J., and Mitchell, J. M. Jr., "On the Power Spectrum of 'Red noise,'" *Journal of the Atmospheric Sciences*, Vol. 20, No. 2, Mar. 1963, pp. 182-184.

Jagannathan, P., and Bhalmé, H. N., "Changes in the Pattern of Distribution of Monsoon Rainfall Over India Associated With Sunspot," *Proceedings of the Symposium on Physical and Dynamic Climatology, Leningrad U.S.S.R., August 16-20, 1971* (to be published by World Meteorological Organization, Geneva, Switzerland, 1973).

Kendall, M. G., and Stuart, A., *The Advanced Theory of Statistics*, Hafner Publishing Company, Vol. 2, New York, N.Y., 1961, p. 676.

Koteswaram, P., and Alvi, S. M. A., "Secular Trends and Periodicities in Rainfall at West Coast Stations in India," *Current Science*, Vol. 38, No. 10, Bangalore, India, May 20, 1969, pp. 229-231.

- Kraus, E. B., "Synoptic and Dynamic Aspect of Climatic Change," *Quarterly Journal of the Royal Meteorological Society*, Vol. 86, No. 367, London, England, Jan. 1960, pp. 1-15.
- Landsberg, H. E., Mitchell, J. M. Jr., Crutcher, H. L., and Quinlan, F. T., "Surface Signs of Biennial Atmospheric Pulse," *Monthly Weather Review*, Vol. 91, Nos. 10-12, Oct.-Dec. 1963, pp. 549-556.
- Pramanik, S. K., and Jagannathan, P., "Climatic Changes in India (1)—Rainfall," *Indian Journal of Meteorology and Geophysics*, Vol. 4, No. 4, New Delhi, Oct. 1953, pp. 291-309.
- Reynolds, G., "Rainfall at Bidston, 1867-1951," *Quarterly Journal of the Royal Meteorological Society*, Vol. 79, No. 390, London, England, Jan. 1953, pp. 137-149.
- Shapiro, Ralph, and Ward, Fred, "A Neglected Cycle in Sunspot Numbers?," *Journal of the Atmospheric Sciences*, Vol. 19, No. 6, Nov. 1962, pp. 506-508.
- Staley, D. O., "A Partial Theory of the 26-month Oscillation of the Zonal Wind in the Equatorial Stratosphere," *Journal of the Atmospheric Sciences*, Vol. 20, No. 6, Nov. 1963, pp. 506-515.
- Tukey, J. W., "Sampling Theory of Power Spectrum Estimates," *Proceedings of the Symposium on Applications of Auto-correlation Analysis to Physical Problems, Woods Hole, Massachusetts, June 13-14, 1949*, Office of Naval Research, NAVEXOS-P-735, Washington, D.C., 1950, pp. 47-67.
- Willett, Hurd C., "Temperature Trends of the Past Century," *Centenary Proceedings of the Royal Meteorological Society*, London, England, Mar.-Apr. 1950, pp. 195-206.
- World Meteorological Organization, "Climatic Change," WMO Technical Note No. 79, WMO-No. 195. TP.-100, Geneva, Switzerland, 1966, 80 pp.

[Received September 30, 1971; revised January 10, 1973]

### CORRECTION NOTICE

Vol. 101, No. 2, Feb. 1973, p. 140, left col: the reference to Deland, Raymond J. (1972b) should read Deland, Raymond J., "Traveling Planetary-Scale Temperature Waves in the Stratosphere," *TR* No. 72-5, NSF Grant GA-25820, Geophysical Sciences Laboratory, New York University, Bronx, N.Y., Aug. 1972b, 63 pp.