Ocean and Ice Shelf Tides from *CryoSat-2* Altimetry

EDWARD D. ZARON

Portland State University, Portland, Oregon

(Manuscript received 21 November 2017, in final form 27 February 2018)

ABSTRACT

A new empirical model of ocean tides has been developed for the Weddell Sea, south of 66°S, between 90°W and 0°, using six years of radar altimeter data from the *CryoSat-2* satellite mission. Because of its long ground-track repeat period (368 days) and its diverse measurement modes, low-rate mode (LRM) over the ocean and synthetic aperture radar interferometric mode (SARin) over ice surfaces and parts of the ocean, the *CryoSat-2* data pose a number of challenges for tidal analysis. The space and time sampling properties of the exact repeat, near-repeat, and crossover ground tracks have been analyzed to discover which tides may be estimated using a combination of conventional harmonic analysis and local spatial regression. Using this information, the M_2 , S_2 , K_2 , N_2 , K_1 , O_1 , P_1 , and Q_1 tides have been mapped for both the ocean and floating ice shelves in this domain. Validation against independent in situ data, along with comparison with existing tide models, finds that the *CryoSat-2*-derived tides are consistent with previous estimates and that they are more accurate than other models at the M_2 and S_2 frequencies. The high inclination of the *CryoSat-2* orbit causes the orbit plane to precess relatively slowly, which leads to significantly less accurate estimates of the K_2 tide. This purely empirical model ought to provide improved tidal corrections for studies of low-frequency variability and secular trends in ice shelf thickness, and it suggests that further increases in quantitative accuracy could be achieved by assimilation of *CryoSat-2* data into dynamical tide models.

1. Introduction

The study of ocean tides in the era of satellite altimetry has changed our understanding of tides in the deep ocean (Fu and Cazenave 2001), led to improved quantitative measures of the rate and distribution of barotropic tidal dissipation (Cartwright and Ray 1991; Egbert and Ray 2003), and is leading to insights into smaller-scale coastal and internal tides and their relationship to wind-driven and thermohaline processes (Ray and Cartwright 2001; Ray and Zaron 2011; Müller et al. 2014). These studies have traditionally utilized data from satellite missions in orbits designed with consideration of tidal periodicities (Parke et al. 1987). For example, the orbits of the TOPEX/Poseidon, *Jason-1*, and *Jason-2* missions were selected to avoid aliasing tidal variability into seasonal, annual, or longer periods in order to clearly separate the tidal signals from

secular change (Fu and Cazenave 2001). Likewise, the orbits of the *Geosat*, *Geosat Follow-On*, and sun-synchronous *Envisat*, *ERS-1*, *ERS-2*, and *Satellite with Argos and AltiKa* (SARAL) missions were designed to trade-off between tidal aliasing, spatial resolution, and other mission priorities. Sun-synchronous orbits are troublesome for tidal studies per se, since the largest semidiurnal solar tide S_2 is aliased to zero frequency (Cheng and Andersen 2011). Nonetheless, the missions just mentioned, which shall be referred to as the exact repeat missions (ERM), have been the primary source of information about tides in the ocean, away from coastlines.

In contrast, there have been other satellite altimeter missions in orbits designed to achieve different objectives, for which tides are not a primary concern. The so-called geodetic mission (GM) phases of *Geosat* and *ERS-1* placed the satellites in orbits that traced out a series of closely spaced ground tracks intended to map the mean sea surface at high spatial resolution to infer the small-scale features of the geoid (Sandwell and Smith 1997). Several contemporary missions have also occupied orbits with closely spaced ground tracks, which can be achieved either through precisely controlled

Supplemental information related to this paper is available at the Journals Online website: <https://doi.org/10.1175/JPO-D-17-0247.s1>.

Corresponding author: Edward Zaron, ezaron@pdx.edu

long-repeat orbits or through less-controlled drifting orbits. The *CryoSat-2* (June 2010–present), *Jason-1* (May 2012–June 2013), *SARAL* (July 2016–present), and *Jason-2* (July 2017–present) missions are recent examples for which high-quality GM altimeter data are available (Marks et al. 2013).

Among these missions, the *CryoSat-2* altimeter was designed for studies of the cryosphere (Wingham et al. 2006). Its 88°-inclination orbit covers higher latitudes than other missions, and the Synthetic Aperture Radar/Interferometric Radar Altimeter (SIRAL-2) instrument on *CryoSat-2* is unique in that it is capable of switching between three modes, low-rate mode (LRM), synthetic aperture radar (SAR) mode, and SAR interferometry (SARin) mode, each optimized for different measurements (Bouzinac 2012; Boy et al. 2017). When in the LRM, the SIRAL-2 instrument operates like a conventional pulse-limited radar for measuring the sea surface height. In SAR mode the altimeter uses Doppler and phase information to increase the spatial resolution of the range measurement in the along-track direction, which is useful for identifying floating ice and leads and for determining ice freeboard. Finally, the SARin mode utilizes two antennas to form an interferometer in the across-track direction, and it is used to map steep and highly variable ice and snow surfaces. Unlike the *SARAL* and *Jason* missions, *CryoSat-2* does not carry a microwave radiometer, so the altimeter range cannot be as accurately corrected for wet troposphere path delays (Francis 2007; Bouzinac 2012).

This paper makes a systematic effort to map ocean tides with the *CryoSat-2* altimeter, taking advantage of its unique orbit and capabilities to extend the tidal maps onto the floating ice shelves of the Weddell Sea at latitudes south of 66° (Fig. 1), the limit of the TOPEX/Poseidon and *Jason* orbits. The motivation is twofold. It is both an exploration of the accuracy attainable from GM altimetry and an effort to improve the precision of tidal estimates at high latitude. The latter are important for the utilization of, for example, GRACE data to study the cryosphere and hydrologic cycle, since aliased tidal variability is a significant source of noise for that mission (Ray and Luthcke 2006; Wiese et al. 2016).

The organization of this paper is as follows: section 2 analyzes the sampling properties of the *CryoSat-2* orbit to determine which tidal frequencies are plausible candidates for mapping; section 3 describes the approach taken to reduce nontidal signals and improve the precision of the tidal analysis; section 4 outlines the methodology, which combines harmonic analysis with spatial regression, for identifying the tides and dealing with the

FIG. 1. Analysis domain. Surface type is water (dark blue), floating ice (light blue), continental ice (gray), and land (red), according to the RTopo-2 database (Schaffer et al. 2016). For reference the sites of in situ GPS and water-level data used for validation are shown with white circles; these are discussed later in the text in the context of Fig. 11.

limitations imposed by the long-repeat period of the *CryoSat-2* orbit; section 5 presents cotidal charts derived from *CryoSat-2* and assesses their accuracy by comparing them with in situ data and other tide models; and, finally, sections 6 and 7 summarize and discuss the results in the context of other studies and make suggestions for further research.

2. The *CryoSat-2* orbit and tidal analysis

The feasibility of performing a tidal analysis of the *CryoSat-2* data for a particular tidal frequency depends not only on the characteristics of the *CryoSat-2* orbit but also on the size and stability of the tides in question. Hence, it is useful to begin by considering what is known about tides in the Weddell Sea, and let this information guide a detailed analysis of the orbit.

Although time series of water and ice shelf elevation within the Weddell Sea are sparse, GPS time series, generally shorter than a year, are available on the ice shelves (King et al. 2011), and a long time series of tide gauge data are available on the western side of the Antarctic Peninsula at the Faraday/Vernadsky Base (65.245°S, 64.267°W). Tidal analysis of GPS records by King et al. (2011) finds that the four largest astronomical tides are M_2 , S_2 , K_1 , and O_1 , followed by K_2 , N_2 , P_1 , and Q_1 . Nonlinear overtides with sufficient signal to noise for identification at some stations are NO_1 , MK_3 , MO_3 , and M_4 . Annual and semiannual variability S_a and S_{sa} are identifiable at some stations but at a level barely above the background of low-frequency variability, and they shall not be considered further.

TABLE 1. *CryoSat-2* orbit parameters.

Orbit attribute	Value
Altitude	717.242 km
Inclination	92°
Repeat cycle	368.24 days
Orbits per cycle	5344
Track spacing at equator	7.5 km
Pseudosubcycles	28.33 days, 2.18 days
Orbit period	1.654 h
Mean local solar time drift	$-179.21 \text{ s day}^{-1}$
Longitude of ascending node	309.37°

When mapping ocean tides with satellites the considerations are, generally, the spacing of the ground tracks relative to the horizontal scales of the tidal elevation field and the alias periods of the tidal harmonics in relation to the length of record of the satellite data. The *CryoSat-2* orbit has a long repeat period of 368 days (Table 1), which covers Earth with an array of 10688 ascending and descending ground tracks with an inter-track spacing of about 7.5 km at the equator (Francis 2007). Track spacing reduces with latitude, allowing the polar regions to be mapped with high spatial resolution approximately every 28 days, the length of a near-repeat pseudosubcycle. Figure 2 illustrates the spatial density of tracks in the southern Weddell Sea and Antarctica during a 30-day period encompassing a pseudosubcycle.

Tidal observations would be impractical if only the exact-repeat orbit aliases of *CryoSat-2* were considered. Instead, consider the closely spaced parallel tracks associated with the pseudosubcycles as well as the intersections of ascending and descending tracks. Figure 3 illustrates the time interval Δt between successive measurements within a 30-km-diameter region at 70°S during a representative 60-day period. There are measurements near $\Delta t = 2$ days and 29 days, the pseudosubcycles, and there are even more measurements near $\Delta t = 7$ days and 20 days associated with crossovers. The sample interval associated with orbit crossovers is a complex function of latitude (Wunsch 1989; Morrow et al. 1994), but the histogram of Δt provides a useful starting point for examining the feasibility of a tidal analysis with *CryoSat-2*.

Given the above sampling periodicities, Table 2 lists the alias periods of the ocean tides identified as candidates for analysis. The *CryoSat-2* record extends from July 2010 to present, approximately 2190 days. All of the linear astronomical tides, excluding Q_1 , have alias periods less than 2190 days and are theoretically resolvable at each point along the nominal repeat orbit ($\Delta t = 368$ days). The sample intervals associated with neighboring tracks and crossovers within the 30-km

FIG. 2. *CryoSat-2* ground tracks during a 30-day period. This enlargement of the southern Weddell Sea and Filchner–Ronne Ice Shelf illustrates the density of ground tracks over a 30-day period encompassing a pseudosubcycle; tracks in black occur within a shorter 3-day period. Line thickness increases over the ocean where the altimeter operates in LRM. Notice that the tracks become somewhat irregular when the altimeter operates in SARin mode, which measures the lateral location of off-nadir reflections. The map uses a polar stereographic projection with easting and northing indicated (km).

region lead to shorter alias periods, which, for the most part, are distinct from each other. The M_2 , K_1 , O_1 , P_1 , MO_3 , and MK_3 tides have aliases between 14 and 16 days, associated with the two smallest (and most frequently occurring) Δt sample intervals. The Q_1 , MO_3 , and M_4 tides have aliases close to the semiannual period S_{sa} , and M_2 and O_1 have aliases separated from the annual period S_a by about 10%. The study of alias periods is not definitive, though, since the cross talk between the frequencies depends on the number of samples with the given Δt as well as the size of the harmonics and the character of the noise associated with them. So, for example, the coincidence of the 16-day aliases of MO_3 and M_2 is problematic for identification of the smaller MO_3 . Conversely, the spurious 1-mm-level contribution of MO_3 to M_2 is smaller than the uncertainty of the latter and may be neglected.

Another factor to consider is the distribution of phases sampled. The failure to uniformly sample the phases can lead to errors in nominally well-resolved harmonic constants. For example, the phases of M_2 and K_2 for measurements within the 30-km-diameter disk used above are highly correlated (Fig. 4a). Even though the alias periods are well-separated, the correlated phases

FIG. 3. Histogram of sample intervals, for $\Delta t < 50$ days during a representative 60-day period, within a 30-km-diameter disk at 70°S . Samples near $\Delta t = 2$ and 29 days are associated with pseudosubcycles. The samples near $\Delta t = 6$ and 20 days are associated with intersecting ascending and descending tracks.

lead to correlated errors between harmonic constants at these frequencies. Estimation of M_2 by itself is problematic since there is an excess of measurements near phases -1 and $+2$ radians but a deficit near -3 and $+1$ radians (Fig. 4b). This phase bias is itself variable in longitude, and if it is not corrected it leads to spurious zonal oscillations in the M_2 harmonic constants, similar to the situation with *Geosat* (Cartwright and Ray 1990).

As an example, consider the harmonic constants for M_2 and K_1 across a zonal section through the Weddell Sea at 70°S (Fig. 5). Two estimates of the harmonic constants are shown, one based on data within small, 30-km-diameter bins and the other based on data within larger, $240\text{ km} \times 30\text{ km}$ zonal bins. The estimates from data within small bins display a zonal waviness that is most evident for M_2 , as suggested above. The larger, zonally elongated bins utilize more data and more uniformly sample the phase of the tides, leading to more accurate and stable tidal estimates.

Some of the tidal frequencies are poorly sampled even with the zonal averaging. Because *CryoSat-2* is in a high-inclination orbit, the orbit plane precesses very slowly and its orientation is nearly fixed with respect to the lunar orbit plane (Capderou 2005). Table 3 lists the phase increments (degrees of phase change) associated with the sample intervals listed in Table 2. All the

TABLE 2. Tidal alias periods. Alias periods (days) are given for sample intervals that occur when *CryoSat-2* data are accumulated within 30-km spatial bins at 70°S .

	Sample interval Δt (days)				
	368.2396	28.9410	19.4246	7.5180	1.9983
M_2	800	371	42	16	14
S_2	768	245	129	209	576
K_2	743	715	438	98	267
N_2	2095	225	113	30	9
K_1	1486	1430	41	16	535
O_1	1262	294	347	638	14
P_1	1591	209	52	15	277
Q_1	5106	195	55	26	9
NO_1	3170	962	86	28	29
MO_3	2187	164	47	16	7
MK_3	1734	500	1682	115	15
M_4	4633	185	288	140	7

frequencies except K_2 and S_2 undergo a phase change of 45° or more for the shortest sample intervals, $\Delta t = 2$ days and $\Delta t = 7.5$ days, most relevant to the spatially binned data. The K_2 is particularly problematic since the phase changes are small for all sample intervals less than the repeat period, $\Delta t = 368$ days. Figure 6 shows the phases of the K_1 and K_2 tides sampled by *CryoSat-2* within the $240\text{ km} \times 30\text{ km}$ bin used above. Although the alias period of K_1 is twice that of K_2 , within any given 20-day period, the K_1 tide is sampled at two phases, approximately 170° apart. In contrast, over the same period, the K_2 tide will be sampled within a narrower range of only about 30° . Consequently, one should expect temporally correlated error, such as elevation change associated with snowfall or the slow movement of ice floes, to have a much greater influence on estimates of K_2 , as compared with the other tides.

Figure 7 illustrates how the accuracy of the harmonic constants is expected to vary as a function of the zonal bin length. Let $\mathbf{F}$ be the design matrix with elements $\zeta_j(t_k) = f_j(t_k)\exp[i[\Omega_j(t_k) + u_j(t_k)]]$ following the customary notation for the nodal amplitude and phase factors f_j and u_j and the phase function Ω_j , computed from the astronomical arguments using the Doodson numbers defining the j_{th} tidal harmonic (Doodson 1921). The elements of the least squares normal matrix, $\mathbf{G} = (\mathbf{F}^T\mathbf{F})^{-1}$, may be used to compute $\tilde{\rho}$, the theoretical cross-correlation matrix for the uncertainty of the harmonic constants:

$$\tilde{\rho} = (\mathbf{G})_d^{-1/2} \mathbf{G} (\mathbf{G})_d^{-1/2}, \quad (1)$$

where $(\mathbf{G})_d$ is equal to the diagonal matrix with elements equal to the main diagonal of $\mathbf{G}$ (Cherniawsky et al. 2001). Let $\rho(X, Y)$ be defined as the maximum

FIG. 4. *CryoSat-2* observations relative to tidal phase within a 30-km-diameter disk of ocean at 70°S. (a) Timing of *CryoSat-2* observations in the M_2 – K_2 phase plane. (b) Probability distribution function (PDF) of M_2 phase as seen by *CryoSat-2*. The nonuniform sampling leads to correlated errors and biased estimates of harmonic constants if *CryoSat-2* data are analyzed within small bins.

correlation of in-phase and quadrature components for tides X and Y , the quantity plotted in Fig. 7. For the latitude of 70°S, a zonal bin length of 240 km corresponds to a local minimum in $\rho(M_2, K_2)$. The zonal average harmonic constants at this scale should be a factor of $1/\sqrt{8}$ more accurate than the estimates using the 30-km-diameter disk because of the additional quantity of data in each estimate, and the error correlation should be reduced by more than a factor of 10. Note that $\rho(M_2, K_2)$ is shown in the figure, but the other correlations involving the semidiurnal tides behave similarly, although they are of smaller magnitude. The error correlation of the

diurnal tides, the largest of which is $\rho(K_1, P_1)$, also can be reduced by zonal averaging, but the main reduction occurs within the first 100 km and there is little improvement with further averaging. Table 2 indicates that NO_1 is likely to be the most easily identified nonlinear overtide; however, preliminary harmonic analyses found that it could not be reliably estimated. Figure 7 shows the $\rho(NO_1, K_1)$ correlation and indicates that additional spatial averaging would be necessary to decorrelate the NO_1 error from the K_1 error. Since it appears that the nonlinear overtones cannot be identified at the scales at which they are likely generated, they

FIG. 5. Tidal harmonic constants computed from *CryoSat-2* data along a section at 70°S. The in-phase (black) and quadrature (red) components of the harmonic constants for K_1 (small circles and dashed lines) and M_2 (large circles and solid lines) are shown. Circles indicate results using data within 30-km-diameter bins; correlated error is evident as noise with a wavelength of about 200 km. Lines indicate results using data within $240 \text{ km} \times 30 \text{ km}$ zonal bins (length scale indicated near 330°E); the noise is greatly reduced.

TABLE 3. Phase increments as a function of sample interval. The change in phase (degrees) associated with the sample periods listed in Table 2 is given.

	Sample interval Δt [days]				
	368.2396	28.9410	19.4246	7.5180	1.9983
M_2	166	28	168	170	50
S_2	173	43	54	13	1
K_2	178	15	16	28	3
N_2	63	46	62	91	76
K_1	89	7	172	166	1
O_1	105	35	20	4	51
P_1	83	50	134	179	3
Q_1	26	54	126	102	77
NO_1	42	11	82	96	25
MO_3	61	64	148	175	101
MK_3	76	21	4	24	49
M_4	29	56	24	19	100

shall henceforth not be considered as candidates for analysis with *CryoSat-2*.

Figure 8 shows how $\rho(M_2, K_2)$ varies as a function of latitude poleward of 66°S. The spatial density of the ground tracks increases poleward up to the orbit inclination of 88°, and this has the dual effect of increasing the diversity of tidal phases sampled and increasing the number of data, both of which lead to a reduction in the error correlation. To optimize the accuracy of the harmonic analysis, below, the data-binning length is permitted to vary as a function of latitude, according to the dashed line, which has been subjectively aligned with the minimum of the error correlation field in Fig. 8. The bin size varies as a linear function of latitude from 75 km at 85°S to 300 km at 66°S:

$$\Delta L = 300 \text{ km} \times (92^\circ - |\theta|)/26^\circ, \quad (2)$$

where θ indicates latitude.

3. Corrections for nontidal signals and the mean surface

The discussion has so far emphasized how the *CryoSat-2* orbit influences the theoretical accuracy of harmonic analysis. Another consideration is the character of nontidal signals in the *CryoSat-2* measurements. When *CryoSat-2* is operating in LRM, the instrumental precision is similar to that of other nadir altimeter missions (Garcia et al. 2014). The main difference is that the satellite does not carry a microwave radiometer for making observations of atmospheric water vapor, so the associated path delay must be estimated from a numerical weather prediction model (Bouzinac 2012). The path delays and other geophysical corrections that are applied to the *CryoSat-2* level-2 baseline-C product

FIG. 6. Phase sampling of K_1 and K_2 . The phases of the K_1 tide (black) and the K_2 tide (red) are shown, as sampled within a 240 km $\times$ 30 km zonal patch in the Weddell Sea. Although the alias period of the K_2 tide is shorter than the alias period of the K_1 tide (Table 2), *CryoSat-2* frequently samples the K_1 tide at two phases almost 180° apart. In contrast, only a small range of phases of the K_2 tide are sampled in any short window.

are listed in Table 4 (Webb and Hall 2016). Note that the data product includes both the elevation measurements and corrections; therefore, the nominal ocean tide correction was removed from the elevation product before subsequent analysis, below.

The elevations of the mean sea and ice surfaces along the orbit ground tracks are not known with the same precision as they are known along the historical ERM ground tracks. A large component of the mean ice surface variability, which can exceed meters of elevation per kilometer, is at scales smaller than the typical ocean tide signals, and if this is not removed it adds significantly to the nontidal noise of the measurements. In principle, the mean surface could be identified as part of the harmonic tidal analysis, since it simply corresponds to the zero frequency, but this approach is not feasible because of the disparate scales of tidal and mean surface variability. The discussion of the orbit indicates that spatial averaging is necessary to reduce error correlations, at least for the present length of the data record. The approach taken here is to separately estimate the mean surface by binning the data at a resolution of about 3.5 km, which is a compromise between minimizing the loss of spatial resolution while improving the stability of the estimate. The number of data per grid cell shows the influence of off-nadir returns in SARin mode, which

FIG. 7. Theoretical error correlation inferred from the normal matrix used in the least squares determination of the harmonic constants. The largest error correlation among each frequency group is as follows: $\rho(M_2, K_2)$ (dashed black line) is the worst combination of semidiurnals, and the other semidiurnal frequency pairs follow a similar pattern; $\rho(K_1, P_1)$ (solid black line) is the worst combination involving diurnals; $\rho(NO_1, K_1)$ (red line) is the worst correlation involving the NO_1 nonlinear overtide. The other overtones have poorer (larger) error correlations and are not shown.

tend to occur from high points in the illuminated scene (Fig. 9). The number of data per grid cell also shows differences as a function of instrument mode, with many more SARin returns compared to LRM, for example.

The standard deviation of elevation within the 3.5-km grid cells provides additional insight into the variability observed by *CryoSat-2*. Figure 10 shows that small-scale features are visible, even over the ocean, which appear to be the result of discrete icebergs. Variability near the grounding lines and other sites of macroscale roughness appears to reflect subgrid-scale features.

In addition to the above considerations, the final component of data processing prior to harmonic analysis involves some basic quality control to identify spurious data. When the satellite is in the vicinity of rough surfaces, there are occasionally large outliers, which deviate by ± 4 m or more from the mean surface, while the typical tidal range is much less than this. It is hypothesized that these outliers are the symptom of a waveform retracker error related to the interferometer beamforming or phase unwrapping in SARin mode. It appears that the off-nadir reflection

FIG. 8. The theoretical error correlation $\rho(M_2, K_2)$ is shown as a function of latitude when the data are binned within a circular disk on the local tangent plane. The dashed white line indicates the optimal data-binning scale, which is approximately a linear function of latitude.

point is not correctly identified since the range measurement is too large (i.e., negative sea or ice surface elevation), as if a distant off-nadir reflection had been mapped to an apparent near-nadir point (Gomez-Enri et al. 2016).

4. Spatially coupled harmonic analysis

Section 2 described how the extent of spatial averaging is related to identification of tidal harmonics, given the orbit characteristics of the *CryoSat-2* altimeter. Harmonic analysis can be directly applied to time series of the data within averaging bins; however, more accurate estimates of the tidal elevation can be obtained using a model for spatial dependence of the tide within the bins. The approach used here is a spatial signal model that consists of a P th-order polynomial in a locally defined tangent plane:

$$\hat{\eta}_j(x, y) = \sum_{p=0}^P \sum_{q=0}^{P-p} a_{jpq} x^p y^q, \quad (3)$$

where $x = (\phi - \phi_0)r_e \cos(\theta_0)$ and $y = (\theta - \theta_0)r_e$ define the local coordinate system around the latitude–longitude coordinates (θ_0, ϕ_0) and r_e is the radius of Earth. Using the same temporal function $\zeta_j(t)$ as above, the complete model for the j th tidal harmonic

TABLE 4. Geophysical and path delay corrections: ESA *Cryosat-2* L2 baseline-C data release. Detailed information concerning the source of each correction is found in Webb and Hall (2016).

Range correction	Model or data source
Dry troposphere	ECMWF
Wet troposphere	ECMWF
Ionospheric path delay	Global Ionosphere Map (GIM) and Bent models
Long-period polar “tide”	Instantaneous Polar Location from CNES
Solid Earth tide	Cartwright
Tidal loading	FES2004
Long-period equilibrium ocean tide	FES2004
Dynamic atmosphere (ocean)	Modèle d’Onde de Gravité 2D (MOG2D)
Inverse barometer (ice shelves)	ECMWF surface pressure
Sea state bias	<i>Jason-2</i> CLS sea state bias (SSB) model

is given by $\eta_j(x, y, t) = \hat{\eta}_j(x, y)\zeta_j(t)$. The coefficients a_{jpq} are computed by finding the least squares solution to

$$\mathbf{h} = \mathbf{F}\mathbf{a}, \quad (4)$$

where $\mathbf{h}$ is the vector $\{h_k\}$ of *CryoSat-2* data within the patch centered on (θ_0, ϕ_0) , $\mathbf{F}$ is the design matrix with entries $x_k^p y_k^q \zeta_j(t_k)$, and $\mathbf{a}$ is the vector $\{a_{jpq}\}$ of coefficients to be determined.

The ordinary least squares solution of the above is $\mathbf{a} = (\mathbf{F}^T \mathbf{F})^{-1} \mathbf{F}^T \mathbf{h}$, and this shall be referred to as the nominal tide estimate. To examine the stability of the estimates, three other tide estimates are also obtained. The first of these is denoted the $2\Delta L$ estimate, and it is computed with double the nominal data bin scale [(2)]. The next estimate is the weighted least squares (WLS) estimator, $\mathbf{a} = (\mathbf{F}^T \mathbf{W} \mathbf{F})^{-1} \mathbf{F}^T \mathbf{W} \mathbf{h}$, where the weight $\mathbf{W}$ is a diagonal matrix equal to the inverse of the square of the variability displayed in Fig. 10; the WLS estimator therefore accounts for spatial variability of the nontidal variance, primarily due to moving sea ice and the ice shelf edge. The final estimate is the iteratively reweighted least squares (IRWLS) estimator, and it is based on the Huber loss function (Huber 1964); the IRWLS estimator robustifies the WLS estimator and reduces the influence of data that contain little tidal variance (Huber 1981).

Local models for η_j are computed on a regular grid of coordinates defining overlapping tangent planes, and the model coefficients are used to estimate the in-phase and quadrature components of the harmonic constants at the origin of the tangent plane (θ_0, ϕ_0) . The diameter of the binning disk in the tangent plane is given by the latitude-dependent scale given by (2) and indicated in

FIG. 9. Number of observations per grid cell for Z_0 , the mean surface. The polygonal boundary between low and high data counts (e.g., near 75°S, 60°W) corresponds to the boundary of the instrument mode mask, with more returns from SARin than LRM.

Fig. 8. It was hypothesized that the bias of the rudimentary estimate based on the averaged data ($P = 0$) could be reduced by using quadratic and higher-order terms ($P \geq 2$) in the spatial model; however, in practice, the estimation of additional model parameters with $P > 1$ led to suboptimal results, even with increased ΔL . Hence, the results shown below simply use the linear spatial model, $P = 1$. Note that some topographic features (e.g., the Antarctic Peninsula) are smaller than the scale of the tangent plane, and if this were not taken into account, the methodology would spuriously combine data from opposite sides of geographic features. To avoid this, as the data are assembled within the tangent plane a test is made to see if continental land or grounded ice is present along a line between the data point and the origin of the tangent plane (θ_0, ϕ_0) , using the Refined Topography dataset, 1-min resolution, version 2 (RTopo-2), surface type definition (Schaffer et al. 2016). As a result, the harmonic constants are based on fewer data and are of reduced accuracy approaching the land boundaries.

5. Results

The above-described methodology was applied to estimating the tides from the *CryoSat-2* measurements.

Table 5 provides a quantitative comparison with in situ GPS (King et al. 2011) and historical tide gauge data (King and Padman 2005; L. Padman et al. 2017, unpublished data) at the locations labeled in Fig. 11, indicated previously in Fig. 1. The error metric tabulated is the root-mean-square vector error (rmsve), the root-mean-square scalar elevation difference computed from the complex-valued harmonic constants. For comparison, the table also displays rmsve for the

FIG. 10. Standard deviation of elevation within each grid cell defining the mean surface elevation Z_0 . The Filchner–Ronne Ice Shelf Front is visibly prominent (cf. Fig. 1); other ice shelf fronts are less visible as they are associated with more sea ice and small-scale variability.

Goddard/Grenoble ocean tide model, version 4.10c (GOT4.10c), and Circum-Antarctic Tidal Simulation (CATS2008) tide models with respect to the same in situ datasets. The GOT4.10c model was developed from a response-type analysis of ERM altimetry (Ray 1999); however, these data were generally too noisy to be used in the Weddell Sea where sea ice is often present, and GOT4.10c reverts to a combination of TPX07.2 (Egbert and Erofeeva 2002) and Finite Element Solution ocean tide model 2004 (FES2004; Lyard et al. 2006). The CATS2008 model is an updated version of the model described in Padman et al. (2002) and employs the same data-assimilative methodology as other high-latitude tide models (Padman and Erofeeva 2004).

The table contains columns corresponding to the four different solutions mentioned previously: the nominal, the enlarged analysis window ($2\Delta L$), the WLS, and the robustified weighted least squares (IRWLS) estimators. The most noteworthy difference among the solutions is the large reduction in error for K_2 using the $2\Delta L$ estimator, but this is not unexpected given the analysis of phase sampling in section 2. In general, though, the differences between the *CryoSat-2*-based estimates are much smaller than the errors with respect to the in situ data, and none of the estimates is unambiguously better than the others, so no attempt is made to analyze the differences in detail. Visually, the $2\Delta L$ and WLS fields are noticeably smoother than the others, and for display purposes the $2\Delta L$ solution shall be used for plotting K_2 and Q_1 . Otherwise, the nominal solution is shown.

Cotidal charts for the four largest tides, M_2 , S_2 , K_1 , and O_1 , are presented in Fig. 12. They are qualitatively very similar to what has been obtained previously using data assimilation of ERM altimetry and in situ data

TABLE 5. Summary comparison of GOT4.10c, CATS2008, and *CryoSat-2* tides vs in situ data at sites indicated in Fig. 11. The columns under *CryoSat-2* correspond to the four estimators mentioned in section 4.

	GOT4.10c	CATS2008	rmsve (cm)			
			<i>CryoSat-2</i>			
			Nominal	$2\Delta L$	WLS	IRWLS
M_2	4.3	4.5	3.1	3.9	3.2	3.2
S_2	8.6	7.6	6.6	6.8	6.5	6.5
N_2	1.3	2.0	1.7	1.5	1.6	1.7
K_2	2.5	2.0	4.5	4.0	4.8	4.8
K_1	4.5	2.4	2.9	2.8	2.6	2.5
O_1	5.6	1.2	2.0	2.1	2.0	1.9
P_1	1.4	1.2	1.4	1.1	1.2	1.1
Q_1	0.8	1.0	1.5	1.1	1.4	1.4

(Padman et al. 2002). The main differences compared to CATS2008 are smaller-scale structures near the grounding line of the Filchner–Ronne Ice Shelf; however, these are probably too close to the analysis boundary to be reliable. Another qualitative difference is the shape of the amphidrome at the edge of the ice shelf; it is more confined to the ice shelf edge in the empirical solution. The channels between the islands in the Filchner–Ronne Ice Shelf are probably too small to reliably estimate the tide within them, and the empirical solution displays some questionable structure, a slight waviness, to the west of Berkner Island. The structure of the K_1 and O_1 tides along the continental shelf break, near 70°S, differs from CATS2008; the region of maximum amplitude is more spread out in the *CryoSat-2* analysis.

Cotidal charts for the smaller tides, K_2 , N_2 , P_1 and Q_1 , are displayed in Fig. 13. The structure of the K_2 amphidrome at the edge of the Filchner–Ronne Ice Shelf is anomalous and is thought to be due to the poor sampling of the K_2 phase combined with the larger level of low-frequency ice surface variability at the edge of the ice shelf. The structure of the N_2 tide is quite similar to the larger semidiurnals, except that it is noisier, presumably due to the larger alias period and the long synodic periods needed to separate this tide from K_1 , P_1 , and Q_1 (5106, 3170, and 6612 days, respectively, for $\Delta t = 368$ sampling). As might be expected, the P_1 and Q_1 tides are also noisy as a consequence of interference with N_2 , but the same structure as the other diurnals is visible.

Errors are very large at two sites, LAR2 and EE55, which lie close to the grounding line, and these sites are excluded from the summary statistics in Table 5. The choice to exclude these sites is somewhat arbitrary, however, since other sites close to land (A020) or the grounding line (FR03) are retained, but the quantity of

FIG. 11. Data sites used for model validation are located (a) on the Larsen Ice Shelf, (b) on the Filchner–Ronne Ice Shelf, and (c) in the eastern Weddell Sea. Errors at sites marked with red dots, LAR2 and EE55, are tabulated in the [appendix](#) but excluded from the root-mean-square summary statistics owing to their close proximity to the ice shelf grounding line.

satellite data going into the tidal analysis at these sites is larger than at the excluded sites because of the shape of the coastline. Amplitude and phase errors for each station are provided in the [appendix](#) (Tables A1–A8), including these two stations. Note that both GOT4.10c and CATS2008 assimilated the L. Padman et al. (2017, unpublished data) data within the southern Weddell Sea subdomain (Fig. 11c; sites named with prefix “A”).

Table 5 indicates that the *CryoSat-2*-derived tides are more accurate than either model for the M_2 and S_2 tides. The rmsve of M_2 is reduced broadly across the Filchner–Ronne Ice Shelf and the western end of the eastern Weddell Sea, at the ocean sites. The rmsve of S_2 is reduced at the eastern Weddell Sea sites, except for the easternmost site, A020, and there is no pattern to changes in the error on the Filchner–Ronne Ice Shelf. On the Larsen Ice Shelf the rmsve of S_2 is

reduced by 3 cm to 6 cm, but it is only reduced at the northernmost station, LAR1, for M_2 . Note that the tidal estimates at LAR2 and LAR3 are based on much less data than at LAR1 owing to their proximity to land.

The rmsve of *CryoSat-2* N_2 falls in between the values for GOT4.10c and CATS2008. An unusual feature of the N_2 tide is that *CryoSat-2* agrees better with the CATS2008 model than it does with the GPS data (e.g., at LAR1, LAR3, FR10, and FR03; Table A3).

The rmsve of K_2 , 4 cm, is about twice the rmsve of the GOT4.10c and CATS2008 models. Given the sampling properties of *CryoSat-2*, it is surprising that the K_2 results are accurate even at this level.

The rmsve of the K_1 , O_1 , and P_1 tides fall between the values for GOT4.10c and CATS2008, where the latter has the lowest error. The rmsve of the Q_1 tides is very

FIG. 12. The four largest tides mapped from *CryoSat-2* data. The amplitude is shown with the color scale, and phase lines are shown in 30° increments. The latitude-dependent averaging scale ΔL is indicated with the scale bars near the eastern boundary of the plots. Note the continuity of the tidal fields across the ice shelf fronts (cf. Figs. 1 and 11).

slightly worse than either of the models. It is noteworthy that the *CryoSat-2* tides are the most accurate of the set at the LAR1 station, which has the best

exposure to the ocean. The largest K_1 error, 7.7 cm, is at the FR10 station, closest to the Filchner–Ronne Ice Shelf edge, and if it were not for this error, the rms

FIG. 13. As in Fig. 12, but for the four smaller tides mapped from *CryoSat-2* data. For the purposes of visualization, the $2\Delta L$ estimate is shown for the K_2 and Q_1 fields.

FIG. 14. *CryoSat-2* vs CATS2008 differences and uncertainty estimates. (top and middle) The root-mean-square vector difference between the *CryoSat-2* and CATS2008 tidal fields, denoted $\Delta(M_2)$, $\Delta(K_1)$, $\Delta(S_2)$, and $\Delta(O_1)$. (bottom) The corresponding formal error estimate for the nominal (ordinary least squares) solutions, denoted $\sigma(M_2)$ and $\sigma(K_1)$. Formal errors are elevated near boundaries and along parts of the ice shelf front where small-scale variability is also large (cf. Fig. 10). Note the different color scales used in each panel.

would be almost the same as that of CATS2008. Errors at stations close to the ice shelf edge are also larger for O_1 , P_1 , and Q_1 .

6. Discussion

The accuracy of tides estimated empirically from *CryoSat-2* are comparable to those computed from ERM altimetry using data assimilation. Only the accuracy of K_2 is seriously deficient, with rmsve substantially larger than that of the CATS2008 model. The spatial distribution of the difference fields between the CATS2008 and the *CryoSat-2*-derived tide, in Fig. 14, displays features similar to the corrections to tidal maps

inferred using GRACE (Han et al. 2007; Egbert et al. 2009) and ICESat data (Padman et al. 2008). The largest differences in the semidiurnal tides occur on the ice shelves, whereas the largest differences in the diurnal tides occur in association with the continental shelf break. The influence of frictional processes on tidal dynamics is significant in precisely these regions, and the difference fields likely indicate errors in the CATS2008 frictional parameterizations (Padman et al. 2008).

The interpretation of the *CryoSat-2* versus CATS2008 difference as error in the CATS2008 model is consistent with the formal estimates of uncertainty provided by the least squares estimates, except near land. Uncertainties

for the M_2 and K_1 tides shown in Fig. 14 have been computed from the residual variance of the nominal estimator (Cherniawsky et al. 2001). Uncertainty is dominated by the proximity to land and variability associated with the ice shelf edge. Unfortunately, the formal uncertainty estimate does not correspond in detail to the actual errors observed where in situ data are available (appendix); however, the relatively small number of sites makes a detailed comparison inconclusive in any case.

The spatial structure of errors in the tide estimates are likely to reflect a combination of factors. These include the geographic variability in the nontidal signals, such as the movement of small-scale icebergs (Fig. 10). There is also likely to be geographic variability in measurement error, such as seasonal changes in radar cross section owing to firn density or surface roughness (Remy and Parouty 2009). As evidence of this possibility, consider the cotidal charts of the annual and semiannual period “tides” in Fig. 15, which were made using the same methodology as described above. It is hypothesized that the signals on the ice shelves, which far exceed their amplitude at the Faraday/Vernadsky tide gauge (not shown) and GPS stations (King et al. 2011), are a measure of seasonal variability of snow and ice properties on the ice shelf. There is also error related to the waveform retracking and geolocation of the range measurement (Nilsson et al. 2016; Wingham et al. 2006), and this is likely to be significant in the vicinity of complex surfaces such as the ice shelf edge or, in some cases, the grounding line. Finally, the last component of geographically correlated error is the systematic error related to the spatial signal model and its ability to represent small-scale features of the tidal fields, particularly near the coastlines and grounding lines.

The signal model used for the spatially coupled harmonic analysis is ad hoc, and it would be preferable to assimilate the *CryoSat-2* data into a hydrodynamic model, building on the work of Egbert and Erofeeva (2002) and Padman and Erofeeva (2004). The spatial structure would then be consistent with the hypothesized tidal hydrodynamics, within the limits imposed by imperfect knowledge of frictional effects and baroclinic dynamics. The use of *CryoSat-2* data would increase by orders of magnitude the amount of data currently assimilated from tide gauges, GPS, and ICESat. It is also conceivable that assimilation might permit some of the smaller linear tides and nonlinear overtones to be identified in the Weddell Sea. Although it was not emphasized above, *CryoSat-2* has favorable sampling characteristics for the MK_3 and M_4 overtones.

FIG. 15. The annual (S_a) and semiannual (S_{sa}) tides mapped from *CryoSat-2*. Since the amplitude of both tides far exceeds the amplitude observed by in situ instruments, it is hypothesized that the apparent S_a and S_{sa} tides are caused by seasonal variations in the surface properties of the ice shelves.

7. Conclusions

An empirical model for the eight largest diurnal and semidiurnal tides in the Weddell Sea and adjoining ice shelves has been developed using *CryoSat-2* altimeter data. The sampling properties of the *CryoSat-2* orbit were used to define the size of the latitude-dependent data window to reduce interference among the tidal frequencies, enabling the stable estimation of tidal amplitude and phase. The tides are estimated with a spatially coupled least squares harmonic analysis in which the in-phase and quadrature components of the tide are assumed to vary linearly across a locally defined tangent plane. The fields are mapped by performing this type of local regression centered at points on a $1/8^\circ \times 1/2^\circ$ (latitude by longitude) grid at latitudes poleward of 66°S , from 90°W to 0° .

The tide model inferred from *CryoSat-2* has been validated using a collection of GPS and historical tide measurements and by comparing it with the GOT4.10c and CATS2008 models. The *CryoSat-2* solution is more accurate than CATS2008 for the

TABLE A1. Comparisons of M_2 at GPS and historical tide stations.

	Amplitude (m)			Phase (°)			Vector error (cm)	
	CAT	C2a	In situ $\pm$ error	CAT	C2a	In situ $\pm$ error	CAT	C2a
LAR3	0.891	0.872	0.890 ± 0.026	-119.2	-113.6	-117.5 ± 1.6	1.8	4.5
LAR2	1.039	0.985	0.966 ± 0.002	-111.3	-96.2	-114.5 ± 0.1	6.5	21.9
LAR1	0.897	0.886	0.902 ± 0.023	-104.7	-108.1	-108.7 ± 1.5	4.4	1.3
EE55	1.168	1.190	1.219 ± 0.004	81.5	84.7	84.9 ± 0.2	6.2	2.0
FR06	0.882	0.878	0.890 ± 0.021	88.4	94.0	94.6 ± 1.4	6.8	1.1
FR07	0.308	0.263	0.256 ± 0.002	112.2	132.4	129.3 ± 0.3	6.9	1.1
FR10	0.265	0.198	0.231 ± 0.007	-12.5	-37.5	-33.7 ± 1.9	6.8	2.5
FR09	0.735	0.704	0.684 ± 0.002	21.5	19.9	19.7 ± 0.2	3.9	1.4
FR02	0.806	0.748	0.765 ± 0.029	-2.8	-5.5	-5.4 ± 2.2	3.8	1.2
FR03	1.154	1.072	1.111 ± 0.026	12.6	8.8	10.6 ± 1.5	4.2	3.6
FR05	1.164	1.176	1.193 ± 0.003	70.1	73.1	73.2 ± 0.1	5.0	1.2
A007	0.551	0.552	0.571	-115.0	-116.1	-118.0	2.5	1.9
A041	0.556	0.565	0.588	-119.4	-119.9	-120.2	2.3	1.6
A042	0.734	0.733	0.772	-110.1	-112.7	-112.3	3.4	2.8
A066	0.695	0.719	0.701	-125.9	-131.1	-127.3	1.3	3.6
A010	0.546	0.548	0.560	-139.1	-138.1	-130.0	6.3	5.6
A009	0.445	0.446	0.460	-155.9	-154.6	-159.0	2.1	2.6
A002	0.430	0.428	0.339	-155.5	-155.8	-162.1	7.2	7.0
A008	0.405	0.386	0.410	-167.3	-166.4	-171.0	1.9	2.8
A020	0.346	0.313	0.347	-175.8	-179.6	-177.0	0.5	2.8

M_2 , S_2 , and N_2 tides, and its accuracy is only slightly worse (1 to 8 mm) for the K_1 , O_1 , P_1 , and Q_1 tides. The high-inclination orbit of *CryoSat-2* poorly samples the K_2 tide, which has errors that are twice as large as those from CATS2008.

The results indicate that *CryoSat-2* should be a useful data source for future data-assimilative tide models. Errors in the present tidal estimates arise from instrument and

retracker errors, as well as from the simplified spatial signal model. Future efforts will investigate *CryoSat-2* data that have been reprocessed using alternative methods to reduce retracker errors (Nilsson et al. 2016); however, assimilation of the present data into a hydrodynamic model ought to improve upon the simple spatial model employed in the present study and, perhaps, enable the identification of even smaller tides.

TABLE A2. Comparisons of S_2 at GPS and historical tide stations.

	Amplitude (m)			Phase (°)			Vector error (cm)	
	CAT	C2a	In situ $\pm$ error	CAT	C2a	In situ $\pm$ error	CAT	C2a
LAR3	0.638	0.599	0.406 ± 0.023	-89.2	-86.1	-81.2 ± 3.3	17.2	14.0
LAR2	0.757	0.647	0.649 ± 0.002	-81.3	-88.1	-85.2 ± 0.2	8.3	2.3
LAR1	0.639	0.597	0.355 ± 0.024	-74.1	-78.7	-75.9 ± 3.5	20.2	17.2
EE55	0.764	0.681	0.777 ± 0.004	113.5	114.1	113.6 ± 0.3	0.9	6.8
FR06	0.566	0.545	0.451 ± 0.020	121.7	127.4	142.3 ± 2.5	15.1	11.2
FR07	0.195	0.170	0.154 ± 0.002	159.4	168.1	171.3 ± 0.7	3.9	1.3
FR10	0.138	0.134	0.140 ± 0.008	-2.7	-43.5	-16.6 ± 2.9	2.4	4.5
FR09	0.449	0.367	0.400 ± 0.002	46.7	43.1	43.6 ± 0.3	3.8	2.3
FR02	0.545	0.471	0.485 ± 0.030	19.3	14.2	18.5 ± 3.3	4.3	2.8
FR03	0.777	0.705	0.816 ± 0.025	36.2	34.0	37.3 ± 2.0	3.0	8.5
FR05	0.758	0.706	0.750 ± 0.002	101.4	99.9	100.7 ± 0.2	0.8	3.2
A007	0.394	0.409	0.388	-92.7	-94.9	-98.0	2.6	2.2
A041	0.395	0.397	0.403	-96.6	-98.1	-97.9	0.8	0.4
A042	0.514	0.516	0.525	-87.9	-88.7	-90.7	2.0	1.4
A066	0.491	0.471	0.484	-103.1	-106.5	-105.4	1.5	1.1
A010	0.391	0.392	0.390	-116.2	-116.9	-118.0	0.9	0.5
A009	0.326	0.333	0.330	-132.8	-134.2	-137.0	1.7	1.2
A002	0.317	0.315	0.255	-132.5	-134.5	-136.1	4.6	4.3
A008	0.302	0.293	0.300	-143.9	-145.3	-149.0	1.9	1.4
A020	0.265	0.256	0.259	-159.3	-166.2	-152.0	2.4	4.5

TABLE A3. Comparisons of N_2 at GPS and historical tide stations.

	Amplitude (m)			Phase (°)			Vector error (cm)	
	CAT	C2a	In situ $\pm$ error	CAT	C2a	In situ $\pm$ error	CAT	C2a
LAR3	0.139	0.139	0.135 ± 0.021	-135.1	-132.1	-142.9 ± 9.5	1.3	1.8
LAR2	0.163	0.179	0.137 ± 0.002	-127.6	-130.4	-132.7 ± 0.9	2.1	3.0
LAR1	0.140	0.135	0.132 ± 0.026	-121.0	-121.7	-134.9 ± 10.4	2.4	2.2
EE55	0.195	0.142	0.183 ± 0.004	64.9	53.1	64.7 ± 1.3	0.9	3.7
FR06	0.149	0.128	0.122 ± 0.021	71.7	81.9	66.0 ± 10.6	2.1	2.5
FR07	0.054	0.056	0.036 ± 0.001	88.1	97.6	101.5 ± 2.6	1.4	1.4
FR10	0.050	0.046	0.036 ± 0.006	-26.2	-22.8	-48.3 ± 11.5	1.5	1.4
FR09	0.126	0.137	0.104 ± 0.002	1.8	5.4	5.2 ± 1.0	1.6	2.3
FR02	0.156	0.126	0.102 ± 0.029	-19.4	-26.2	-18.1 ± 17.4	3.8	2.0
FR03	0.211	0.186	0.141 ± 0.028	-5.9	-5.9	1.0 ± 12.7	5.1	3.5
FR05	0.194	0.178	0.175 ± 0.003	53.5	46.8	56.1 ± 0.8	1.4	2.0
A007	0.087	0.095	0.092	-123.9	-125.9	-133.0	1.1	0.9
A041	0.088	0.094	0.089	-129.4	-132.2	-130.5	0.1	0.4
A042	0.122	0.103	0.119	-120.1	-119.2	-122.8	0.5	1.2
A066	0.113	0.112	0.102	-137.0	-147.5	-143.1	1.1	0.9
A010	0.087	0.085	0.090	-150.7	-154.1	-151.0	0.2	0.5
A009	0.070	0.062	0.070	-168.1	-163.1	-167.0	0.1	0.7
A002	0.068	0.070	0.052	-167.8	-169.4	174.2	1.7	1.8
A008	0.064	0.056	0.080	179.6	178.3	180.0	1.1	1.7
A020	0.055	0.057	0.055	160.5	163.5	154.0	0.4	0.7

Acknowledgments. Discussions with Laurie Padman and Richard Ray are gratefully acknowledged regarding the ice shelf tidal dynamics and altimeter sampling. The *CryoSat-2* level-2 baseline-C surface elevation product was provided by Sara Fleury at the Center for Topographic studies of the Ocean and Hydrosphere (CTOH), who also provided valuable insight concerning the data quality flags (ctoh_product@legos.obs-mip.fr). The data

are available by request (<http://ctoh.legos.obs-mip.fr/products/alongtrack-data/cryosat2-request-form>). The Faraday/Vernadsky tide gauge time series was obtained from the University of Hawaii Sea Level Center archive of research-quality water-level records (<http://uhslc.soest.hawaii.edu>). This project was supported by the NASA Ocean Surface Topography Science Team, Grant NNX13AH06G.

TABLE A4. Comparisons of K_2 at GPS and historical tide stations.

	Amplitude (m)			Phase (°)			Vector error (cm)	
	CAT	C2a	In situ $\pm$ error	CAT	C2a	In situ $\pm$ error	CAT	C2a
LAR3	0.175	0.156	0.121 ± 0.018	-88.5	-58.1	-81.7 ± 8.0	4.0	4.7
LAR2	0.210	0.509	0.193 ± 0.002	-80.2	-62.9	-84.9 ± 0.5	1.6	23.9
LAR1	0.176	0.138	0.106 ± 0.018	-73.0	-52.6	-76.4 ± 8.9	5.0	4.2
EE55	0.213	0.250	0.235 ± 0.003	114.9	148.2	112.5 ± 0.9	1.7	10.6
FR06	0.159	0.155	0.134 ± 0.014	123.1	148.9	141.8 ± 7.1	3.8	2.0
FR07	0.059	0.146	0.051 ± 0.001	155.4	-156.2	169.3 ± 1.5	1.1	7.6
FR10	0.041	0.028	0.045 ± 0.005	5.7	-73.2	-16.8 ± 6.8	1.2	2.6
FR09	0.127	0.092	0.120 ± 0.001	45.5	100.1	43.7 ± 0.7	0.5	7.3
FR02	0.165	0.099	0.147 ± 0.022	20.2	29.1	17.6 ± 9.2	1.4	3.8
FR03	0.226	0.132	0.238 ± 0.022	36.4	50.2	37.2 ± 5.4	0.8	8.0
FR05	0.210	0.188	0.223 ± 0.002	102.7	138.3	100.2 ± 0.5	1.2	9.8
A007	0.106	0.116	0.108	-89.9	-86.2	-97.0	0.9	1.6
A041	0.107	0.109	0.121	-94.5	-111.8	-98.3	1.1	2.1
A042	0.143	0.156	0.156	-83.5	-88.8	-90.7	1.6	0.4
A066	0.136	0.170	0.152	-100.1	-112.7	-108.6	1.9	1.5
A010	0.107	0.126	0.110	-114.8	-124.1	-119.0	0.6	1.3
A009	0.091	0.114	0.100	-131.4	-121.9	-138.0	1.0	2.3
A002	0.088	0.105	0.073	-131.3	-128.0	-133.7	1.1	2.3
A008	0.084	0.095	0.080	-142.6	-152.2	-146.0	0.4	1.3
A020	0.074	0.113	0.074	-159.4	-168.3	-160.0	0.1	2.9

TABLE A5. Comparisons of K_1 at GPS and historical tide stations.

	Amplitude (m)			Phase (°)			Vector error (cm)	
	CAT	C2a	In situ $\pm$ error	CAT	C2a	In situ $\pm$ error	CAT	C2a
LAR3	0.310	0.299	0.320 ± 0.018	5.8	5.3	9.8 ± 3.4	1.7	2.3
LAR2	0.352	0.356	0.362 ± 0.002	7.1	-4.8	5.6 ± 0.4	0.9	4.6
LAR1	0.342	0.348	0.365 ± 0.018	19.7	10.4	15.3 ± 2.6	2.6	2.4
EE55	0.371	0.390	0.433 ± 0.003	73.7	71.0	72.4 ± 0.4	4.4	3.2
FR06	0.343	0.335	0.390 ± 0.026	76.5	79.3	76.0 ± 3.8	3.3	4.2
FR07	0.314	0.388	0.376 ± 0.002	62.6	64.6	59.7 ± 0.3	4.6	2.5
FR10	0.278	0.200	0.309 ± 0.018	43.5	42.4	41.2 ± 3.5	2.3	7.7
FR09	0.312	0.325	0.352 ± 0.002	48.5	46.8	46.5 ± 0.3	3.0	1.9
FR02	0.356	0.381	0.399 ± 0.022	40.4	38.0	39.8 ± 3.5	3.1	1.6
FR03	0.412	0.501	0.457 ± 0.027	44.4	40.4	42.3 ± 3.7	3.4	3.3
FR05	0.370	0.410	0.424 ± 0.003	68.6	66.7	67.6 ± 0.3	3.9	1.1
A007	0.350	0.317	0.361	12.5	7.9	17.0	2.1	4.9
A041	0.314	0.327	0.338	11.9	12.0	14.6	2.0	1.3
A042	0.301	0.313	0.317	11.8	11.7	11.9	1.2	0.3
A066	0.295	0.308	0.298	9.7	8.2	9.4	0.3	0.8
A010	0.280	0.280	0.290	5.9	3.4	5.0	0.8	0.9
A009	0.261	0.269	0.270	-0.1	3.4	-2.0	0.9	1.8
A002	0.256	0.253	0.257	-0.2	1.6	3.2	1.1	0.6
A008	0.262	0.249	0.270	-5.0	-2.3	-8.0	1.1	2.4
A020	0.246	0.255	0.253	-9.4	-14.4	-9.0	0.5	1.7

APPENDIX

CryoSat-2 and CATS2008 Tides at in situ Stations

Tables A1 through A8 present tidal amplitude and phase at each of the sites where in situ data were available for comparison. The sites, LAR1, LAR2, LAR3, EE5, FR02, FR03, FR05, FR07, FR09, and

FR10, are taken from the supplementary material of King et al. (2011); they are tides inferred from the GPS elevation records. As mentioned in the text, the sites LAR2 and EE55, where the largest errors are found, are also the closest sites to the grounding line, and errors at these sites are not used when computing the rmsve summary reported in Table 5.

TABLE A6. Comparisons of O_1 at GPS and historical tide stations.

	Amplitude (m)			Phase (°)			Vector error (cm)	
	CAT	C2a	In situ $\pm$ error	CAT	C2a	In situ $\pm$ error	CAT	C2a
LAR3	0.273	0.263	0.277 ± 0.017	-3.3	1.2	0.2 ± 3.4	1.2	1.0
LAR2	0.300	0.346	0.297 ± 0.002	-4.4	-18.0	-7.9 ± 0.4	1.3	5.3
LAR1	0.318	0.312	0.317 ± 0.018	8.1	-0.3	3.7 ± 2.9	1.7	1.6
EE55	0.382	0.364	0.415 ± 0.002	62.8	92.3	62.4 ± 0.3	2.3	14.6
FR06	0.358	0.359	0.388 ± 0.027	65.0	69.8	64.0 ± 3.7	2.1	3.4
FR07	0.351	0.338	0.373 ± 0.002	50.5	54.9	47.6 ± 0.3	2.0	4.1
FR10	0.299	0.282	0.301 ± 0.016	34.7	27.4	33.0 ± 3.0	0.7	2.4
FR09	0.319	0.337	0.332 ± 0.002	40.5	41.8	39.0 ± 0.3	1.1	1.2
FR02	0.355	0.356	0.373 ± 0.022	32.2	31.2	31.1 ± 3.7	1.4	1.2
FR03	0.409	0.414	0.418 ± 0.028	36.0	35.1	35.5 ± 3.8	0.7	0.3
FR05	0.377	0.401	0.402 ± 0.002	58.5	64.1	58.3 ± 0.4	1.7	2.8
A007	0.310	0.312	0.302	4.2	5.7	6.0	0.9	0.7
A041	0.329	0.318	0.342	5.1	5.6	6.3	1.0	1.7
A042	0.314	0.301	0.320	4.6	3.1	5.2	0.5	1.6
A066	0.309	0.290	0.311	2.2	6.4	2.8	0.3	2.0
A010	0.298	0.287	0.310	-2.1	-2.6	-1.0	0.9	1.7
A009	0.282	0.288	0.290	-8.2	-2.8	-8.0	0.6	1.9
A002	0.278	0.271	0.271	-8.6	-4.2	-5.6	1.1	0.5
A008	0.285	0.262	0.290	-13.0	-11.8	-13.0	0.4	2.0
A020	0.270	0.266	0.270	-18.4	-18.4	-17.0	0.5	0.6

TABLE A7. Comparisons of P_1 at GPS and historical tide stations.

	Amplitude (m)			Phase (°)			Vector error (cm)	
	CAT	C2a	In situ $\pm$ error	CAT	C2a	In situ $\pm$ error	CAT	C2a
LAR3	0.096	0.103	0.103 ± 0.022	25.8	3.3	9.5 ± 11.0	2.1	0.8
LAR2	0.102	0.132	0.117 ± 0.003	28.4	−9.9	4.9 ± 1.1	3.3	2.5
LAR1	0.100	0.112	0.118 ± 0.020	34.8	12.5	15.0 ± 10.6	2.9	0.6
EE55	0.126	0.113	0.143 ± 0.003	72.2	58.6	74.4 ± 1.1	1.2	3.2
FR06	0.117	0.124	0.132 ± 0.027	74.8	73.9	75.8 ± 12.6	1.1	0.6
FR07	0.113	0.092	0.122 ± 0.002	61.1	44.8	56.3 ± 0.9	1.0	2.6
FR10	0.097	0.106	0.102 ± 0.021	42.4	26.9	40.9 ± 10.7	0.4	1.8
FR09	0.107	0.128	0.118 ± 0.002	47.8	47.8	50.7 ± 1.1	0.9	0.8
FR02	0.121	0.125	0.136 ± 0.028	40.5	44.9	39.7 ± 10.2	1.0	1.1
FR03	0.139	0.143	0.154 ± 0.033	44.1	40.3	42.5 ± 11.3	1.1	0.9
FR05	0.125	0.144	0.144 ± 0.003	67.4	68.7	69.7 ± 1.1	1.4	0.2
A007	0.102	0.103	0.119	10.6	10.3	17.0	1.5	1.5
A041	0.102	0.104	0.113	13.6	12.3	13.8	0.8	0.7
A042	0.100	0.085	0.105	12.4	6.7	12.1	0.4	1.5
A066	0.097	0.077	0.103	10.2	15.9	9.3	0.5	2.0
A010	0.092	0.100	0.100	6.8	2.9	4.0	0.6	0.1
A009	0.086	0.069	0.090	0.7	−15.3	−4.0	0.6	1.9
A002	0.084	0.078	0.087	0.6	−9.3	2.0	0.2	1.3
A008	0.086	0.079	0.090	−4.2	−3.6	−9.0	0.6	1.0
A020	0.081	0.056	0.084	−9.0	0.3	−10.0	0.2	2.2

The sites A002, A007, A008, A009, A010, A020, A041, A042, and A066 are historical sites taken from the Antarctic Tide Gauge Database (King and Padman 2005; L. Padman et al. 2017, unpublished data). The sites used are ones where the tide was computed from bottom pressure gauges, rather than tiltmeters.

The sites are listed in order clockwise around the Larsen and Filchner–Ronne Ice Shelves, starting from the west, and from west to east in the eastern Weddell Sea.

Values tabulated under the “CAT” column are from the CATS2008 model, and values under the “C2a” column are from the nominal estimate discussed in the text.

TABLE A8. Comparisons of Q_1 at GPS and historical tide stations.

	Amplitude (m)			Phase (°)			Vector error (cm)	
	CAT	C2a	In situ $\pm$ error	CAT	C2a	In situ $\pm$ error	CAT	C2a
LAR3	0.066	0.027	0.063 ± 0.018	6.4	20.4	-17.7 ± 16.4	1.9	3.2
LAR2	0.069	0.050	0.061 ± 0.002	7.8	−17.9	-16.9 ± 2.1	2.1	0.8
LAR1	0.072	0.070	0.070 ± 0.018	13.8	−17.4	-19.1 ± 14.9	2.8	0.1
EE55	0.086	0.247	0.093 ± 0.003	53.2	76.5	56.0 ± 1.4	0.6	11.5
FR06	0.081	0.063	0.077 ± 0.025	55.1	61.9	44.0 ± 18.9	1.1	1.8
FR07	0.081	0.059	0.086 ± 0.002	40.6	51.2	37.5 ± 1.5	0.5	2.2
FR10	0.069	0.078	0.063 ± 0.017	26.3	0.2	16.2 ± 14.6	0.9	1.7
FR09	0.073	0.084	0.075 ± 0.002	31.7	41.7	31.7 ± 1.6	0.2	1.1
FR02	0.080	0.082	0.077 ± 0.024	24.4	32.6	22.5 ± 16.3	0.3	1.0
FR03	0.092	0.073	0.082 ± 0.029	27.9	38.3	31.0 ± 22.3	0.8	1.0
FR05	0.085	0.091	0.093 ± 0.003	49.0	41.4	50.5 ± 1.6	0.6	1.0
A007	0.073	0.074	0.074	−0.9	0.6	−2.0	0.1	0.2
A041	0.076	0.068	0.082	−1.4	3.4	−0.4	0.5	1.1
A042	0.072	0.047	0.075	−1.7	−29.9	−1.3	0.2	2.9
A066	0.071	0.065	0.069	−4.1	−1.4	−5.8	0.2	0.5
A010	0.069	0.071	0.080	−8.4	−9.0	−8.0	0.8	0.7
A009	0.066	0.076	0.070	−14.6	−17.0	−11.0	0.4	0.7
A002	0.065	0.066	0.065	−14.8	−19.4	−15.7	0.1	0.3
A008	0.066	0.076	0.080	−19.9	−24.0	−19.0	1.0	0.6
A020	0.063	0.041	0.063	−24.7	−6.3	−26.0	0.1	2.0

REFERENCES

- Bouzinac, C., 2012: CryoSat product handbook. ESRIN-ESA and Mullard Space Science Laboratory Rep., 90 pp., <http://emits.sso.esa.int/emits-doc/ESRIN/7158/CryoSat-PHB-17apr2012.pdf>.
- Boy, F., J.-D. Desjonquères, N. Picot, T. Moreau, and M. Raynal, 2017: CryoSat-2 SAR-mode over oceans: Processing methods, global assessment, and benefits. *IEEE Trans. Geosci. Remote Sens.*, **55**, 148–158, <https://doi.org/10.1109/TGRS.2016.2601958>.
- Capderou, M., 2005: *Satellites Orbits and Missions*. Springer, 545 pp.
- Cartwright, D. E., and R. D. Ray, 1990: Oceanic tides from Geosat altimetry. *J. Geophys. Res.*, **95**, 3069–3090, <https://doi.org/10.1029/JC095iC03p03069>.
- , and —, 1991: Energetics of global ocean tides from Geosat altimetry. *J. Geophys. Res.*, **96**, 16 897–16 912, <https://doi.org/10.1029/91JC01059>.
- Cheng, Y., and O. B. Andersen, 2011: Multimission empirical ocean tide modeling for shallow waters and polar seas. *J. Geophys. Res.*, **116**, C11001, <https://doi.org/10.1029/2011JC007172>.
- Cherniawsky, J. Y., M. G. G. Foreman, W. R. Crawford, and R. F. Henry, 2001: Ocean tides from TOPEX/Poseidon sea level data. *J. Atmos. Oceanic Technol.*, **18**, 649–664, [https://doi.org/10.1175/1520-0426\(2001\)018<0649:OTFTPS>2.0.CO;2](https://doi.org/10.1175/1520-0426(2001)018<0649:OTFTPS>2.0.CO;2).
- Doodson, A. T., 1921: The harmonic development of the tide-generating potential. *Philos. Trans. Roy. Soc. London*, **100A**, 305–329, <https://doi.org/10.1098/rspa.1921.0088>.
- Egbert, G. D., and S. Y. Erofeeva, 2002: Efficient inverse modeling of barotropic ocean tides. *J. Atmos. Oceanic Technol.*, **19**, 183–204, [https://doi.org/10.1175/1520-0426\(2002\)019<0183:EIMOBO>2.0.CO;2](https://doi.org/10.1175/1520-0426(2002)019<0183:EIMOBO>2.0.CO;2).
- , and R. D. Ray, 2003: Semi-diurnal and diurnal tidal dissipation from TOPEX/Poseidon altimetry. *Geophys. Res. Lett.*, **30**, 1907–1910, <https://doi.org/10.1029/2003GL017676>.
- , S. Y. Erofeeva, S.-C. Han, S. B. Luthcke, and R. D. Ray, 2009: Assimilation of GRACE tide solutions into a numerical hydrodynamic inverse model. *Geophys. Res. Lett.*, **36**, L20609, <https://doi.org/10.1029/2009GL040376>.
- Francis, C. R., 2007: CryoSat mission and data description. ESTEC Noordwijk Tech. Doc. CS-RP-ESA-SY-0059, 82 pp.
- Fu, L.-L., and A. Cazenave, 2001: *Satellite Altimetry and Earth Sciences*. International Geophysics Series, Vol. 69, Academic Press, 463 pp.
- Garcia, E. S., D. T. Sandwell, and W. H. Smith, 2014: Retracking CryoSat-2, Envisat and Jason-1 radar altimetry waveforms for improved gravity field recovery. *Geophys. J. Int.*, **196**, 1402–1422, <https://doi.org/10.1093/gji/ggt469>.
- Gomez-Enri, J., A. Scozzari, F. Soldovieri, J. Coca, and S. Vignudelli, 2016: Detection and characterization of ship targets using CryoSat-2 altimeter waveforms. *Remote Sens.*, **8**, 193, <https://doi.org/10.3390/rs8030193>.
- Han, S.-C., R. D. Ray, and S. B. Luthcke, 2007: Ocean tidal solutions in Antarctica from GRACE inter-satellite tracking data. *Geophys. Res. Lett.*, **34**, L21607, <https://doi.org/10.1029/2007GL031540>.
- Huber, P. J., 1964: Robust estimation of a location parameter. *Ann. Math. Stat.*, **35**, 73–101, <https://doi.org/10.1214/aoms/1177703732>.
- , 1981: *Robust Statistics*. John Wiley and Sons, 312 pp.
- King, M. A., and L. Padman, 2005: Accuracy assessment of ocean tide models around Antarctica. *Geophys. Res. Lett.*, **32**, L23608, <https://doi.org/10.1029/2005GL023901>.
- , —, K. Nicholls, P. J. Clarke, G. H. Gudmundsson, B. Kulesa, A. Shepherd, and N. Gourmelen, 2011: Ocean tides in the Weddell Sea: New observations on the Filchner-Ronne and Larsen C ice shelves and model validation. *J. Geophys. Res.*, **116**, C06006, <https://doi.org/10.1029/2011JC006949>.
- Lyard, F., F. Lefevre, T. Letellier, and O. Francis, 2006: Modelling the global ocean tides: Modern insights from FES2004. *Ocean Dyn.*, **56**, 394–415, <https://doi.org/10.1007/s10236-006-0086-x>.
- Marks, K. M., W. H. Smith, and D. T. Sandwell, 2013: Significant improvements in marine gravity from ongoing satellite missions. *Mar. Geophys. Res.*, **34**, 137–146, <https://doi.org/10.1007/s11001-013-9190-8>.
- Morrow, R., R. Coleman, J. Church, and D. Chelton, 1994: Surface eddy momentum flux and velocity variances in the Southern Ocean from Geosat altimetry. *J. Phys. Oceanogr.*, **24**, 2050–2071, [https://doi.org/10.1175/1520-0485\(1994\)024<2050:SEMAV>2.0.CO;2](https://doi.org/10.1175/1520-0485(1994)024<2050:SEMAV>2.0.CO;2).
- Müller, M., J. Cherniawsky, M. G. Foreman, and J.-S. von Storch, 2014: Seasonal variation of the M2 tide. *Ocean Dyn.*, **64**, 159–177, <https://doi.org/10.1007/s10236-013-0679-0>.
- Nilsson, J., A. Gardner, L. Sandberg Sørensen, and R. Forsberg, 2016: Improved retrieval of land ice topography from CryoSat-2 data and its impact for volume-change estimation of the Greenland ice sheet. *Cryosphere*, **10**, 2953–2969, <https://doi.org/10.5194/tc-10-2953-2016>.
- Padman, L., and S. Erofeeva, 2004: A barotropic inverse tidal model for the Arctic Ocean. *J. Geophys. Res.*, **31**, L02303, <https://doi.org/10.1029/2003GL019003>.
- , H. A. Fricker, R. Coleman, S. Howard, and S. Erofeeva, 2002: A new tidal model for the Antarctic ice shelves and seas. *Ann. Glaciol.*, **34**, 247–254, <https://doi.org/10.3189/172756402781817752>.
- , S. Y. Erofeeva, and H. A. Fricker, 2008: Improving Antarctic tide models by assimilation of ICESat laser altimetry over ice shelves. *Geophys. Res. Lett.*, **35**, L22504, <https://doi.org/10.1029/2008GL035592>.
- Parke, M. E., R. H. Stewart, D. L. Farless, and D. E. Cartwright, 1987: On the choice of orbits for an altimetric satellite to study ocean circulation and tides. *J. Geophys. Res.*, **92**, 11 693–11 707, <https://doi.org/10.1029/JC092iC11p11693>.
- Ray, R. D., 1999: A global ocean tide model from TOPEX/Poseidon altimetry: GOT99.2. NASA Tech. Memo. NASA/TM-1999-209478, 58 pp., http://denali.gsfc.nasa.gov/personal_pages/ray/MiscPubs/19990089548_1999150788.pdf.
- , and D. E. Cartwright, 2001: Estimates of internal tide energy fluxes from TOPEX/Poseidon altimetry: Central North Pacific. *Geophys. Res. Lett.*, **28**, 1259–1262, <https://doi.org/10.1029/2000GL012447>.
- , and S. B. Luthcke, 2006: Tide model errors and GRACE gravimetry: Towards a more realistic assessment. *Geophys. J. Int.*, **167**, 1055–1059, <https://doi.org/10.1111/j.1365-246X.2006.03229.x>.
- , and E. D. Zaron, 2011: Non-stationary internal tides observed with satellite altimetry. *Geophys. Res. Lett.*, **38**, L17609, <https://doi.org/10.1029/2011GL048617>.
- Remy, F., and S. Parouty, 2009: Antarctic ice sheet and radar altimetry: A review. *Remote Sens.*, **1**, 1212–1239, <https://doi.org/10.3390/rs1041212>.
- Sandwell, D. T., and W. H. F. Smith, 1997: Marine gravity anomaly from Geosat and ERS 1 satellite altimetry. *J. Geophys. Res.*, **102**, 10 039–10 054, <https://doi.org/10.1029/96JB03223>.

- Schaffer, J., R. Timmermann, J. E. Arndt, C. Mayer, M. Morlighem, and D. Steinhage, 2016: A global, high-resolution data set of ice sheet topography, cavity geometry, and ocean bathymetry. *Earth Syst. Sci. Data*, **8**, 543–557, <https://doi.org/10.5194/essd-8-543-2016>.
- Webb, E., and A. Hall, 2016: Geophysical corrections in level 2 CryoSat data products. European Space Agency Rep. ESRIN-EOP-GMQ/IDEAS+/IDEAS-VEG-IPF-MEM-1288, 27 pp., http://wiki.services.esoportal.org/tiki-download_wiki_attachment.php?attId=4287.
- Wiese, D. N., B. Killett, M. M. Watkins, and D.-N. Yuan, 2016: Antarctic tides from GRACE satellite accelerations. *J. Geophys. Res. Oceans*, **121**, 2874–2886, <https://doi.org/10.1002/2015JC011488>.
- Wingham, D. J., and Coauthors, 2006: CryoSat: A mission to determine the fluctuations in Earth's land and marine ice fields. *Adv. Space Res.*, **37**, 841–871, <https://doi.org/10.1016/j.asr.2005.07.027>.
- Wunsch, C., 1989: Sampling characteristics of satellite orbits. *J. Atmos. Oceanic Technol.*, **6**, 891–907, [https://doi.org/10.1175/1520-0426\(1989\)006<0891:SCOSO>2.0.CO;2](https://doi.org/10.1175/1520-0426(1989)006<0891:SCOSO>2.0.CO;2).