Advanced Baseline Imager Cloud-Top Trajectories and Properties of Electrified Snowfall Flash Initiation

SEBASTIAN S. HARKEMA,^a EMILY B. BERNDT,^b JOHN R. MECIKALSKI,^a AND ALANA CORDAK^c

^a *Department of Atmospheric and Earth Science, University of Alabama in Huntsville, Huntsville, Alabama*

^b *NASA Marshall Space Flight Center, Huntsville, Alabama*

^c *Atmospheric Sciences Program, The University of Georgia, Athens, Georgia*

(Manuscript received 5 January 2021, in final form 30 September 2021)

ABSTRACT: Using gridded and interpolated derived motion winds from the Advanced Baseline Imager (ABI), a Lagrangian cloud-feature tracking technique was developed to create and document trajectories associated with electrified snowfall and changes in cloud characteristics leading up to the initiation of lightning, respectively. This study implemented a thundersnow detection algorithm and defined thundersnow initiation (TSI) as the first group in a flash detected by the Geostationary Lightning Mapper when snow was occurring. A total of 10 ABI channels and four multispectral [e.g., red–green–blue (RGB)] composites were analyzed to investigate characteristics that lead up to TSI for 16 644 thundersnow (TSSN) flashes. From the 10.3- μm channel, TSI trajectories were associated with a median decrease of 12.2 K in brightness temperature (T_B) 1 h prior to TSI. Decreases in the reflectance component of the 3.9- μm channel indicated that TSI trajectories were associated with ice crystal collisions and/or particle settling at cloud top. The nighttime microphysics, day cloud phase distinction, differential water vapor, and airmass RGBs were examined to evaluate the microphysical and environmental changes prior to TSI. For daytime TSI trajectories, the predominant colors associated with the day cloud phase distinction RGB transitioned from cyan to yellow/green, physically representing cloud growth and glaciation at cloud top. Gold/orange hues in the differential water vapor RGB indicated that some trajectories were associated with dry upper-level air masses prior to TSI. The analysis of ABI characteristics prior to TSI, and subsequently relating those characteristics to physical processes, inherently increases the fundamental understanding and ability to forecast TSI; thus, providing additional lead time into changes in surface conditions (i.e., snowfall rates).

KEYWORDS: Cloud tracking/cloud motion winds; Lightning; Snowfall; Cloud microphysics; Glaciation; Satellite observations; Nowcasting

1. Introduction

Many studies have examined the electrification mechanisms of thunderstorms that precede the initiation of lightning flashes (i.e., Paluch and Sartor 1973; Jayarante et al. 1983). The leading in-cloud charging theory (i.e., noninductive charging) involves the collisions of graupel and ice crystals in the presence of supercooled liquid water and water vapor (Reynolds et al. 1957; Takahashi 1978; Saunders et al. 2006). The Bergeron process relates these ice collisions and supercooled liquid water to the growth of ice crystals in the mixed-phase region and can serve as a proxy for noninductive charging (Reynolds et al. 1957; Harris et al. 2010). Therefore, understanding the transition from a water cloud to an ice cloud (i.e., glaciation) and fluctuations in a cloud's updraft depth/volume plays a pivotal role in understanding where lightning is initiated (Tilles et al. 2019). From a meteorological satellite perspective, the solar reflected component of near-infrared imagery can be useful for understanding glaciation, while infrared (IR) imagery can be used as a proxy for

cloud updraft strength (Adler and Fenn 1979; Strabala et al. 1994; Mecikalski et al. 2010a,b).

In contrast to warm season convective storms, few studies have investigated the potential relationships between satellite observations and lightning initiation in snowfall [i.e., thundersnow (TSSN)]. Cold-season clouds may contain graupel, ice crystals, and supercooled liquid water at the same temperatures over a deep layer, like that observed in warm-season clouds. Therefore, electrified snowfall is possible if charge separation is enabled within cold-season clouds (e.g., Takahashi et al. 1999; Trapp et al. 2001; Rauber et al. 2014; Schrom and Kumjian 2016; Takahashi et al. 2017). Winter storms are also associated with weaker updrafts that have a more slanted vertical structure in comparison to summertime convection; thus, TSSN events are rare in comparison to summertime lightning activity, and have been more challenging to quantify with regards to the underlying physical processes responsible for their occurrence (e.g., Schultz 1999; Market et al. 2002). Even though TSSN is relatively rare—compared to summertime lightning—studies have found that TSSN can be useful to forecast surface snowfall rates (e.g., Market and Becker 2009) and estimate snowfall accumulation totals (e.g., Crowe et al. 2006). Additionally, TSSN flashes are statistically significant with regards to longer flash durations, higher optical energies, and larger areas from a Geostationary Lightning Mapper (GLM; Goodman et al. 2013) perspective (Harkema et al. 2019). Therefore, understanding cloud microphysical

Supplemental information related to this paper is available at the Journals Online website: <https://doi.org/10.1175/WAF-D-21-0003.s1>.

Corresponding author: Sebastian S. Harkema, sebastian.harkema@nasa.gov

changes prior to the development of TSSN can help provide additional lead time to changes in surface conditions.

To identify relationships between lightning initiation and satellite observations for summertime convective clouds, [Harris et al. \(2010\)](#) examined 10 *Geostationary Operational Environmental Satellite (GOES)-12* IR fields and their trends an hour leading up to lightning initiation. Of the 10 IR fields examined (i.e., single channel, channel differences, and temporal trends), 8 were shown to be statistically significant in predicting lightning initiation in manually tracked cloud features. Similar results were found using the same 10 IR fields in *Meteosat Second Generation (MSG)-8* ([Matthee and Mecikalski 2013](#)). In a recent study, [Thiel et al. \(2020\)](#) compared GOES-East Advanced Baseline Imager (ABI; [Schmit et al. 2005](#)) data and gridded GLM products (e.g., flash extent density) and found that summertime lightning production was favored when 10.3- μm brightness temperature (T_B) were colder than 240 K. [Elsenheimer and Gravelle \(2019\)](#) used high-resolution GOES-East data, specifically the day cloud phase distinction ([Shimizu 2015](#)) multispectral composite, to show value in short-term forecasting the onset of summertime cloud-to-ground lightning in storms. Prior to the initial cloud-to-ground lightning activity, [Elsenheimer and Gravelle \(2019\)](#) found that on multiple occasions the day cloud phase distinction red-green-blue (RGB) multispectral imagery transitioned from cyan to a yellow/green color, which physically represents the development of an updraft and glaciation at cloud top. Additionally, this color change was used to identify locations where noninductive charging occurs in a thundercloud, like techniques used in [Harris et al. \(2010\)](#) and [Matthee and Mecikalski \(2013\)](#). These studies demonstrate how useful satellite imagery, related to convective cloud updraft strength, cloud depth, and glaciation, can be when trying to understand lightning initiation when associated with convective storms.

Electrified snowfall characteristics have been examined through surface observations (e.g., [Market et al. 2002](#)), and ground- (e.g., [Schultz et al. 2018](#)) and space-based ([Harkema et al. 2019, 2020](#)) lightning detection systems. In recent years, new advances in satellite technology (e.g., lightning observations from geostationary orbit) have allowed for the development of a thundersnow detection algorithm (TDA; [Harkema et al. 2019, 2020](#)) that can objectively identify TSSN flashes, by fusing GLM and derived snowfall rates via the merged snowfall rate product ([Meng et al. 2017a,b](#)). The GLM data structure consists of events, groups, and flashes where GLM events are single pixels illuminated beyond a background threshold. In contrast, GLM groups are adjacent GLM events in a single 2-ms frame. The TDA uses GLM groups to identify TSSN flashes in a variety of storm environments (e.g., lake-effect snow, synoptic-scale heavy-banded snowfall). [Cordak et al. \(2020\)](#) used GLM flash data that was processed through the TDA to examine the associated cloud characteristics in ABI single channel and multispectral composite imagery for the 13–17 April 2018 winter storm investigated in [Harkema et al. \(2019, 2020\)](#). The purpose of their analysis was to demonstrate potential satellite-based precursors to TSSN initiation (TSI; as defined below). Approximately half of a total of 44 flashes observed in their study were associated with

decreases in ABI T_B , corresponding to TSI. In an initial effort to understand TSI, [Cordak et al. \(2020\)](#) analyzed cloud-top characteristics in an Eulerian framework where cloud features were tracked independently from TSSN flash occurrences and categorized by single ABI pixels. Changes in cloud properties (e.g., height, phase) were tracked only at locations of TSSN. As a consequence of tracking TSSN relative cloud-property changes, the complete temporal evolution of the cloud characteristics leading up to TSI were not considered.

Prior to this study, the TDA was updated to enhance the determination of rain-to-snow transition regions to improve identification of TSSN. The TDA uses the underlying radar mosaic data of the merged snowfall rate product ([Meng et al. 2017a,b](#)), which has been noted that it may not represent the rain-to-snow transition zone well ([S. Martinaitis 2019](#), CIMMS/NSSL, personal communication). Misrepresentation of the rain-to-snow transition zone resulted in missing data (i.e., pixels not being classified as snow), thus leading to an underestimation of the number of merged snowfall rate pixels associated with TSSN. To alleviate these concerns, the TDA threshold of a GLM flash being classified as TSSN was lowered from an average of ≥ 700 merged snowfall rate pixels to an average of ≥ 500 merged snowfall rate pixels associated with GLM group/flash data. The average merged snowfall rate pixel count of 500 represents $\sim 70\%$ of pixels, within the TDA search radius around GLM data, being classified as snow. As a result, the reduction in the TDA threshold allowed for more GLM flashes to be classified along the edges of snowbands (i.e., rain-to-snow transition region), and/or on the edges of merged snowfall rate data gaps. For a complete description of the TDA methodology, see [Harkema et al. \(2019, 2020\)](#).

For this study, we made use of the improved TDA and define TSI as the first occurrence of a GLM group for each of the GLM flashes classified as TSSN. Our definition of TSI, as the first GLM group, was based on the concept used to identify the initiation locations of lightning flashes defined by the initial radiation source emission recorded by lightning mapping arrays ([Maggio et al. 2005](#)). However, the actual initial breakdown location may be different compared to the first GLM group location because GLM group locations represent radiance weighted centroids of the underlying GLM event data. Further research would be needed to test the validity of this assumption but is beyond the scope of this work.

To expand upon the work of [Cordak et al. \(2020\)](#), this study examined T_B trends (ΔT_B) and cloud processes prior to TSI in a Lagrangian framework. A methodology to track cloud features was developed to observe and improve our understanding of the full cloud-top evolution prior to TSI. The ability to track cloud features, however, is nontrivial and involves some level of subjectivity in how to define them. Some techniques track discrete cloud features using the coldest T_B —or coldest $\sim 25\%$ of T_B values—around a feature of interest per imagery scene from geostationary satellite data (e.g., [Lensky and Rosenfeld 2006](#)). Other studies track convection based on a blend of radar and satellite data (e.g., [Mecikalski and Bedka 2006](#); [Mecikalski et al. 2013](#); [Sieglaff et al. 2013](#); [Cintineo et al.](#)

2020). These techniques work well for discrete meso- γ (2–20 km) cloud features but can be problematic when tracking clouds within meso- β - to synoptic-scale systems (200–2000 km). For example, stratiform regions in the comma head of a wintertime extratropical cyclone can be challenging due to the complex flow regimes. A possible solution is provided by Daniels et al. (2018) in which multiple GOES-16 ABI spectral channels were used to identify and track cloud, or moisture, features to estimate atmospheric motion at different levels. These ABI atmospheric motion vector winds can depict synoptic and mesoscale motions, and those associated with wintertime midlatitude cyclones (i.e., synoptic conveyor belts; Bedka et al. 2009). A need for a robust cloud tracking algorithm plays a pivotal role in understanding ABI characteristics associated with snowfall and TSI. Hanna et al. (2008) demonstrated the importance of the cloud-top T_B distributions for precipitating winter clouds, and the differences in T_B distributions of different precipitation types, in having correct measurements to facilitate a better understanding in cloud evolution prior to TSI.

The availability of high temporal, spatial, and spectral resolution geostationary satellite imagery over the United States, along with the recent development of the TDA, provides an opportunity to better understand TSI. Relating trends in T_B , reflectance values, and RGB colors to physical processes within the cloud structure increases our ability to identify important cloud and microphysical characteristics of TSSN in advanced imagery leading up to its development, and expand conceptual models of electrified snowfall events from a satellite-based framework. Henceforth, the questions this study addressed were: 1) How do cloud-top phase and ice crystal sizes change leading up to TSI? 2) What can be inferred from RGB composite imagery prior to TSI? 3) Are there similarities between cloud-top characteristics related to TSI and those associated with lightning initiation in summertime convection? Thus, the objectives of this study were to:

- 1) Build upon the work of Cordak et al. (2020) to develop a Lagrangian technique that uses ABI atmospheric motion vectors to objectively track cloud features that eventually lead to TSI;
- 2) Quantize the trends in ABI T_B , ΔT_B , reflectance, and RGB color/byte values prior to TSI and compare those distributions to Hanna et al. (2008), Harris et al. (2010), Matthee and Mecikalski (2013), and Elsenheimer and Gravelle (2019);
- 3) Relate channel and RGB trends to physical processes in tracked cloud features that produce TSSN.

A collection of electrified snowfall events was investigated, including numerous heavy-banded snowfall events associated with midlatitude cyclones. This study built upon Harkema et al. (2019, 2020) by examining their cases from January 2018 to April 2018, as well as incorporating TSSN flashes that occurred in January–April 2019 and October 2019–April 2020.

2. Datasets

a. The Geostationary Lightning Mapper

GLM detects total lightning by examining optical properties on the 777.4-nm wavelength with spatial resolution of 8 km at nadir with data components: events, groups, and flashes (Goodman et al. 2013). GLM lightning detection efficiency changes from day to night with higher probability of detection occurring at night (Bateman and Mach 2020). This study used the Level-2 GLM data that is available through National Oceanic and Atmospheric Administration's Comprehensive Large Array-Data Stewardship System (NOAA CLASS) and Amazon web Services.

b. The Advanced Baseline Imager

The ABI radiometer provides the continental United States (CONUS) with 0.5–2-km spatial resolution meteorological satellite imagery every 5 min (Schmit et al. 2017). Table 1 includes the 10 ABI spectral channels used in this study, which were chosen as they provide insight into cloud-top phase, ice particle size, cloud-top temperature, vertical extent of water vapor, and/or were useful in the construction of RGBs. The Level-2 ABI data used in this study were collected from Amazon web Service and NOAA CLASS.

1) DERIVED MOTION WINDS

Estimating atmospheric wind vectors using GOES imagery has been around since the 1970s, so the ABI derived motion winds (ABI DMW) algorithm builds upon those techniques and provides derived wind vectors every 15 min for the entire CONUS (Nieman et al. 1997; Velden et al. 1997; Bresky et al. 2012; Daniels et al. 2018, 2019). The current ABI DMW algorithm uses six different ABI channels, independently. Cloud and moisture features are tracked using three sequential scenes, or images derived from the ABI channels, and are used to estimate the cloud motion derived winds at various levels, which are consistent with the wavelength of a given satellite channel. Estimates of the wind vectors at various levels are based on how features move between scenes at each level. For example, the longwave IR band (i.e., 11.2 μm , channel 14) tracks all cloud features to estimate winds at the cloud-top level (100–950 hPa) for optically thick clouds (i.e., clouds that are thick enough so that ground-based temperatures are not observed through the clouds within a satellite pixel), with a horizontal spatial resolution of 38 km (Daniels et al. 2019). Readers can refer to Daniels et al. (2019) for a complete description of the ABI DMW algorithm. The channel 14 (11.2 μm) ABI DMW data used in this study were downloaded from NOAA CLASS.

2) MULTISPECTRAL IMAGERY

The combination of single channels and/or channel differences to form RGBs allowed for the satellite imagery to be physically interpreted, with image colors designed to help represent a specific feature or atmospheric phenomena (e.g., air masses). Lensky and Rosenfeld (2008) and EUMETSAT User Services (2009) applied a standard Eq. (1) to create a

TABLE 1. GOES-East spectral channels used in this study.

Channel (frequency)	Nickname	Primary purpose
2 (0.64 μm)	“Red” visible	Cloud optical thickness
5 (1.6 μm)	“Snow/ice” near-IR	Cloud-top phase
6 (2.2 μm)	“Cloud particle size” near-IR	Cloud-top phase/size
7 (3.9 μm)	“Shortwave window” IR	Clouds, low fog/stratus
8 (6.2 μm)	“Upper level water vapor” IR	Water vapor features
10 (7.3 μm)	“Low level water vapor” IR	Water vapor features
11 (8.4 μm)	“Cloud-top phase” IR	Cloud-top phase
12 (9.6 μm)	“Ozone” IR	Day/night dynamics
13 (10.3 μm)	“Clean” IR	Cloud-top temperatures
15 (12.3 μm)	“Dirty” IR	Clouds

24-bit composite image that was first applied to MSG satellite imagery, which has been applied to additional polar-orbiting and geostationary sensors (e.g., [Hillger et al. 2011](#); [Zavodsky et al. 2013](#); [Elmer 2015](#); [Fuell et al. 2016](#); [Elmer et al. 2016, 2019](#); [Berndt et al. 2018](#); [Miller et al. 2020](#)). The RGB images were created by combining the 8-bit red, green, and blue components for which the RGB byte values were calculated separately by using the following equation:

$$\text{byte} = 255 \times \left[\frac{(T_B, \Delta T_B, R, \Delta R) - \text{MIN}}{\text{MAX} - \text{MIN}} \right]^{1/\gamma}, \quad (1)$$

where T_B and R represent single channel brightness temperature and reflectance values, respectively. In contrast, ΔT_B and ΔR represent channel differences between T_B and reflectance values, respectively. The MIN and MAX terms represent the minimum and maximum thresholds (in K) in the specified channel or channel difference to constrain the resulting color intensities over a specified range of values. Gamma (γ) is an enhancement parameter and can be applied to adjust the spacing of the range of byte values (e.g., equally spaced, convex, or concave) to allow finer detail in the lower or upper portion of the byte range when gamma is not equal to 1. [Table 2](#) shows the mathematical properties of each RGB component used in this study.

The differential water vapor RGB ([Shimizu 2015](#); [Jedlovec et al. 2018](#); [Berndt et al. 2019](#)) can provide insights into the vertical distribution of water vapor to help diagnose the depth of moisture boundaries and associated trough/ridge patterns. The differential water vapor RGB is dominated by three colors: white, orange, and gray/blue ([Fig. 1a](#)). These colors represent thick clouds, dry regions, and moist regions with no clouds, respectively ([Shimizu 2015](#)). While the differential water vapor RGB is used to diagnose the vertical distribution of upper-level air masses, the air-mass RGB ([Lensky and Rosenfeld 2008](#); [Zavodsky et al. 2013](#); [Elmer 2015](#)) can be used to diagnostically examine upper-level air mass characteristics particularly in regions of rapid cyclogenesis ([Fig. 1b](#)). Clouds in the air-mass RGB differ in color depending on the height in the atmosphere and the characteristics of the overlaying air mass. High thick clouds are white, midlevel clouds are tan, and low-level clouds are blue or green depending on whether cold or warm air is present. Low to midlevel clouds can also appear tan or whitish red when dry air is aloft. Greens are representative of warm, moist, low-ozone air masses, and blues indicate polar, ozone-rich air masses; in contrast, warm, dry, ozone-rich air masses are represented by orange/tan colors and warm, dry air masses will appear olive/orange.

TABLE 2. Various RGB components and related information needed to calculate the appropriate R, G, and B color intensities or assigned byte values. When the RGB component is inverted, the left-hand side of Eq. (1) is subtracted from 255 and allows features to be enhanced (de-emphasized) by reversing the range of values to assign high (low) color intensity.

RGB name/component	Band(s)	Min (max) value	Gamma (γ)	Inverted
Day cloud phase distinction/R	13	219.65 (280.65)	1	Yes
Day cloud phase distinction/G	2	0 (78)	1	No
Day cloud phase distinction/B	5	1 (59)	1	No
Nighttime microphysics/R	15–13	−6.7 (2.6)	1	No
Nighttime microphysics/G	13–7	−3.1 (5.2)	1	No
Nighttime microphysics/B	13	243.6 (292.6)	1	No
Differential water vapor/R	10–8	−3 (30)	0.29	Yes
Differential water vapor/G	10	213.15 (278.15)	0.4	Yes
Differential water vapor/B	8	208.5 (243.9)	0.4	Yes
Airmass/R	8–10	−26.2 (0.6)	1	No
Airmass/G	12–13	−43.2 (6.7)	1	No
Airmass/B	8	208.5 (243.9)	1	Yes

FIG. 1. Four-panel display of (a) differential water vapor, (b) air-mass, (c) nighttime microphysics, and (d) day cloud phase distinction RGB on 11 Apr 2019. The top three panels show the developing system at 0631 UTC. In contrast, the bottom panel displays the mature system at 1856 UTC. Two different times are shown so proper interpretation of the nighttime microphysics and day cloud phase distinction RGBs are possible.

Unlike other RGBs, the nighttime microphysics (Lensky and Rosenfeld 2008; Berndt et al. 2018) can only be used to identify different cloud types at night (Fig. 1c). The reflectance component of the $3.9\text{-}\mu\text{m}$ shortwave IR channel leads to inaccurate and noisy T_B values during the daytime (Lindsey et al. 2006). Dark reds in this RGB are representative of high thick clouds, and red features with intermittent yellow features are representative of very cold clouds (i.e., $\sim -30^\circ\text{C}$) due to saturation of the $3.9\text{-}\mu\text{m}$ channel. In contrast, midlevel clouds associated with water/ice have a dull green and/or tan color (Lensky and Rosenfeld 2008). The day cloud phase distinction RGB composite (Shimizu 2015; Elsenheimer and Gravelle 2019) was used to evaluate the phase of cooling cloud-tops—during the day—and can be useful for monitoring storm growth/decay and convective storm initiation (Fig. 1d). In this RGB, light green

shading is associated with glaciating clouds, and the yellow (red/orange) colors signify thick (thin) high clouds with ice particles. Low-level clouds associated with water droplets are represented by a cyan/lavender color (Shimizu 2015).

3. Methodology

a. Cloud-top trajectories from derived motion wind

The tracking of cloud features associated with TSI was based on the native tracking capability within the ABI DMW product. Essentially, the cloud features associated with the first GLM group were tracked backward using gridded back trajectory winds derived from ABI DMW data up to 2 h prior to TSI. This process created a cloud feature trajectory, whereby the cloud evolution can be tracked x min prior to the TSI time and location. The $11.2\text{-}\mu\text{m}$ channel has the capability to estimate atmospheric motion vectors from cloud-top T_B at varying heights as it is less sensitive to errors due to water vapor absorption. The ABI DMW point data were first gridded using a Cressman interpolation within MetPy (Cressman 1959; May et al. 2020). Next, the DMW grid was reprojected into the WGS84 latitude/longitude with a spatial resolution of 10 km to approximately match the GLM resolution. Finally, individual u - and v -component grids of the wind field were extracted.

The feature tracking method involved finding the nearest gridded ABI DMW location to a TSI designated GLM group location, and converting the u and v winds from meters per second to degrees per 5 min. These converted winds were subtracted from the GLM group location (i.e., latitude and longitude) and signify the new location of the cloud feature 5 min prior. This backward trajectory process continued until the feature was tracked for 120 min. Figure 2 represents a schematic of how the backward trajectories were computed. Updrafts associated with snowfall are known to be weaker and often very slanted in the vertical. These slanted updrafts are therefore associated with slower in-cloud charging rates and with much slower/less drastic cloud evolution and movement (see supplementary video for an example). As a result, tracking cloud features in satellite observations only 1 h prior to TSI, like the methods found in Harris et al. (2010) and Matthee and Mecikalski (2013), may be too short to quantify the cloud-top characteristics associated with electrified snowfall.

b. Assigning T_B , reflectance, and RGB byte values to TSI trajectories

ABI data were also transformed to the same WGS84 latitude/longitude coordinate system with a 2-km spatial resolution. The T_B and reflectance values were extracted at 5-min intervals from each of the grids using a 10-km search radius around the DMW-derived trajectory center at each time. The coldest T_B (mean of the lowest 25th percentile of pixels in the individual search radius) and highest reflectance (mean

FIG. 2. Schematic of how the TSI trajectories are calculated from the gridded ABI DMW and the initial GLM group location.

above the 75th percentile) from the 20-km window were assigned to capture the timing of glaciation. Individual RGB byte values were then calculated using these values in Eq. (1). Additionally, the 3.9- μm channel was separated into the reflectance and emitted components, during the day, following the methodology described in Setvák and Doswell (1991) and Lindsey et al. (2006).

Assigning the correct T_B , reflectance, and RGB byte values, and properly interpreting those values are two separate matters. For example, the day cloud phase distinction RGB and visible bands are only physically useful during the day. In contrast, the nighttime microphysics green component contains the 3.9- μm ABI channel; thus, the nighttime microphysics RGB was designed specifically for nighttime use (Lensky and Rosenfeld 2008; Berndt et al. 2018). Therefore, the ability to differentiate between day and night within a satellite image is paramount, and solar zenith angle plays a role in this differentiation. A solar angle of 90° represents sunrise/sunset and can differentiate day and night. However, the quality of the visible band values slightly degrades before/after sunset and sunrise and may not be defined as the true reflectance of the cloud tops (Lindsey et al. 2006; Miller et al. 2020). In this study, daytime was representative of locations where the solar angle is $\leq 80^\circ$; night was assigned to locations where the solar angle is $\geq 92^\circ$. When comparing the solar angle to reflectance values, it was found that reflectance was at or near zero for solar angles greater than 92°, and the T_B difference between 3.9 μm (uncorrected) and 10.3 μm sharply decreased between solar angles of 80° and 92°. Last, outliers in T_B , reflectance, and RGB byte values were neglected for any statistical analysis.

4. Results

a. Qualitative accuracy of cloud tracking method and TSSN flash classification

At 5-min intervals, the tracked cloud features were assigned T_B and reflectance values at each location of the TSI trajectory. These assigned values were then used to create RGB byte values, which were stacked together to produce a color. One such TSI trajectory traversed from Iowa into Wisconsin (supplementary video). The tracked cloud feature within the yellow oval demonstrates the effectiveness of the cloud-tracking technique. At the bottom of the main plot in the supplementary video is a subplot that demonstrates how the airmass RGB color changes with time as the cloud feature is tracked. A red box highlights the specific airmass RGB color that corresponds to the location and time that is highlighted in the top plot of the video. Specifically for this TSI trajectory, the tracked cloud feature in the airmass RGB transitioned from a green to olive green, then to a tan color. In the airmass RGB, the olive green color of the low- to midlevel clouds is an indicator of the temperature/moisture characteristics of the air mass above the cloud feature, and midlevel clouds are typically a tan color. At 120 min prior to TSI, the cloud feature was associated with a warm/moist upper-level air mass (i.e., green hues). In contrast, at the time of TSI, the cloud feature was associated with a different (i.e., drier) upper-level air mass (i.e., orange hues). This physically represents the cloud feature moving from a warm upper-level air mass with some moisture and then evolving into a midlevel cloud associated with a drier upper-level air mass prior to TSI.

The TDA identified 22 067 TSSN flashes and represents a nearly tenfold increase over the number of TSSN flashes analyzed in Harkema et al. (2019, 2020). The tenfold increase in the number of TSSN flashes was a result of analyzing TSSN from additional seasons (e.g., 2019 and 2020) and the improved characterization of the rain-to-snow transition region. Each flash and associated TSI trajectory was classified as day (solar angle $< 80^\circ$; 5197 flashes) or night (solar angle $> 92^\circ$; 11 447 flashes). The higher percentage of nighttime flashes could be due to better detection efficiency by the GLM overnight (e.g., Bateman and Mach 2020) and/or the decreased hours of daylight in the Northern Hemisphere winter.

b. Trends in ABI channels prior to TSI

Violin plots for channels 8 (6.2 μm), 10 (7.3 μm), and 12 (9.6 μm) were examined to diagnose environmental characteristics that impact storm development (Fig. 3). For example, the T_B distributions become colder—prior to TSI—for the upper-level water vapor (i.e., 6.2 μm , Fig. 3a) and low-level water vapor (i.e., 7.3 μm , Fig. 3b) channels. The decreases in T_B in these channels signify the cloud structure deepening prior to TSI. The multimodal distribution in the upper-level water vapor channel, at 120 min prior to TSI, suggests four distinct T_B regions where these trajectories originated, which indicates various degrees of moisture and cloud development in the mid and upper-levels. The 9.6- μm ozone channel was

FIG. 3. Violin plots related to the T_B distributions of the Advanced Baseline Imager water vapor and infrared channels used in this study: (a) channel 8 ($6.2\ \mu\text{m}$), (b) channel 10 ($7.3\ \mu\text{m}$), (c) channel 12 ($9.6\ \mu\text{m}$), (d) channel 11 ($8.4\ \mu\text{m}$), (e) channel 13 ($10.3\ \mu\text{m}$), and (f) channel 15 ($12.3\ \mu\text{m}$) leading up to thundersnow initiation.

associated with a weak bimodal distribution and transitioned to a gamma distribution tailing off toward warmer T_B at the time of TSI (Fig. 3c). The shift in the distribution represents the deepening of clouds prior to TSI. The changes in the bimodal and multimodal IR and water vapor T_B distribution suggest the TSI trajectories originated in either: 1) regions of preexisting convection, or 2) outside deep convection and associated with clear air conditions. See Table 3 for the median values for the various violin plot distributions mentioned above and hereafter.

The 8.4- (Fig. 3d), 10.3- (Fig. 3e), and 12.3- μm (Fig. 3f) channels are nearly identical in shape; this is not unexpected because these channels are within the IR atmospheric window. Slight variations between these distributions are caused by differences in water vapor absorption at each frequency. Related to the clouds being analyzed for this study, the IR cloud-top phase channel (i.e., $8.4\ \mu\text{m}$) has two peaks in the T_B distribution 2 h prior to TSI (Fig. 3d). Similarly, the clean (i.e., $10.3\ \mu\text{m}$) and dirty (i.e., $12.3\ \mu\text{m}$) atmospheric window channels (Figs. 3e,f) have peaks at the same T_B , 2 h prior to TSI, with each peak corresponding to clear air conditions or shallow clouds, or optically thick clouds, respectively. Additionally, all three of the panels indicate that the distributions develop more prominent peak(s) at colder T_B , suggesting that

TSI was more preferential to cloud features associated with cloud T_B values between 200 and 230 K. Fluctuations in the T_B values for all three channels over time provide insight into the changes in the cloud structure. The 60-min differences in the $10.3\text{-}\mu\text{m}$ cloud-top T_B showed that T_B decreased by a median value of $12.2\ \text{K}$ prior to TSI. During this time period, 23.8% of TSI trajectories were associated with increases in T_B prior to TSI (not shown). Over the course of 2 h, 85.6% of TSSN flashes observed by GLM were associated with a decrease in $10.3\text{-}\mu\text{m}$ T_B , and the range in cloud-top T_B differences [e.g., max increase (decrease): $42.4\ \text{K}$ ($70.2\ \text{K}$)] demonstrates that TSSN can occur in a variety of cloud features. Increases of $10.3\text{-}\mu\text{m}$ T_B prior to TSI were likely caused by potential errors in tracking the cloud features back in time, particularly, if the trajectory crossed a boundary with low-level clouds and upper-level clouds moving in different directions. In comparison, Harris et al. (2010) found that GOES satellite $10.7\text{-}\mu\text{m}$ 15-min trends in T_B decreased by a median of approximately $16\ \text{K}$ prior to lightning initiation in summertime convection. Thus, the changes in T_B were small compared to those associated with intense summertime convective clouds, which likely were caused by slantwise ascent of air as common within winter systems.

TABLE 3. The 15-min intervals of the median T_B (K) or reflectance value (%) for various ABI bands leading up to TSI (i.e., $t - 0$). Bands 2–6 are all reflectance values (%), and bands 8–15 are T_B (K). “Band 7—refl” is in percent, while “band 7—emit” and band 7 are T_B in kelvin.

ABI band	Time	$t - 120$	$t - 105$	$t - 90$	$t - 75$	$t - 60$	$t - 45$	$t - 30$	$t - 15$	$t - 0$
2 (0.64 μm)	Day	54.2	57.7	60.5	63.3	65.6	68.3	71.0	71.8	72.6
5 (1.6 μm)	Day	27.3	28.4	29.0	29.0	28.0	26.5	25.3	24.0	23.5
6 (2.2 μm)	Day	25.8	26.4	27.3	28.0	28.2	28.5	28.7	29.1	29.2
7 (3.9 μm)—refl	Day	4.1	3.5	3.1	2.7	2.4	2.2	2.2	2.2	2.3
7 (3.9 μm)—emit	Day	248.3	247.8	246.1	234.0	238.2	234.2	230.7	228.3	225.4
7 (3.9 μm)	Night	254.7	252.3	248.9	245.1	239.7	233.8	229.9	226.8	222.7
8 (6.2 μm)	Day/night	232.3	231.4	230.3	228.7	227.1	225.6	224.2	222.7	220.8
10 (7.3 μm)	Day/night	243.4	241.3	239.1	236.0	232.3	229.4	227.0	225.0	222.1
11 (8.4 μm)	Day/night	250.7	248.6	245.7	241.2	235.9	231.8	228.7	226.2	222.8
12 (9.6 μm)	Day/night	236.0	235.0	233.5	231.5	229.3	227.8	226.6	225.6	224.2
13 (10.3 μm)	Day/night	252.2	249.8	246.9	242.2	236.7	232.5	229.4	226.8	223.4
15 (12.3 μm)	Day/night	250.6	248.1	244.7	240.0	234.6	230.8	228.0	225.5	222.2

At night, the shortwave IR channel (3.9 μm)—as well as the emitted daytime component—can be interpreted similarly as the other IR channels. In contrast, the reflected daytime component of the shortwave IR channel provides insight into cloud-top microphysics (e.g., glaciation). As a result, the nighttime shortwave IR and emitted daytime T_B values are distributed similarly to the clear window channel (i.e., 10.3 μm) because they show the same information (not shown). The median 3.9- μm reflectance values decrease from 4.0% to 2.3% 2 h prior to TSI (Table 3). These values are similar to those in convective storms (Setvák et al. 2003). The decrease in the variability in reflectance (see Fig. 4) is likely, in part, a result of water drops glaciating into ice crystals, as water clouds are more reflective at 3.9 μm compared to ice clouds (Turk et al. 1998). Several TSI trajectory 3.9- μm reflectance values were plotted within an hour of TSI and 5-min differences were found to be as much as $\pm 3\%$ (not shown). Furthermore, GOES-East ABI cloud optical depth data were plotted for a single timeframe at 1501 UTC 11 April 2019; all 59 TSI flashes—within a 5-min period prior to the ABI data—

occurred in or around optically thick clouds (i.e., cloud optical depth ≥ 38 ; not shown). From this result, it was assumed that all TSI flashes within this study occurred within optically thick clouds. Therefore, assuming changes in solar angle were small and clouds were optically thick, any decrease (increase) in 3.9- μm reflectance—after glaciation has occurred—may be associated with ice particles becoming larger (smaller) over time (Lindsey et al. 2006). These 3.9- μm reflectance characteristics suggest that: 1) cloud-top phase transitioned from water to ice, and 2) ice crystals were colliding with each other prior to TSI and/or there is evidence that large ice particle sedimentation (i.e., particle settling) was occurring at cloud top.

At 120 min prior to TSI, the visible and near IR channels had a distinct bimodal distribution (Fig. 5). As TSI approaches, the bimodal distribution becomes more compact for the 0.64- (Fig. 5a) and 2.2- μm channels (Fig. 5c), and Gaussian for the 1.6- μm channel (Fig. 5b). In the 0.64- μm channel (Fig. 5a), the daytime cloud features’ median reflectance increased from 54.2% to 72.6% over the course of 2 h (Table 3). This increase in reflectance physically represents the cloud features becoming more optically thick over time, which implies increasing trends of stronger updrafts reaching to cloud top, and/or thickening cloud structures in general that are increasingly supportive of lightning production. The range in the snow/ice band (i.e., 1.61 μm) reflectance values decreased (Fig. 5b) for the 2-h duration leading up to initiation with the median reflectance values remaining between 23% and 27% (Table 3). Like the reflected component of the 3.9- μm channel, decreasing reflectance values in the 1.61- μm channel represent glaciation. Finally, the median reflectance values for the cloud particle size band (i.e., 2.24 μm , Fig. 5c) slightly increased from 25.8% to 29.2%, suggesting cloud particles in the median cloud feature associated with TSI became slightly smaller throughout the duration of the trajectory (Table 3).

c. Trends in ABI RGB prior to TSI

The trends in median RGB byte values can be used to better understand how changes correspond to the microphysical

FIG. 4. Violin plot related to daytime reflectance (%) of the Advanced Baseline Imager shortwave IR channel (3.9 μm) leading up to thundersnow initiation. Note that the y axis is logarithmic to better show the reflectance distributions.

FIG. 5. Violin plots related to the reflectance (%) distributions of the Advanced Baseline Imager visible and near-infrared channels used in this study: (a) channel 2 ($0.64\ \mu\text{m}$), (b) channel 5 ($1.6\ \mu\text{m}$), and (c) channel 6 ($2.2\ \mu\text{m}$) leading up to thundersnow initiation.

properties or air mass characteristics (Table 4). These trends can be used to imply how the red, green, and blue components of the satellite imagery change over time, and how these changes relate to the microphysical characteristics of the clouds or to the air mass in the scene (depending on the specific RGB). The nighttime microphysics and day cloud phase distinction RGB provide insight into cloud microphysics, and over the course of 2 h, the median red and blue components for the nighttime microphysics RGB (nighttime only) changed little over time. As a result, the change (i.e., decrease) in the

green component (i.e., $10.3\text{--}3.9\ \mu\text{m}$) of the nighttime microphysics was the most influential for any color change. The decrease in this byte value physically represents liquid water clouds glaciating into ice crystal-dominated clouds. The median day cloud phase distinction (daytime only) red and green byte values increased whereas the median blue byte value decreased slightly (Table 4). This means that the largest contributions to the color changes in the day cloud phase distinction RGB were a result of cloud tops becoming cooler (i.e., red component; $10.3\ \mu\text{m}$), and becoming deeper and more optically thick (i.e., green component; $0.64\ \mu\text{m}$).

In contrast, the differential water vapor and air mass RGB provide insight into the vertical and horizontal extent of water vapor associated with different air masses, respectively, and do not provide information on cloud microphysics. However, relative cloud height can be inferred from the air mass RGB. As the time of TSI approaches, the median red, green, and blue byte values increase for the tracked trajectory for both the air mass and differential water vapor RGB (Table 4). The increases in the median red (i.e., $6.2\text{--}7.3\ \mu\text{m}$) and green (i.e., $9.6\text{--}10.3\ \mu\text{m}$) byte values for the air mass RGB indicate that the cloud features were becoming higher and optically thicker. The increases in the blue component (i.e., $6.2\ \mu\text{m}$) indicate that the cloud-top T_B were cooling prior to TSI. The trends in the differential water vapor byte values confirm the results from the air mass RGB. These median byte values for the nighttime microphysics, day cloud phase distinction, air mass, and differential water vapor RGBs likely do not correlate with the same trajectory. Therefore, these median RGB byte values cannot be combined to produce a meaningful color related to the physical interpretation of the RGB.

For each trajectory, the combination of the RGB byte values into a color was done to provide greater insight into the changes in the cloud structure that contribute to TSI. Figures 6 and 7 were produced to better visualize the RGB color trends for all TSI trajectories that did not contain any missing RGB component(s) for night and day, respectively. These RGB color-transition plots indicate the color of a specified RGB at a specific time (x axis) for a specific TSI trajectory (y axis). Moving from left (i.e., 120 min prior to TSI) to right (i.e., time of TSI)—while maintaining a constant y value (i.e., a single back-tracked trajectory)—shows the RGB color transition

TABLE 4. The 15-min intervals of the median byte values for various RGB components leading up to TSI (i.e., $t - 0$).

RGB name/component	$t - 120$	$t - 105$	$t - 90$	$t - 75$	$t - 60$	$t - 45$	$t - 30$	$t - 15$	$t - 0$
Day cloud phase distinction/R	130	133	141	155	177	193	208	218	230
Day cloud phase distinction/G	174	187	198	208	214	223	232	235	237
Day cloud phase distinction/B	115	120	122	123	118	112	106	101	98
Nighttime microphysics/R	153	154	156	157	157	157	158	158	159
Nighttime microphysics/G	102	96	87	82	78	78	76	79	84
Nighttime microphysics/B	3	0	0	0	0	0	0	0	0
Differential water vapor/R	248	249	252	253	254	254	254	254	254
Differential water vapor/G	228	231	237	242	247	249	251	252	253
Differential water vapor/B	181	187	195	204	213	221	228	234	238
Airmass/R	155	162	176	191	207	219	228	233	240
Airmass/G	140	145	155	169	184	196	205	213	225
Airmass/B	84	90	98	110	121	131	141	151	168

FIG. 6. Three-panel display of the changes in RGB color leading up to TSI during the night, as seen from the (a) differential water vapor, (b) airmass, and (c) nighttime microphysics RGBs. The TSI trajectories in the panels are sorted by differential water vapor color at 120 min prior to TSI, and the sorting order is applied to the other two RGBs. These color transition plots represent how cloud features transition from different environments and moisten over time, and how cloud top particle sizes and phase change leading up to TSI.

that a TSI trajectory experienced while being tracked. In other words, the color changes in the plots show the progression of changing cloud characteristics in time up to TSI. Specifically, these color transitions represent the development of updrafts that lead to cloud-top glaciation, the lofting of ice particles enough to produce in-cloud charging that eventually leads to TSI. Both color-transition plots (Figs. 6 and 7) were sorted by the differential water vapor RGB color 2 h prior to TSI (i.e., leftmost side of the plot), and that sort order was applied to the other two RGBs. Therefore, the RGB color-transition plots allow for visual inspection of the TSI trajectories and the relation of RGB color changes to physical cloud processes that lead to TSI. See the supplementary video for a detailed example on how the color transition plot are properly read and how they relate to the TSI trajectory.

At night, the differential water vapor RGB trajectories (Fig. 6a) consist of two main colors 2 h prior to TSI, orange/tan and white, and represent regions of low-level clouds associated with a dry layer in the mid-to-upper atmosphere, and thick-high clouds, respectively. At 60 min prior to TSI, tints of gold/orange became pronounced in the differential water vapor RGB, representing cloud development in the midlevels with a dry upper level (Berndt et al. 2019). At time of TSI, thick-high clouds (i.e., white/grays) were present for the majority of the TSI trajectories. The coloration in the airmass color-transition plot (Fig. 6b) is dominated by tan/brown hues with light browns becoming more prevalent at time periods closer to TSI. Like the differential water vapor color transitions, the airmass colorations suggest that the trajectories

FIG. 7. Three-panel display of the changes in RGB color leading up to TSI during the day, as seen from the (a) differential water vapor, (b) airmass, and (c) day cloud phase distinction RGBs. The TSI trajectories in the panels are sorted by differential water vapor color at 120 min prior to TSI, and the sorting order is applied to the other two RGBs. These color transition plots represent how cloud features transition from different environments and moisten over time, and how cloud top particle sizes and phase change leading up to TSI.

were more likely to originate in low-level clouds. The lack of pure white colors near the time of TSI also indicates that TSI was more likely to occur in midlevel cloud features associated with drier upper-level air masses. In other words, TSI was not preferential to the tallest cloud features because the airmass RGB coloring was not a bright white color (Fig. 6b). The increased extent of the tan/brown colors are indicative of mid-level clouds associated with upper-level warm air masses with less moisture. As for the nighttime microphysics color transition plot (Fig. 6c), the main RGB colors 2 h prior to TSI are sea-green (low- to midlevel water cloud), red (high-thick cloud), yellow (very cold optically thick cloud), and purple (high, very thin, ice cloud). As time of TSI approaches (i.e., approaching 0 min prior to TSI), these colors quickly transition into two predominate colors, yellow and red both with orange hues. The lack of pure yellows and dark reds suggests that the clouds became glaciated, if they were not already, and that they were not the coldest and thickest clouds possible.

During the day and 2 h prior to TSI, the differential water vapor RGB trajectories consist of mostly white, tan colors, and, to lesser extents, bright orange colors (Fig. 7a). Interestingly, there was a greater extent of gold/orange hues as the time of TSI approached for the differential water vapor RGB during the day, suggesting that these trajectories were associated with midlevel clouds associated with a slightly less dry upper-level compared to those during the night. This was also seen in the airmass color-transition plot (Fig. 7b) where the coloring appears to favor a lighter brown hue. Even during

the day, TSI does not appear to occur in the coldest and thickest clouds as represented by the differential water vapor and airmass RGBs. At 2 h prior to TSI, the day cloud phase distinction RGB colors were predominately red/orange, yellow, and cyan (Fig. 7c). Cyan and orange/red colors represent low-level clouds with water droplets and thin high-level clouds with ice particles, respectively. At time of TSI, the day cloud phase distinction RGB colors were primarily hues of yellow/green and, to lesser extents, cyan and orange/red (Fig. 7c). The yellow/green colors represent regions of mid-to-high level clouds with ice particles and/or clouds associated with glaciation. From these results, it is clear that many back-tracked cloud features—that eventually lead to TSI—began as low-level clouds with water droplets, which eventually became glaciated prior to TSI. This cloud-top glaciation therefore is indicative of higher in-cloud supercooled water and ice concentrations, which lead to in-cloud charging and eventual lightning.

5. Discussion

Over 76% of TSSN flashes in this study were associated with decreases in $10.3\text{-}\mu\text{m } T_B$ 1 h prior to TSI, which exceeded the $\sim 50\%$ value found in Cordak et al. (2020). The primary reason for this discrepancy was that this study used a Lagrangian tracking method compared to the Eulerian method in Cordak et al. (2020). Even though it was snowing at the surface, the $10.3\text{-}\mu\text{m } T_B$ distributions shown within this study were more comparable to the rain distributions than to the snow distributions in Hanna et al. (2008). At 60 min prior to TSI, the two peaks in the bimodal distribution for the $10.3\text{-}\mu\text{m } T_B$ (Fig. 3e) closely match that of the light rain distribution from Hanna et al. (2008) (Fig. 2b in Hanna et al.). In comparison, at the time of TSI, the $10.3\text{-}\mu\text{m } T_B$ distribution appear to be qualitatively similar to the moderate rain distribution from Hanna et al. (2008) (Fig. 2c in Hanna et al.). These similarities show the typical TSSN environment is convectively driven and likely contains more supercooled water than typical snow environments without lightning (Schultz 1999; Market et al. 2006).

At the time of TSI, the median T_B for the $10.3\text{-}\mu\text{m}$ channel was 222.2 K, which is comparable to the median value of 223.7 K from Matthee and Mecikalski (2013); furthermore, the nonoutlier data also ranged from approximately 260 K to 200 K and had a similar distribution shape to the $10.3\text{-}\mu\text{m } T_B$ results in Thiel et al. (2020). These results again show that T_B values for convective clouds in winter storms are distributed similarly to nonwinter convective clouds. This result also aligns with the conclusions of Takahashi et al. (1999) regarding charge centers in winter clouds. Takahashi et al. (1999) noted that while charge centers are at lower heights during winter, they are typically at the same temperature as warmer convection. At 1 h prior to TSI, the median $10.3\text{-}\mu\text{m } T_B$ decreased by 12.2 K, which is less than the 15 min $10.7\text{-}\mu\text{m } T_B$ decrease of 16 K for summertime convection as noted in Harris et al. (2010). This difference is likely caused by the fact that TSSN cases are more likely to be associated with

slantwise ascent and lower instability; thus, the changes in cloud-top $10.3\text{-}\mu\text{m } T_B$ values reflect larger meso- and synoptic-scale cloud feature evolution in the TSSN environment rather than individual updrafts associated with TSSN events.

TSI events have similar day cloud phase distinction RGB color trends (i.e., cyan to yellow) compared to cloud features associated with lightning initiation in summertime convection (Fig. 7c; Elseneheimer and Gravelle 2019). The changes in cloud-top phase and optical thickness that lead up to TSI should be studied for summertime lightning flashes at night by using the nighttime microphysics RGB. Additionally, coupling this information with electric field measurements or modeling would be helpful in further understanding the electrification process in winter storms. The light brown colors and gold/orange hues in the airmass and differential water vapor RGB, respectively, indicate the development of middle-level clouds associated with dry upper-layer air masses prior to TSI. Dry air aloft can aid in the continued development of the mixed-phase region in clouds via dry air entrainment, latent heating, and increased updraft velocities. In turn, these processes cause turbulent in-cloud motions that can cause water and ice hydrometeors to collide with each other. These collisions within the mixed-phase region enhance the electric field through noninductive charging and—if charge separation occurs—increases the potential for lightning initiation (Reynolds et al. 1957; Takahashi 1978; Saunders et al. 2006). In the context of a mature extratropical cyclone (see Fig. 1d), these colorations in the airmass and differential water vapor RGBs were likely associated with descending stratospheric air (i.e., the dry conveyor belt). In part, the dry conveyor belt plays a role in the development of heavy-banded snowfall (Nicosia and Grumm 1999; Moore et al. 2005). Thus, the color transitions in the airmass and differential water vapor RGBs suggest that the dry conveyor belt may influence the environment that is conducive for TSI. Furthermore, the synergistic interactions of the dry, warm, and cold conveyor belts likely provide the necessary conditions ripe for instability and lift. However, more analysis would be needed to fully understand how these interactions directly influence the atmosphere to produce an environment that can support TSI.

The noninductive charging theory (Reynolds et al. 1957; Takahashi 1978; Saunders et al. 2006) states that ice particles, supercooled liquid water, and graupel need to be present for lightning initiation. The normalization of the $1.61\text{-}\mu\text{m}$ reflectance values peaking around 0.25 suggests that cloud features associated with TSI contain both ice crystals and (supercooled liquid) water droplets with ice particles having a dominant presence (Fig. 5b). Assuming clouds were optically thick near and at time of TSI and changes in solar angle were small, the variability in the $3.9\text{-}\mu\text{m}$ reflectance values over 5-min intervals indicates that some TSI trajectories were associated with ice crystal collisions which likely resulted in enhancements in the electric field. Therefore, these results suggest that the TSI trajectories were associated with ice crystals collisions and the coexistence of (supercooled liquid) water droplets and ice crystals at cloud top. However, this study did not provide conclusive evidence of graupel (or lack thereof) and associated charge separation within the cloud feature beyond the fact

that TSSN was observed. In situ observation would be needed to determine locations of graupel and the presence of charge separation through electric field measurements in/above winter clouds (e.g., [Schultz et al. 2021](#)).

6. Conclusions

Studies have examined TSSN to aid in forecasting heavy-banded snowfall and understanding flash characteristics, but fewer studies have investigated the use of satellite observations to understand TSI. In this study, TSI was defined as the first GLM group in a GLM flash that was identified by a TDA ([Harkema et al. 2019, 2020](#)). ABI DMW L2 data were gridded and interpolated in order to develop a Lagrangian cloud feature tracking method to track clouds associated with TSI. A total of 10 ABI bands ([Table 1](#)) and four RGBs (i.e., differential water vapor, airmass, nighttime microphysics, and day cloud phase distinction; [Table 2](#)) were examined to understand the characteristics and microphysics associated with TSI and examine how TSSN may be connected with the air masses and synoptic-scale motions around midlatitude cyclones (i.e., conveyor belts). ABI T_B , reflectance, and RGB byte values and colors were assigned to the locations within the TSI trajectories.

The goal of this study was to answer questions about changes in cloud-top phase and ice particle sizes prior to TSI; how RGB satellite imagery might be useful to understand TSI; and potential similarities between cloud-top characteristic of TSI and lighting initiation in summertime convection. Answering the questions posed at the end of the introduction:

- 1) Values from the TSI trajectories showed water droplets glaciated into ice particles. Prior to TSI, some trajectories had large variability in 3.9- μm reflectance over short time periods (i.e., 5 min) and indicated that ice collisions were likely occurring at cloud top. IR channels depicted cloud features with vertical growth and decreased T_B .
- 2) RGB color changes from day cloud phase distinction and nighttime microphysics indicated the potential in using patterns such as cyan to yellow (day cloud) to predict TSSN. RGB patterns from some of the differential water vapor and airmass trajectories showed the cloud features were associated with upper-level dryness (e.g., gold/orange hues in differential water) prior to TSI.
- 3) Similarities between the TSSN T_B distributions and rain T_B distributions in [Hanna et al. \(2008\)](#) and non-TSSN lightning T_B distributions in [Harris et al. \(2010\)](#) and [Matthee and Mecikalski \(2013\)](#) suggested that the TSSN environment was fairly convective, and hence, cloud features associated with TSSN may contain more super-cooled water than traditionally stratiform snow events.

Finally, this study provided evidence that the development of TSSN may be connected to the dry conveyor belt of mature midlatitude extratropical cyclones by finding that some tracked cloud features were associated with dry upper-level air masses—prior to TSI—as represented by gold/orange color hues in the differential water vapor RGB. The

interactions between the dry, warm, and cold conveyor belts likely lead to the environmental conditions necessary for TSI. But the analysis between these interactions is well beyond the scope of this study. Overall, understanding the changes in cloud-top characteristics prior to TSI provides insight into cloud electrification and can be useful for nowcasting potential changes in surface conditions.

Acknowledgments. This work is part of a larger investigation of electrified snowfall that is directly supported through NASA's Future Investigators in NASA Earth and Space Science and Technology (FINESST) program 80NSSC20K1623. Additionally, this work was also supported by Dr. Tsengdar Lee of NASA's Research and Analysis Program, Weather and Atmospheric Dynamics Focus Area, as part of the Short-term Prediction Research and Transition (SPoRT) Center at Marshall Space Flight Center and supported through the NASA-UAH Cooperative Agreement NNM11AA01A. The authors thank the three anonymous reviewers for providing valuable insight and suggestions; especially, with regards to the RGB analysis.

REFERENCES

- Adler, R. F., and D. D. Fenn, 1979: Thunderstorm intensity as determined from satellite data. *J. Appl. Meteor.*, **18**, 502–517, [https://doi.org/10.1175/1520-0450\(1979\)018<0502:TIADFS>2.0.CO;2](https://doi.org/10.1175/1520-0450(1979)018<0502:TIADFS>2.0.CO;2).
- Bateman, M., and D. Mach, 2020: Preliminary detection efficiency and false alarm rate assessment of the Geostationary Lightning Mapper on the *GOES-16* satellite. *J. Appl. Remote Sens.*, **14**, 032406, <https://doi.org/10.1117/1.JRS.14.032406>.
- Bedka, K. M., C. S. Velden, R. A. Petersen, W. F. Feltz, and J. R. Mecikalski, 2009: Comparisons of satellite-derived atmospheric motion vectors, rawinsondes, and NOAA wind profiler observations. *J. Appl. Meteor. Climatol.*, **48**, 1542–1561, <https://doi.org/10.1175/2009JAMC1867.1>.
- Berndt, E. B., N. Elmer, L. Schultz, and A. Molthan, 2018: A methodology to determine recipe adjustments for multispectral composites derived from next-generation advanced satellite imagers. *J. Atmos. Oceanic Technol.*, **35**, 643–664, <https://doi.org/10.1175/JTECH-D-17-0047.1>.
- , G. Jedlovec, K. K. Fuell, F. J. LaFontaine, and N. J. Elmer, 2019: Analysis and applications of water vapor–derived multispectral composites for geostationary satellites. *15th Annual Symp. on New Generation Operational Environmental Satellite Systems*, Phoenix, AZ, Amer. Meteor. Soc., 227, <https://ams.confex.com/ams/2019Annual/webprogram/Paper350911.html>.
- Bresky, W. C., J. M. Daniels, A. A. Bailey, and S. T. Wanzong, 2012: New methods toward minimizing the slow speed bias associated with atmospheric motion vectors. *J. Appl. Meteor. Climatol.*, **51**, 2137–2151, <https://doi.org/10.1175/JAMC-D-11-0234.1>.
- Cintineo, J. L., M. J. Pavolonis, J. M. Sieglaff, L. Cronic, and J. Brunner, 2020: NOAA ProbSevere v2.0—ProbHail, ProbWind, and ProbTor. *Wea. Forecasting*, **35**, 1523–1543, <https://doi.org/10.1175/WAF-D-19-0242.1>.
- Cordak, A., E. B. Berndt, and S. S. Harkema, 2020: Assessing cloud-top brightness temperatures and cloud types to identify the precursors to thundersnow development. *19th Annual*

- Student Conf.*, Boston, MA, Amer. Meteor. Soc., S205, <https://ams.confex.com/ams/2020Annual/meetingapp.cgi/Paper/371641>.
- Cressman, G. P., 1959: An operational objective analysis system. *Mon. Wea. Rev.*, **87**, 367–374, [https://doi.org/10.1175/1520-0493\(1959\)087<0367:AOOAS>2.0.CO;2](https://doi.org/10.1175/1520-0493(1959)087<0367:AOOAS>2.0.CO;2).
- Crowe, C., P. Market, B. Pettegrew, C. Melick, and J. Podzimek, 2006: An investigation of thundersnow and deep snow accumulations. *Geophys. Res. Lett.*, **33**, L24812, <https://doi.org/10.1029/2006GL028214>.
- Daniels, J. M., W. C. Bresky, A. Bailey, A. Allegrino, S. Wanzong, and C. S. Velden, 2018: Introducing atmospheric motion vectors derived from the *GOES-16* Advanced Baseline Imager (ABI). *14th Annual Symp. on New Generation Operational Environmental Satellite Systems*, Austin, TX, Amer. Meteor. Soc., 11A.3, <https://ams.confex.com/ams/98Annual/webprogram/Paper331856.html>.
- , —, S. Wanzong, C. Velden, and H. Berger, 2019: GOES-R Advanced Baseline Imager (ABI) algorithm theoretical basis document for derived motion winds. NOAA/NESDIS, 103 pp, https://www.star.nesdis.noaa.gov/goesr/documents/ATBDs/Baseline/ATBD_GOES-R_Winds_v3.1_Feb2019.pdf.
- Elmer, N. J., 2015: Limb Correction of individual infrared channels for the improved interpretation of RGB composites. Dept. of Atmospheric Science, University of Alabama in Huntsville, 90 pp.
- , E. Berndt, and G. J. Jedlovec, 2016: Limb correction of MODIS and VIIRS infrared channels for the improved interpretation of RGB composites. *J. Atmos. Oceanic Technol.*, **33**, 1073–1087, <https://doi.org/10.1175/JTECH-D-15-0245.1>.
- , —, and K. Fuell, 2019: Limb correction of geostationary infrared imagery in clear and cloudy regions to improve interpretation of RGB composites for real-time applications. *J. Atmos. Oceanic Technol.*, **36**, 1675–1690, <https://doi.org/10.1175/JTECH-D-18-0206.1>.
- Elsenheimer, C. B., and C. M. Gravelle, 2019: Introducing lightning threat messaging using the *GOES-16* day cloud phase distinction RGB composite. *Wea. Forecasting*, **34**, 1587–1600, <https://doi.org/10.1175/WAF-D-19-0049.1>.
- EUMETSAT User Services, 2009: Best practices for RGB compositing of multi-spectral imagery. European Organisation for the Exploitation of Meteorological Satellites, 8 pp., http://oiswww.eumetsat.int/idds/html/doc/best_practices.pdf.
- Fuell, K. K., B. J. Guyer, D. Kann, A. L. Molthan, and N. Elmer, 2016: Next generation satellite RGB dust imagery leads to operational changes at NWS Albuquerque. *J. Oper. Meteor.*, **4**, 75–91, <https://doi.org/10.15191/nwajom.2016.0406>.
- Goodman, S. J., and Coauthors, 2013: The GOES-R Geostationary Lightning Mapper (GLM). *Atmos. Res.*, **125–126**, 34–49, <https://doi.org/10.1016/j.atmosres.2013.01.006>.
- Hanna, J. W., D. M. Schultz, and A. R. Irving, 2008: Cloud-top temperatures for precipitating winter clouds. *J. Appl. Meteor. Climatol.*, **47**, 351–359, <https://doi.org/10.1175/2007JAMC1549.1>.
- Harkema, S. S., C. J. Schultz, E. B. Berndt, and P. M. Bitzer, 2019: Geostationary Lightning Mapper flash characteristics of electrified snowfall events. *Wea. Forecasting*, **34**, 1571–1585, <https://doi.org/10.1175/WAF-D-19-0082.1>.
- , E. B. Berndt, and C. J. Schultz, 2020: Characterization of snowfall rates, totals, and snow-to-liquid ratios in electrified snowfall events identified by the Geostationary Lightning Mapper. *Wea. Forecasting*, **35**, 673–689, <https://doi.org/10.1175/WAF-D-19-0126.1>.
- Harris, R. J., J. R. Mecikalski, W. M. MacKenzie, P. A. Durkee, and K. E. Nielsen, 2010: The definition of GOES infrared lightning initiation interest fields. *J. Appl. Meteor. Climatol.*, **49**, 2527–2543, <https://doi.org/10.1175/2010JAMC2575.1>.
- Hillger, D. W., L. Grasso, S. Miller, R. Brummer, and R. DeMaria, 2011: Synthetic advanced baseline imager true-color imagery. *J. Appl. Remote Sens.*, **5**, 053520, <https://doi.org/10.1117/1.3576112>.
- Jayarante, E. R., C. P. R. Saunders, and J. Hallett, 1983: Laboratory studies of the charging of soft-hail during ice crystal interactions. *Quart. J. Roy. Meteor. Soc.*, **109**, 609–630, <https://doi.org/10.1002/qj.49710946111>.
- Jedlovec, G., E. Berndt, K. Fuell, F. LaFontaine, and N. Elmer, 2018: Analysis and applications of water vapor-derived multispectral composites for geostationary satellites. *AGU 2018 Fall Meeting*, Washington, DC, Amer. Geophys. Union, <https://www.essoar.org/doi/abs/10.1002/essoar.10500213.1>.
- Lensky, I. M., and D. Rosenfeld, 2006: The time-space exchangeability of satellite retrieved relations between cloud top temperature and particle effective radius. *Atmos. Chem. Phys.*, **6**, 2887–2894, <https://doi.org/10.5194/acp-6-2887-2006>.
- , and —, 2008: Clouds-Aerosols-Precipitation Satellite Analysis Tool (CAPSAT). *Atmos. Chem. Phys.*, **8**, 6739–6753, <https://doi.org/10.5194/acp-8-6739-2008>.
- Lindsey, D. T., D. W. Hillger, L. Grasso, J. A. Knaff, and J. F. Dostalek, 2006: GOES climatology and analysis of thunderstorms with enhanced 3.9- μ m reflectivity. *Mon. Wea. Rev.*, **134**, 2342–2353, <https://doi.org/10.1175/MWR3211.1>.
- Maggio, C., and Coauthors, 2005: Lightning-initiation locations as a remote sensing tool of large thunderstorm electric field vectors. *J. Atmos. Oceanic Technol.*, **22**, 1059–1068, <https://doi.org/10.1175/JTECH1750.1>.
- Market, P. S., and Coauthors, 2006: Proximity soundings of thundersnow in the central United States. *J. Geophys. Res.*, **111**, D19208, <https://doi.org/10.1029/2006JD007061>.
- , and A. E. Becker, 2009: A study of lightning flashes attending periods of banded snowfall. *Geophys. Res. Lett.*, **36**, L01809, <https://doi.org/10.1029/2008GL036317>.
- , C. E. Halcomb, and R. L. Ebert, 2002: A climatology of thundersnow events over the contiguous United States. *Wea. Forecasting*, **17**, 1290–1295, [https://doi.org/10.1175/1520-0434\(2002\)017<1290:ACOTEO>2.0.CO;2](https://doi.org/10.1175/1520-0434(2002)017<1290:ACOTEO>2.0.CO;2).
- Mathee, R., and J. R. Mecikalski, 2013: Geostationary infrared methods for detecting lightning-producing cumulonimbus clouds: IR methods for detecting lightning. *J. Geophys. Res. Atmos.*, **118**, 6580–6592, <https://doi.org/10.1002/jgrd.50485>.
- May, R. M., S. C. Arms, P. Marsh, E. Bruning, J. R. Leeman, K. Goebbert, J. E. Thielen, and Z. S. Bruck, 2020: Metpy: A Python Package for Meteorological Data. Unidata, accessed 14 May 2020, <https://github.com/Unidata/MetPy>.
- Mecikalski, J. R., and K. M. Bedka, 2006: Forecasting convective initiation by monitoring the evolution of moving cumulus in daytime GOES imagery. *Mon. Wea. Rev.*, **134**, 49–78, <https://doi.org/10.1175/MWR3062.1>.
- , W. M. MacKenzie, M. Koenig, and S. Muller, 2010a: Cloud-top properties of growing cumulus prior to convective initiation as measured by Meteosat second generation. Part I: Infrared fields. *J. Appl. Meteor. Climatol.*, **49**, 521–534, <https://doi.org/10.1175/2009JAMC2344.1>.
- , —, M. König, and —, 2010b: Cloud-top properties of growing cumulus prior to convective initiation as measured by Meteosat Second Generation. Part II: Use of visible reflectance. *J. Appl. Meteor. Climatol.*, **49**, 2544–2558, <https://doi.org/10.1175/2010JAMC2480.1>.

- , X. Li, L. D. Carey, E. W. McCaul, and T. A. Coleman, 2013: Regional comparison of GOES cloud-top properties and radar characteristics in advance of first-flash lightning initiation. *Mon. Wea. Rev.*, **141**, 55–74, <https://doi.org/10.1175/MWR-D-12-00120.1>.
- Meng, H., J. Dong, R. Ferraro, B. Yan, L. Zhao, C. Kongoli, N.-Y. Wang, and B. Zavadsky, 2017a: A 1DVAR-based snowfall rate retrieval algorithm for passive microwave radiometers: Microwave snowfall rate algorithm. *J. Geophys. Res. Atmos.*, **122**, 6520–6540, <https://doi.org/10.1002/2016JD026325>.
- , —, C. Kongoli, R. Ferraro, B. Yan, K. White, E. Berndt, and B. Zavadsky, 2017b: NESDIS snowfall rate product and its applications. JPSS Science Seminar, 41 pp., http://www.jpss.noaa.gov/assets/pdfs/science_seminars/presentations/Snowfall_Rate_Meng_White_Berndt_2017.pdf.
- Miller, S. D., D. T. Lindsey, C. J. Seaman, and J. E. Solbrig, 2020: GeoColor: A blending technique for satellite imagery. *J. Atmos. Oceanic Technol.*, **37**, 429–448, <https://doi.org/10.1175/JTECH-D-19-0134.1>.
- Moore, J. T., C. E. Graves, S. Ng, and J. L. Smith, 2005: A process-oriented methodology toward understanding the organization of an extensive mesoscale snowband: A diagnostic case study of 4–5 December 1999. *Wea. Forecasting*, **20**, 35–50, <https://doi.org/10.1175/WAF-829.1>.
- Nicosia, D. J., and R. H. Grumm, 1999: Mesoscale band formation in three major Northeastern United States snowstorms. *Wea. Forecasting*, **14**, 346–368, [https://doi.org/10.1175/1520-0434\(1999\)014<0346:MBFTM>2.0.CO;2](https://doi.org/10.1175/1520-0434(1999)014<0346:MBFTM>2.0.CO;2).
- Nieman, S. J., W. P. Menzei, C. M. Hayden, D. Gray, S. T. Wanzong, C. S. Velden, and J. Daniels, 1997: Fully automated cloud-drift winds in NESDIS operations. *Bull. Amer. Meteor. Soc.*, **78**, 1121–1134, [https://doi.org/10.1175/1520-0477\(1997\)078<1121:FACDWI>2.0.CO;2](https://doi.org/10.1175/1520-0477(1997)078<1121:FACDWI>2.0.CO;2).
- Paluch, I. R., and J. D. Sartor, 1973: Thunderstorm electrification by the inductive charging mechanism: I. Particle charges and electric fields. *J. Atmos. Sci.*, **30**, 1166–1173, [https://doi.org/10.1175/1520-0469\(1973\)030<1166:TEBTIC>2.0.CO;2](https://doi.org/10.1175/1520-0469(1973)030<1166:TEBTIC>2.0.CO;2).
- Rauber, R. M., and Coauthors, 2014: Stability and charging characteristics of the comma head region of continental winter cyclones. *J. Atmos. Sci.*, **71**, 1559–1582, <https://doi.org/10.1175/JAS-D-13-0253.1>.
- Reynolds, S. E., M. Brook, and M. Foulks Gourley, 1957: Thunderstorm charge separation. *J. Meteor.*, **14**, 426–436, [https://doi.org/10.1175/1520-0469\(1957\)014<0426:TCS>2.0.CO;2](https://doi.org/10.1175/1520-0469(1957)014<0426:TCS>2.0.CO;2).
- Saunders, C. P. R., H. Bax-Norman, C. Emersic, E. E. Avila, and N. E. Castellano, 2006: Laboratory studies of the effect of cloud conditions on graupel/crystal charge transfer in thunderstorm electrification. *Quart. J. Roy. Meteor. Soc.*, **132**, 2653–2673, <https://doi.org/10.1256/qj.05.218>.
- Schmit, T. J., M. M. Gunshor, W. P. Menzel, J. J. Gurka, J. Li, and A. S. Bachmeier, 2005: Introducing the next-generation advanced baseline imager on GOES-R. *Bull. Amer. Meteor. Soc.*, **86**, 1079–1096, <https://doi.org/10.1175/BAMS-86-8-1079>.
- , P. Griffith, M. M. Gunshor, J. M. Daniels, S. J. Goodman, and W. J. Lebar, 2017: A closer look at the ABI on the GOES-R series. *Bull. Amer. Meteor. Soc.*, **98**, 681–698, <https://doi.org/10.1175/BAMS-D-15-00230.1>.
- Schrom, R. S., and M. R. Kumjian, 2016: Connecting microphysical processes in Colorado winter storms with vertical profiles of radar observations. *J. Appl. Meteor. Climatol.*, **55**, 1771–1787, <https://doi.org/10.1175/JAMC-D-15-0338.1>.
- Schultz, C. J., T. J. Lang, E. C. Bruning, K. M. Calhoun, S. Harikema, and N. Curtis, 2018: Characteristics of lightning within electrified snowfall events using lightning mapping arrays. *J. Geophys. Res. Atmos.*, **123**, 2347–2367, <https://doi.org/10.1002/2017JD027821>.
- , and Coauthors, 2021: Remote sensing of electric fields observed within winter precipitation during the 2020 Investigation of Microphysics and Precipitation for Atlantic Coast-Threatening Snowstorms (IMPACTS) Field Campaign. *J. Geophys. Res. Atmos.*, **126**, e2021JD034704, <https://doi.org/10.1029/2021JD034704>.
- Schultz, D. M., 1999: Lake-effect snowstorms in northern Utah and western New York with and without lightning. *Wea. Forecasting*, **14**, 1023–1031, [https://doi.org/10.1175/1520-0434\(1999\)014<1023:LESINU>2.0.CO;2](https://doi.org/10.1175/1520-0434(1999)014<1023:LESINU>2.0.CO;2).
- Setvák, M., and C. A. Doswell, 1991: The AVHRR channel 3 cloud top reflectivity of convective storms. *Mon. Wea. Rev.*, **119**, 841–847, [https://doi.org/10.1175/1520-0493\(1991\)119<0841:TACCTR>2.0.CO;2](https://doi.org/10.1175/1520-0493(1991)119<0841:TACCTR>2.0.CO;2).
- , R. Rabin, C. A. Doswell, and V. Levizzani, 2003: Satellite observations of convective storm tops in the 1.6, 3.7 and 3.9 μm spectral bands. *Atmos. Res.*, **67–68**, 607–627, [https://doi.org/10.1016/S0169-8095\(03\)00076-0](https://doi.org/10.1016/S0169-8095(03)00076-0).
- Shimizu, A., 2015: Introduction of JMA VLab support site on RGB composite imagery and tentative RGBs. *Sixth Asia/Oceania Meteorological Satellite Users' Conf.*, Tokyo, Japan, Japan Meteorological Agency, 39 pp., http://www.data.jma.go.jp/mscweb/en/aomsuc6_data/oral/st1-07.pdf.
- Sieglaff, J. M., D. C. Hartung, W. F. Feltz, L. M. Counce, and V. Lakshmanan, 2013: A satellite-based convective cloud object tracking and multipurpose data fusion tool with application to developing convection. *J. Atmos. Oceanic Technol.*, **30**, 510–525, <https://doi.org/10.1175/JTECH-D-12-00114.1>.
- Strabala, K. I., S. A. Ackerman, and W. P. Menzel, 1994: Cloud properties inferred from 8–12- μm data. *J. Appl. Meteor.*, **33**, 212–229, [https://doi.org/10.1175/1520-0450\(1994\)033<0212:CPIFD>2.0.CO;2](https://doi.org/10.1175/1520-0450(1994)033<0212:CPIFD>2.0.CO;2).
- Takahashi, T., 1978: Riming electrification as a charge generation mechanism in thunderstorms. *J. Atmos. Sci.*, **35**, 1536–1548, [https://doi.org/10.1175/1520-0469\(1978\)035<1536:REAACG>2.0.CO;2](https://doi.org/10.1175/1520-0469(1978)035<1536:REAACG>2.0.CO;2).
- , T. Tajiri, and Y. Sonoi, 1999: Charges on graupel and snow crystals and the electrical structure of winter thunderstorms. *J. Atmos. Sci.*, **56**, 1561–1578, [https://doi.org/10.1175/1520-0469\(1999\)056<1561:COGASC>2.0.CO;2](https://doi.org/10.1175/1520-0469(1999)056<1561:COGASC>2.0.CO;2).
- , S. Sugimoto, T. Kawano, and K. Suzuki, 2017: Riming electrification in Hokuriku winter clouds and comparison with laboratory observations. *J. Atmos. Sci.*, **74**, 431–447, <https://doi.org/10.1175/JAS-D-16-0154.1>.
- Thiel, K. C., K. M. Calhoun, A. E. Reinhart, and D. R. MacGorman, 2020: GLM and ABI characteristics of severe and convective storms. *J. Geophys. Res. Atmos.*, **125**, e2020JD032858, <https://doi.org/10.1029/2020JD032858>.
- Tilles, J. N., and Coauthors, 2019: Fast negative breakdown in thunderstorms. *Nat. Commun.*, **10**, 1648, <https://doi.org/10.1038/s41467-019-09621-z>.
- Trapp, R. J., D. M. Schultz, A. V. Ryzhkov, and R. L. Holle, 2001: Multiscale structure and evolution of an Oklahoma winter precipitation event. *Mon. Wea. Rev.*, **129**, 486–501, [https://doi.org/10.1175/1520-0493\(2001\)129<0486:MSAEOA>2.0.CO;2](https://doi.org/10.1175/1520-0493(2001)129<0486:MSAEOA>2.0.CO;2).
- Turk, J., J. Vivekanandan, T. Lee, P. Durkee, and K. Nielsen, 1998: Derivation and applications of near-infrared cloud reflectances from GOES-8 and GOES-9. *J. Appl. Meteor.*

- Climatol.*, **37**, 819–831, [https://doi.org/10.1175/1520-0450\(1998\)037<0819:DAAONI>2.0.CO;2](https://doi.org/10.1175/1520-0450(1998)037<0819:DAAONI>2.0.CO;2).
- Velden, C. S., C. M. Hayden, S. J. W. Nieman, W. Paul Menzel, S. Wanzong, and J. S. Goerss, 1997: Upper-tropospheric winds derived from geostationary satellite water vapor observations. *Bull. Amer. Meteor. Soc.*, **78**, 173–196, [https://doi.org/10.1175/1520-0477\(1997\)078<0173:UTWDFG>2.0.CO;2](https://doi.org/10.1175/1520-0477(1997)078<0173:UTWDFG>2.0.CO;2).
- Zavodsky, B., A. Molthan, and M. Folmer, 2013: Multispectral imagery for detecting stratospheric air intrusions associated with mid-latitude cyclones. *J. Oper. Meteor.*, **1**, 71–83, <https://doi.org/10.1519/nwajom.2013.0107>.