

CORRIGENDUM**DAVID J. STEPHENS***Department of Agriculture and Food WA, South Perth, Western Australia, Australia***HARRY VAN LOON***Colorado Research Associates/Northwest Research Associates, Boulder, Colorado***MALCOLM H. LAMOND***Lamond Weather Services, Karrinyup, Western Australia, Australia***NICOLA P. TELCIK***Yarra Valley Water, Mitcham, Victoria, Australia*

Due to an error, an incorrect version of Fig. 8 in Stephens et al. (2007) was published in which the x -axis labels were not accurately placed. The correct version of Fig. 8 is presented below.

REFERENCES

Stephens, D. J., M. J. Meuleners, H. van Loon, M. H. Lamond, and N. P. Telcik, 2007: Differences in atmospheric circulation between the development of weak and strong warm events in the Southern Oscillation. *J. Climate*, **20**, 2191–2209.




FIG. 8. (a) Plot of normalized Rapa Island SLP anomalies since 1951 (black line, left y axis) vs actual Niño-3 SST anomalies (red and blue shading, right y axis). (b) Plot of normalized Niño-3 SST anomalies (red curve) vs normalized Rapa Island SLP anomalies (blue curve) and normalized 850-hPa west Pacific trade winds (purple curve), 1979–2004. All curves have been smoothed using a 5-month running mean.