

Atlantic Hurricane Season of 1974

JOHN R. HOPE

National Hurricane Center, National Weather Service, NOAA, Miami, Fla. 33124

ABSTRACT

A general overview of the 1974 hurricane season in the North Atlantic is presented together with detailed accounts of individual storms.

1. General summary

The relative lull in tropical storm activity that began in the Atlantic in 1970 continued in 1974. Seven tropical cyclones were named during the season, including four hurricanes. The averages for the past 30 years are nine storms including six hurricanes.

Tracks of 1974 North Atlantic tropical and subtropical cyclones are shown in Figs. 1 and 2. A summary of 1974 hurricane statistics is shown in Table 1.

The extensive upper-level westerlies and below normal sea-surface temperatures over large portions of the hurricane-generating areas, which are believed to have been factors inhibiting tropical cyclone development during the past two seasons (Simpson and Hebert, 1973; Hebert and Frank, 1974) were in evidence again, but to a lesser extent. Figures 3 and 4 are charts of the mean vertical shear of the tropospheric horizontal wind (derived by subtracting the 1000 to 600 mb mean from the 600 to 200 mb mean, and from which the strength of the upper westerlies can be inferred), and the mean sea-surface temperature anomalies during the most active part of the hurricane season, 15 August–15 September. Comparison of the 1974 chart with the two previous seasons shows the vertical shear over much of the tropical Atlantic and Caribbean Sea to be less than in 1972 and 1973. In 1974 the 10 kt isotach covers a relatively small area east-northeast of the Leeward Islands, whereas during the 1972 and 1973 seasons it included a much larger area east of the Lesser Antilles as well as most of the Caribbean Sea. Gray (1967) has shown that large vertical shear of the horizontal wind is unfavorable for tropical cyclone development, since the heat released by convection in this situation does not remain in a vertical column to the extent necessary to produce falling surface pressures and the development of a circulation.

In 1974 all the named systems reached tropical storm intensity in areas where the mean vertical shear during 15 August–15 September was less than 10 kt, and less than 5 kt where Alma, Becky, Elaine, Dolly, and Gertrude were named. Gertrude, however, occurred

after 15 September, and Alma had its beginning just before 15 August.

Mean sea-surface temperatures, while continuing to be slightly below normal over much of the tropical Atlantic, were nevertheless generally above Palmén's (1948) threshold value for development. All seven 1974 named storms developed in areas where the mid-season sea-surface temperatures exceeded 27°C.

There were 19 hurricane days during the 1974 season compared to an average of 29 days for the past 20 years. A decline in the Atlantic of the number of hurricane days per season since the mid-1960's is indicated on Fig. 5, in which five-year running averages for the past 20 years are shown.

Hurricane Carmen, the only named tropical cyclone to strike the United States in 1974, was the most intense Atlantic Hurricane since Camille of 1969. Thousands of deaths attributed to Hurricane Fifi in Honduras were the result of rainfall-induced flooding. Although becoming a moderately intense hurricane, Becky was a threat only to ships at sea. Gertrude, the final hurricane of the season, acquired hurricane force winds for only a brief period before reaching the Lesser Antilles.

In addition, four systems during the 1974 season were classified as subtropical storms. All had their origins in initially baroclinic zones, exhibiting some tropical characteristics as latent heat played an increasingly greater role in their development. Subtropical storm statistics are shown in Table 2.

There were no tropical storms in June or July. In June the mid-tropospheric circulation was well amplified, with a trough over east-central United States and a ridge over the western Atlantic (Taubensee, 1974a). Neither the positive height anomaly at 700 mb from the Ohio Valley eastward, nor the negative anomaly from Newfoundland into the western Caribbean, shown by Ballenzweig (1958) to be favorable for development in these respective areas, was present in June. In July, the mean features over the eastern United States and the western Atlantic progressed eastward (Wagner, 1974). The mean trough along the east coast of the




Fig. 1. Tracks of North Atlantic tropical cyclones, 1974.




FIG. 2. Tracks of North Atlantic subtropical cyclones, 1974.

United States, considered favorable for development in the western Caribbean, was present but did not extend far enough into the western Caribbean itself.

All of the 1974 tropical storm developments occurred in August and September. In August the July mid-tropospheric ridge over central United States was replaced by a mean trough, and a strong subtropical ridge developed from the Gulf of Mexico eastward over the Atlantic (Dickson, 1974). The 700 mb height anomaly pattern resembled that shown by Ballenzweig to

favor development near the Lesser Antilles, in which area Alma and Carmen originated. Once formed, the westward path of these storms in the Caribbean was assured by the presence of the strong subtropical ridge.

Since the aforementioned subtropical ridge held and strengthened somewhat in September (Taubensee, 1974b), hurricanes Fifi and Gertrude maintained westward courses. Strong zonal westerlies north of latitude 35°N caused Becky, Elaine, and Dolly to move north-eastward after attaining storm intensity.

TABLE 1. Summary of North Atlantic tropical cyclone statistics, 1974.

No.	Name	Class	Dates	Maximum sustained winds (kt)	Lowest pressure (mb)	U. S. damage (\$ millions)	Deaths
1	Alma	T	Aug. 12-15	55	1007		Trinidad, 2
2	Becky	H	Aug. 26-Sept. 2	100	977		
3	Carmen	H	Aug. 29-Sept. 10	130	928	150	U. S., 1
4	Dolly	T	Sept. 2-5	45	1005		
5	Elaine	T	Sept. 4-13	60	1001		
6	Fifi	H	Sept. 14-22	95	971		*Honduras,
7	Gertrude	H	Sept. 28-Oct. 3	65	999		3,000-10,000

* The Red Cross has confirmed 3,000 fatalities, other estimates range up to 10,000.




FIG. 3. NHC tropospheric mean vertical shear field (600-200 mb upper mean minus the 1000-600 mb lower mean) for the period 15 August-15 September 1974. Dashed lines are isotachs.

Amplification and strengthening of the mean flow and eastward progression of the trough over the western Atlantic in October (Wagner, 1975) permitted sweeps of cool, dry air into the October hurricane-generating areas, inhibiting development there.

2. Individual named storms

a. Tropical Storm Alma, 12-15 August

The initial tropical storm of the season developed unusually far south, forming from a strong ITCZ disturbance in the Eastern Atlantic and reaching tropical storm intensity near 10°N 52°W on 13 August. Only

one storm of record, occurring in 1933, crossed Trinidad farther south than did Alma.

The westward course at low latitudes was maintained by the circulation around a strong subtropical ridge located somewhat to the south of its normal August position.

After the storm center moved westward over Trinidad it continued into northern Venezuela, where its circulation was destroyed by the mountainous terrain on 15 August.

The highest wind reported by a land station was 30 kt with gusts to 42 kt at Piarco Airport on Trinidad, while the maximum associated with the storm during




FIG. 4. Sea-surface temperature anomalies (°C) (solid lines) for the period 15 August-15 September 1974. The dashed line is the 27°C isotherm.

its existence was estimated at 55 kt based on aerial reconnaissance reports. The lowest pressure was 1007 mb on 13 August.

There were two fatalities as a direct result of the storm on Trinidad. On the Isla de Magarita off the Venezuelan coast, 47 persons perished when a passenger aircraft crashed as the leading edge of the rain shield associated with Alma passed over the island. There were reports of considerable property and crop damage on Trinidad.

b. Hurricane Becky, 26 August–2 September

Becky attained tropical storm status west of Bermuda, near 30°N, 68°W, on the morning of 28 August after having acquired a circulation several hundred n mi north of Puerto Rico on the 26th. Hurricane strength was reached on the afternoon of the 28th, followed by further intensification during the next two days as Becky moved slowly northeastward over the open sea and around the periphery of a well-developed surface anticyclone centered just west of the Azores. Maximum strength was observed on the 30th, when reconnaissance reports indicated a minimum central pressure of 977 mb and sustained winds of 100 kt. At this time, 200 mb charts showed a well-developed warm ridge over the hurricane, and satellite loops indicated extensive outflow at the cirrus level.

On 2 September the system lost tropical characteristics after it had accelerated eastward ahead of an approaching frontal trough and merged with a frontal zone northeast of the Azores.

Never a threat to land, Becky was of concern only to shipping, because of its presence astride the main north Atlantic shipping lanes for several days.

c. Hurricane Carmen, 29 August–10 September

Carmen was the most severe Atlantic hurricane since Camille of 1969.

TABLE 2. Summary of North Atlantic subtropical cyclone statistics 1974.

No.	Dates	Maximum sustained winds (kt)	Lowest pressure (mb)
1	June 25–27	60	1000
2	July 16–19	50	1005
3	August 10–16	50	992
4	October 3–8	50	1005

The densely populated areas near Carmen’s two landfalls were spared by last minute changes in the hurricane’s course. First, the hurricane’s center veered away from Belize City, Belize, and brushed Chetumal, Mexico, after crossing the coast in a sparsely populated area. After the weakened Carmen crossed the Yucatan peninsula, it regained strength as it moved northward across the Gulf of Mexico. However, the hurricane turned toward the northwest as it reached the Louisiana coast south of New Orleans, sparing that city from considerable damage.

The forerunner of Carmen was an easterly wave which moved off the African coast on 23 August and became a depression centered about 180 n mi east of Guadeloupe on 29 August. It intensified gradually, under a classic outflow pattern produced by a 200 mb anticyclone that moved westward with the hurricane across the Leeward Islands into the eastern Caribbean. The depression deposited up to five inches of rain in the Virgin and Leeward Islands, and over five inches in some sections of Puerto Rico.

Carmen acquired tropical storm strength south of the Mona Passage on 30 August as it moved westward 80 n mi south of Puerto Rico and Hispaniola, and became a minimal hurricane the following day just south of Jamaica. Intensification had not been rapid since the system entered the Caribbean because the low-level inflow pattern was not well established and much of its circulation was over Hispaniola and eastern Cuba.

The center passed a short distance north of Islas del Cisne (Swan Island) on 1 September, where gale force winds were experienced for a period of four hours, and highest sustained winds reached 50 kt, gusting to 60 kt.

Rapid development ensued as the hurricane center approached the Yucatan peninsula on 2 September with most of its circulation over the very warm waters of the northwest Caribbean Sea while a favorable outflow pattern aloft persisted. Sustained winds reached 130 kt and the pressure fell to 928 mb just before landfall as the Belize radar showed the hurricane’s course shifting to a more northwesterly direction, sparing the cities of Belize and Corozal. Figure 6 is an example of the excellent presentation on the recently-installed Belize radar as Carmen skirted the city.

The center passed a few n mi north of Chetumal, Mexico, where sustained winds reached 118 kt, accompanied by a minimum pressure of 956 mb. Although




FIG. 5. Five-year running average of North Atlantic hurricane days, 1955–1974.




FIG. 6. Radar photograph of Hurricane Carmen from Belize, 0534 GMT 2 September 1974 (courtesy Belize Meteorological Service).

Chetumal likely was under the eye wall of the hurricane, the city was on the storm's weaker side. Winds of 57 kt were reported at San Pedro, Belize, and 87 kt at Corozal.

After moving inland over the Yucatan peninsula, Carmen weakened over land and it was not until 5 September that it regained hurricane strength as it began its northward movement over the Gulf of Mexico




FIG. 7. Radar photograph of Hurricane Carmen from Slidell, La., 0545 GMT 8 September 1974.

in response to surface pressure falls over Texas and rises over Florida and adjacent waters. Reconnaissance reports indicated that sustained winds again reached 130 kt over the northern Gulf of Mexico, while the surface pressure dropped to 937 mb. However, some weakening occurred just before landfall on 8 September as a decrease in echo coverage on land-based radars suggested that cooler, drier air was entering the system. Figure 7 is a photograph of the radar depiction at Slidell, La., at 0545 GMT 8 September, illustrating the absence of echoes in the hurricane's eastern semicircle. Rapid weakening followed landfall on the Louisiana coast. The highest sustained wind measured over Louisiana was 75 kt at Morgan City. Rainfall amounts over land were mostly six inches or less. Reported storm tides ranged up to six feet along the Louisiana coast. Meteorological data associated with the United States landfall of Carmen are shown in Table 3.

Damage caused by Carmen in Puerto Rico was estimated in excess of two million dollars, mainly in flash floods and in a tornado accompanying the developing storm.

In Louisiana crop damage was estimated at 90 million dollars, primarily to sugar cane. Some losses were suffered by offshore oil installations and the shrimp industry. Total damages in Louisiana were about 150 million dollars.

Over 75,000 people, mostly in Louisiana and Mississippi, evacuated low-lying areas ahead of the hurricane.

The only fatality caused directly by Carmen was an electrocution by a fallen power line in Louisiana.

d. Tropical Storm Dolly, 2–5 September

The cloud system that was to evolve into Tropical Storm Dolly was first evident on satellite pictures when it was located 700 n mi east southeast of Bermuda on 30 August. Drifting northwestward for the next few days, the system slowly became better organized and the development of a depression was confirmed on 2 September when a ship reported southwesterly winds south of the developing center. On the following day the storm was named, after a reconnaissance flight reported 45 kt winds and a surface pressure of 1005 mb. Following tropical storm development, Dolly recurved to the northeast around the periphery of the Azores-Bermuda high and ahead of an approaching frontal trough, losing its tropical structure when it merged with the cold front on 5 September. However, the remnants of Dolly, as a frontal wave, produced sustained northerly winds of 33 kt with gusts to 54 kt on Sable Island.

Environmental conditions were not particularly favorable for intensification as a tropical system during the slow transition from depression to a minimal tropical storm, since the system was under a 200 mb trough. However, as the storm moved northward it moved under 200 mb anticyclonic flow. Further tropical de-

TABLE 3. Hurricane Carmen, 29 August–10 September 1974. Meteorological data.

Station	Date	Pressure (mb)		Fastest mile	Wind (kt)			Time (CST)	Tide (ft)		Rainfall (inches)	
		Low	Time (CST)		Time (CST)	Gusts	Highest MSL		Time (CST)	Storm total	Dates	
Florida												
Pensacola WSO	7	1009	1756				43	8/0122			5.09	7-8
Alabama												
Dauphin Island	7	1009	1800			SSW	33	1800			5.28	7-8
Evergreen											11.60	7-9
Mobile WSO	7	1008	1756	ENE 19	1159	NE	33	1404	3.5	1950	7.58	7-8
Mississippi												
Bay St. Louis	8	1006	0230			SSE	42	0235	4.6	0300		
Pascagoula	8	1006	0330			SE	34	0230	4.6	0300		
Louisiana												
Alexandria WSO	8	1000	0902			E	42	1042			1.59	8
Baton Rouge WSO	8	996	0349	NE 28	0030	E	44	0323			0.95	7-8
Bayou Boeuf	8								5.1	0800		
Bayou Lafourche	8								5.9	0400		
Boothville WSO	7	1001	1930			NE	38	0623			7.81	6-7
Cote Blanc Bay	8								4.5	0615		
Grand Isle CG	7	996	1845	48	1900		62	2025	4.0			
Lafayette	8			NE 45	0655	NE	71	0655			3.50	7-8
Lake Charles WSO	8	993	1154			NNW	44	0519			1.84	7-8
Morgan City	8	982					75					
New Orleans WSFO	8	999	0200				49				4.00	7-8
New Orleans WSMO	8	998	0247	NNE 29	7/1732	NE	45	7/2008			3.64	7-8
New Orleans Lakefront Airport	7					NNE	62	7/1639				
Shell Beach (Miss. River Outlet)	8								6.0			
Vermilion Lock (on Intra-Coastal Waterway)	8								4.0			
Wax Lake Outlet (Vermilion Bay)	8								5.3	0815		
Texas												
Galveston WSO	8	1003	1700	NW 28	1604	NW	42	1606	2.4	7/1800	0.13	8
Port Arthur WSO	8	998	1700	N 24	1120	NW	34	1401			0.38	8

velopment was halted as cooler air entered the low-level circulation.

e. Tropical Storm Elaine, 4–13 September.

The convective cloud pattern that developed into Elaine was traced by satellite pictures from the African coast on 30 August to 600 n mi east of the Leeward Islands on 4 September, where a closed circulation was detected by reconnaissance aircraft.

The storm was named during the late afternoon of 9 September, while centered 225 n mi east southeast of Cape Hatteras, on the basis of wind reports from reconnaissance flights. Thereafter, Elaine moved north-eastward, ahead of an approaching frontal trough, losing tropical characteristics as it neared the cold front 350 n mi southeast of Cape Race, Newfoundland, on 13 September. No land areas were threatened by Elaine. Elaine reached storm strength in an area of light vertical shears and under an upper-level anticyclone. As the storm reached higher latitudes, it moved away from the upper anticyclone and under a zone of relatively strong upper westerlies.

The lowest pressure measured in Elaine was 1001 mb on 10 September. Highest sustained winds were esti-

mated to be 60 kt. Some higher winds reported by reconnaissance aircraft are believed to have been transitory and not representative of the strength of the system.

f. Hurricane Fifi, 14–22 September

Hurricane Fifi will be remembered chiefly because of the appalling loss of life in Honduras caused by inland flooding of rivers and streams. The influx of moist air from the Caribbean and the Pacific into the hurricane as it skirted the Honduran coast produced torrential rains up to 20 inches or more in the mountains. In addition to drownings, many perished in mud slides as tilled slopes gave way and the avalanche buried all in its path.

The precursor of Fifi moved off the African coast as an organized convective mass on 8 September. Moving steadily across the tropical Atlantic, it became a well-organized tropical depression on the 15th, centered 60 n mi south of Haiti, and a tropical storm on the 16th just south of Jamaica. Strengthening continued and Fifi became a hurricane 250 n mi east of Islas del Cisne (Swan Island) on 17 September. On the next day the hurricane had acquired its maximum sustained

winds of 95 kt. The lowest pressure of 971 mb was observed on the morning of 19 September; however, slight weakening took place prior to landfall on the coast of southern Belize late that afternoon. Rapid weakening to storm and then to depression strength followed landfall. After crossing Belize and southern Mexico, the system regained strength in the Pacific where it was renamed Orlene. Further strengthening to a hurricane occurred before its final landfall on the west coast of Mexico.

The track of Fifi was controlled by a large high pressure system which persisted over the central Gulf of Mexico throughout the life of the hurricane. Although the 200 mb pattern appeared to become somewhat more favorable as Fifi intensified from depression to hurricane, the upper-level northeasterly flow around the Gulf of Mexico anticyclone inhibited outflow and probably prevented the rapid deepening observed in Carmen and many other hurricanes in the western Caribbean.

Although Honduras received the brunt of Fifi's devastation there were heavy rains in other areas. More than 8 inches fell in two days over southern Jamaica. Heavy rain associated with the hurricane also fell over El Salvador, Guatemala, and southern Belize. Among the highest winds reported at land stations were 115 kt on the island of Guanaja off the north coast of Honduras, 85 kt on *Islas del Cisne* (Swan Island), and 85 kt with gusts to 100 kt in southern Belize.

The actual number of deaths caused by Fifi may never be determined. Estimates range from 3000 to 10000. The Red Cross has confirmed by actual count the lower figure, which is high enough to place Fifi among the most disastrous hurricanes ever to strike in the western hemisphere.

g. Hurricane Gertrude, 28 September–3 October

The seventh and last named storm of the season developed from an Intertropical Convergence Zone disturbance which moved off the African coast on 22 September. By the 26th, satellite pictures showed increasing organization, and a depression had formed by early morning of the 28th. Reconnaissance flights reported hurricane force winds that afternoon.

Gertrude never developed the characteristics of a well organized hurricane. No wall cloud ever formed. The lowest pressure was 999 mb on 28 September. It is concluded that the hurricane force winds observed occurred during short-period accelerations.

Weakening began on the 30th, when satellite pictures showed the low-level circulation center separating from the main convective cloud mass as the storm slowly approached an upper trough and increasing vertical shears. The system was downgraded to a tropical depression on 2 October, just prior to reaching the Windward Islands, and its circulation disappeared over the southeast Caribbean on the following day.

Locally heavy showers, well east of the remnants of Gertrude, continued over the Windward Islands through 4 October.

There are no reports of substantial damage or casualties attributed to Gertrude.

3. Subtropical storms

a. Subtropical storm # 1 24–27 June

The development of this subtropical storm in the eastern Gulf of Mexico followed by several days the formation of a tropical depression off the Mexican coast near Vera Cruz on 22 June

By the evening of 24 June the convective system associated with the southwestern Gulf of Mexico tropical depression had weakened while convective cloudiness had increased and a new low center had formed in the eastern Gulf of Mexico. The new system moved northeastward at 30–35 kt during the night of 24–25 June and crossed the Florida peninsula during the morning of the 25th. Sustained winds of 30–40 kt, with gusts to 45–55 kt, were reported in the Fort Myers-Naples area as the system approached. Tides 2–4 feet above normal occurred from Everglades City to the Tampa Bay area.

At midday on the 25th, after the storm moved over the Atlantic east of Florida, reconnaissance aircraft and ships found a band of 45–60 kt winds about 150 n mi wide to the east of the storm center. As it moved northeastward during the 25th, the system weakened as it became frontal in character, and became difficult to identify by late evening.

Total rainfall from the storm and the tropical depression, which crossed the coast two days later, amounted to 20 inches in the Tampa Bay area and 10 inches or more over much of west central Florida.

There were three deaths by drowning in Florida. Damages caused by tidal and heavy rainfall flooding has been estimated at \$10,000,000 in western Florida.

b. Subtropical storm # 2, 16–19 July

Satellite pictures on 15 July suggested that a weak circulation was developing northeast of the Bahamas in an area of convective cloudiness that had been associated with a quasistationary front.

The circulation, as indicated on satellite pictures, became increasingly better organized as it moved northeastward through 18 July; however, the area covered by the circulation remained small through this period and there was no evidence from ship reports that supported satellite-picture evidence of a significant weather disturbance until the ship *Export Adventurer* encountered winds of 47 kt and a pressure of 1006 mb at 2100 GMT 18 July.

The ship's barogram and hourly weather reports between 2000 GMT 18 July and 0400 GMT 19 July

indicated a closed circulation, and that the ship's barometer fell 18 mb in 6 hours.

The low became absorbed in the circulation of a large extratropical low-pressure system off Newfoundland by 20 July, as it continued rapidly northeastward.

c. Subtropical storm # 3, 10–16 August

The convective area which evolved into this subtropical storm was associated on 10 August with a wave located 300 n mi southeast of southern New England. The wave had developed along a stationary front extending northeastward from near Cape Hatteras over the North Atlantic.

As the system drifted northeastward, slowly deepening, the air mass along its northern periphery gradually became warmed; 200 mb ridging, and developing light vertical shears, suggested that the low was acquiring some tropical structure.

At 0000 GMT 11 August, ship 4YH reported sustained winds of 40 kt as the low center passed about 200 n mi east of the ship. During the next two days, several ships reported winds in excess of gale force as the low deepened to an estimated 992 mb. The low passed over Cape Race, Newfoundland, on 15 August, after satellite pictures showed the vortex was less well defined; the remaining cloudiness assumed a configuration typical of a frontal band.

d. Subtropical storm # 4, 3–8 October

There were indications of a low center forming over extreme eastern Cuba during the afternoon of 3 October as a cold front that had been moving southeastward across eastern Cuba was becoming quasistationary. An extensive area of cloudiness and rain persisted behind the front. A large high pressure cell was centered over the middle Atlantic states, and the strong pressure gradient between the high center and the front was already producing northeasterly winds up to 30 kt over the eastern Gulf of Mexico, the Florida Straits, and over the Bahamas and adjacent waters.

By the afternoon of 5 October, as the low center moved into the central Bahamas, a number of ships reported winds in excess of gale force, as did land stations in the Bahamas. Although the storm began to weaken as it moved northeastward away from the Florida coast during the afternoon of 7 October, ships

reported winds up to 40 kt on 7–8 October. By the afternoon of 8 October, the weakening storm had merged with a cold front 350 n mi east of Cape Hatteras.

The main effects of the storm on land areas were tidal flooding and some beach erosion along the Florida east coast, caused by the strong and persistent northeast winds, and some local flooding as a result of heavy rainfall. Tides ranged up to three feet above normal along portions of the Florida east coast. Rainfall amounts over a three-day period included 10 inches at Cocoa, 11.5 inches at Deerfield Beach, and 14 inches at Boca Raton.

Total damage along the Florida east coast is estimated at less than one million dollars.

Acknowledgments. Portions of the individual storm accounts were extracted from preliminary reports prepared by Messrs. G. B. Clark, P. J. Hebert, M. B. Lawrence, and J. M. Pelissier at the National Hurricane Center, and by C. Conner and José Colon of the New Orleans and San Juan Hurricane Warning Offices, respectively. D. Martin and R. Powell prepared the graphics. L. Wilson typed the manuscript.

REFERENCES

- Ballenzweig, E. M., 1958: Formation of tropical storms related to anomalies of the long period circulation. National Hurricane Research Report No. 21, U. S. Weather Bureau, 16 pp.
- Dickson, R. R., 1974: Weather and circulation of August 1974—Relief from heat and drought in the central United States. *Mon. Wea. Rev.*, **102**, 807–811.
- Gray, W. M., 1967: Global view of the origin of tropical disturbance and storms. Atmospheric Science Paper No. 114, Colorado State University, 105 pp.
- Hebert, P. J., and N. L. Frank, 1974: Atlantic hurricane season of 1973. *Mon. Wea. Rev.*, **102**, 280–289.
- Palmén, E. H., 1948: On the formations and structure of tropical hurricanes. *Geophysica*, **3**, 26–38.
- Simpson, R. H., and P. J. Hebert, 1973: Atlantic hurricane season of 1972. *Mon. Wea. Rev.*, **101**, 323–333.
- Taubensee, R. E., 1974a: Weather and circulation of June, 1974—Increasing drought in the Southwest. *Mon. Wea. Rev.*, **102**, 662–666.
- , 1974b: Weather and circulation of September, 1974—Record September cold in the central United States. *Mon. Wea. Rev.*, **102**, 871–876.
- Wagner, A. J., 1974: Weather and circulation of July, 1974—Heat wave and drought over the middle third of the country. *Mon. Wea. Rev.*, **102**, 736–742.
- , 1975: Weather and circulation of October, 1974—Record early fall cold over the eastern half of the United States. *Mon. Wea. Rev.*, **103**, 78–85.