

CORRIGENDUM

FLORIAN SÉVELLEC

*Ocean and Earth Science, National Oceanography Centre Southampton, University of Southampton,
Southampton, United Kingdom*

JOËL J.-M. HIRSCHI AND ADAM T. BLAKER

National Oceanography Centre, Southampton, Southampton, United Kingdom

Two production errors occurred in [Sévellec et al. \(2013\)](#). In both [Figs. 5](#) and [6](#), titles above the figures panels were incorrectly removed, leaving the two figures without units (hour and m s^{-1} , respectively). Those figures appear below as they were meant to appear originally.

The staff of the *Journal of Physical Oceanography* regrets any inconvenience these errors may have caused.

REFERENCES

- Blaker, A. T., J. J.-M. Hirschi, B. Sinha, B. de Cuevas, S. Alderson, A. Coward, and G. Madec, 2012: Large near-inertial oscillations of the Atlantic meridional overturning circulation. *Ocean Modell.*, **42**, 50–56.
- Sévellec, F., J. J.-M. Hirschi, and A. T. Blaker, 2013: On the near-inertial resonance of the Atlantic meridional overturning circulation. *J. Phys. Oceanogr.*, **43**, 2661–2672, doi:[10.1175/JPO-D-13-092.1](https://doi.org/10.1175/JPO-D-13-092.1).

Corresponding author address: Florian Sévellec, Ocean and Earth Science, University of Southampton, Waterfront Campus, European Way, Southampton, SO14 3ZH, United Kingdom.
E-mail: florian.sevellec@noc.soton.ac.uk




FIG. 5. Period (hour) of the max amplitude response of the vertical shear for the geo- (black lines) and ageostrophic (gray lines) terms following (11) and (16), respectively. Solid lines represent the result for $\epsilon = 2 \times 10^{-5} \text{ s}^{-1}$, the gray lines in Figs. 3b and 4b, respectively. Dashed lines represent the result for $\epsilon = 5 \times 10^{-5} \text{ s}^{-1}$, the gray lines in Figs. 3c and 4c, respectively. The thin black line indicates the inertial period. The crosses denote the AMOC variability period from the $1/4^\circ$ OGCM experiments by Blaker et al. (2012).




FIG. 6. Phase and group velocities [c_p (solid lines) and c_g (dashed lines), respectively] of the Poincaré waves for a 5° wavelength as a function of latitude (m s^{-1}). These results follow the analytical expression of (22), where a negative value of these curves is also a possible solution. The crosses denote the phase velocities from the $1/4^\circ$ OGCM experiments by Blaker et al. (2012).